

Entanglement, steering, and separability in Compton-scattered annihilation photons

Peter Caradonna^{1,*} and Kenji Shimazoe^{1,2}

¹*Department of Nuclear Engineering and Management, The University of Tokyo, Bunkyo-ku, Tokyo, Japan*

²*Department of Bioengineering, The University of Tokyo, Bunkyo-ku, Tokyo, Japan*



(Received 8 May 2025; accepted 22 August 2025; published 5 September 2025)

This work analyzes quantum correlations of photon pairs initially prepared in a Bell state, where one photon undergoes Compton scattering from a stationary, unpolarized electron. The evolved two-photon density matrix is analytically derived and spectrally decomposed. Concurrence and the Clauser-Horne-Shimony-Holt (CHSH) inequality evaluations identify angular regions demonstrating quantum steering correlations, entanglement without steering, and complete separability. The Peres conjecture is applied to interpret regions of entanglement lacking steering, emphasizing their limited practicality for quantum-information tasks. These results provide details potentially useful for employing Compton-scattered annihilation photons in various quantum protocols.

DOI: [10.1103/j7p7-ckp9](https://doi.org/10.1103/j7p7-ckp9)

I. INTRODUCTION

The 511-keV emission line, resulting from electron-positron annihilation in the singlet state, is a focus of study in astrophysics, particularly those emitted from the Galactic center region [1]. Observational campaigns using balloon-borne detectors and satellite missions have mapped this emission line across the Galaxy, revealing complexities in its spatial distribution, spectral properties, and temporal stability. Despite this work, uncertainties persist regarding the astrophysical origins of positrons, their interactions with the interstellar medium, and how photon polarization and coherence might encode information about local magnetic fields, matter density distributions, and energetic processes. Importantly, existing research generally does not account for the quantum entanglement inherently present between annihilation photon pairs.

In other studies by Berera [2] and Berera and Calderón-Figueroa [3], it was demonstrated that photons could maintain quantum coherence and entanglement across interstellar distances. While these studies primarily addressed environmental decoherence and highlighted quantum teleportation [4] as a method for quantum communication and potential detection of intelligent extraterrestrial signals, they did not explicitly analyze intrinsic Compton scattering, a dominant interaction channel at the 511-keV photon energy critical for understanding quantum correlations at these energies in astrophysical contexts.

Annihilation photons are also practically utilized in materials science [5,6] and positron emission tomography (PET) [7,8], which is prominently used for early cancer detection. Compton scattering frequently occurs in PET imaging [9,10],

negatively affecting accuracy, sensitivity, and image contrast [11–13]. Quantum entanglement has long been recognized to influence scattering distributions [14–16], stimulating interest in exploiting entanglement via Compton polarimetry to enhance PET imaging. A recently developed theoretical model [10] of photon decoherence during scattering has been supported by preliminary experimental validation [17,18], encouraging further investigation into entanglement-based methods for improving PET imaging.

Building on the formalism from [16], the present study develops a quantum mechanical approach to photon-electron interactions, typically described by 4×4 Mueller matrices [19]. This approach employs a spin-averaged Compton scattering matrix, tracing over electron spin states and introducing mixedness into the photon subsystem due to unobserved degrees of freedom within a closed quantum system undergoing unitary evolution [20]. By addressing intrinsic Compton scattering explicitly, this work complements earlier studies of environmental decoherence. Analysis of entanglement dynamics through concurrence and quantum steering [via violations of the Clauser-Horne-Shimony-Holt (CHSH) inequality] reveals how these quantum correlations distinctly respond to scattering processes. Results from this study may inform gamma-ray astrophysics, indicate new possibilities for astronomical-scale quantum communication, and provide PET research with a detailed quantum description of annihilation photons, enabling the advancement of entanglement-based imaging techniques. Additionally, these findings carry implications for high-energy quantum optics and the development of advanced coincidence-based Compton polarimetry architectures [21–28].

II. SPECTRAL DECOMPOSITION OF THE EVOLVED STATE

This section extends the analysis of Compton scattering presented in Sec. V D of Ref. [16], transforming the evolved Stokes density matrix into a quantum mechanical description of the scattering process. Detailed derivations, including

*Contact author: petercaradonna65@gmail.com

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.




FIG. 1. Compton scattering of a maximally entangled incident (emitted) annihilation beam: A photon from subsystem 2, initially propagating along the z_2 axis with energy $E_0 = 1$ (in natural units) and momentum $\vec{k}_0$, scatters off a stationary electron. The electron recoils with energy ϵ and momentum $\vec{p}$, emerging at a polar angle φ relative to the momentum $\vec{k}_0$ of the incident photon. The scattered photon (signal photon) is deflected into the z'_2 direction, with kinetic energy $E(\theta)$ and momentum $\vec{k}$, where θ and ϕ are the scattering and azimuthal angles, respectively. Each signal photon is correlated with an idler photon, both originating from the same electron-positron annihilation event. For detailed kinematic definitions, see Ref. [16].

definitions of momenta and energies of incident photons, scattered photons, and recoiling electrons, are provided therein.

Consider an initial two-photon system comprising annihilation photons prepared in the maximally entangled Bell state $|\psi_c^-\rangle$ (as denoted in Ref. [16]), given by

$$|\psi_c^-\rangle = \frac{1}{\sqrt{2}}|R, R\rangle - \frac{1}{\sqrt{2}}|L, L\rangle, \quad (1)$$

where the computational basis states $|R\rangle$ and $|L\rangle$ correspond to right- and left-handed circular polarizations, respectively.

Let the density matrix of Eq. (1) be denoted as $\rho = |\psi_{-}^{-}\rangle\langle\psi_{-}^{-}|$. The evolved Stokes density matrix $S'(\rho)$ describes the polarization state of a photon subensemble defined by measuring signal photons scattered in a specified direction and detected in coincidence with corresponding idler photons, as illustrated in Fig. 1. This conditional (postselected) measurement defines a polarization subensemble, as established previously in Ref. [16]. The matrix is given by

$$S'(\rho) = \frac{r_o^2}{2} \left(\frac{E(\theta)}{E_0} \right)^2 \times \begin{bmatrix} t_{11} & t_{12} & 0 & 0 \\ -t_{12} \cos 2\phi - (2 - t_{12}) \cos 2\phi & t_{33} \sin 2\phi & 0 & 0 \\ t_{12} \sin 2\phi & (2 - t_{12}) \sin 2\phi & t_{33} \cos 2\phi & 0 \\ 0 & 0 & 0 & t_{44} \end{bmatrix}, \quad (2)$$

where the explicit form of the t_{ij} for the spin-average case of the electron has been obtained by the method used by Wightman [29].

To obtain the quantum mechanical density matrix ρ' in Hilbert space from the Stokes density matrix $S'(\rho)$, the transformation of Abouraddy *et al.* [30] is applied, such that

$$\rho' = \frac{\sum_{i,j=0}^3 S'_{ij}(\rho) \sigma_i \otimes \sigma_j}{\text{Tr}[\sum_{i,j=0}^3 S'_{ij}(\rho) \sigma_i \otimes \sigma_j]},$$

where $S'_{ij}(\rho)$ ($i = 0, 1, 2, 3$) denote the matrix elements of $S'(\rho)$ and σ_i are the standard Pauli matrices, with $\sigma_0 = \mathbb{I}_2$ representing the 2×2 identity matrix. By construction, ρ' is Hermitian, trace-normalized, and positive semi-definite.

Symbolically, ρ' can be expressed as

$$\rho' = \begin{bmatrix} \rho'_{11}(\theta) & \rho'_{12}(\theta) & \rho'_{13}(\theta, \phi) & \rho'_{14}(\theta, \phi) \\ \rho'^{*}_{12}(\theta) & \rho'_{22}(\theta) & \rho'_{23}(\theta, \phi) & \rho'_{24}(\theta, \phi) \\ \rho'_{13}(\theta, \phi) & \rho'^{*}_{23}(\theta, \phi) & \rho'_{33}(\theta) & \rho'_{34}(\theta, \phi) \\ \rho'^{*}_{14}(\theta, \phi) & \rho'^{*}_{23}(\theta, \phi) & \rho'^{*}_{24}(\theta, \phi) & \rho'_{44}(\theta) \end{bmatrix}. \quad (3)$$

Diagonal elements depend solely on θ , with the functional forms of all the matrix elements provided in Appendix A.

To facilitate the analysis of ρ' given in Eq. (3), introduce the following orthonormal Bell-like states, incorporating an azimuthal phase factor $e^{-2i\phi}$

$$\begin{aligned}
|\Phi_1^-\rangle &= \frac{1}{\sqrt{2}}|R, R\rangle - \frac{e^{-2i\phi}}{\sqrt{2}}|L, L\rangle, \\
|\Psi_2^-\rangle &= \frac{1}{\sqrt{2}}|R, L\rangle - \frac{e^{-2i\phi}}{\sqrt{2}}|L, R\rangle, \\
|\Phi_3^+\rangle &= \frac{1}{\sqrt{2}}|R, R\rangle + \frac{e^{-2i\phi}}{\sqrt{2}}|L, L\rangle, \\
|\Psi_4^+\rangle &= \frac{1}{\sqrt{2}}|R, L\rangle + \frac{e^{-2i\phi}}{\sqrt{2}}|L, R\rangle.
\end{aligned} \tag{4}$$

The states given in Eq. (4) constitute a maximally entangled orthonormal basis that are dependent on the scattering-plane orientation via the azimuthal phase factor.

The spectral decomposition of ρ' naturally splits into two distinct sets of eigenstates, which can be expressed as

$$\rho' = \sum_{i=1}^2 P_i(\theta) |\psi_i\rangle \langle \psi_i| + \sum_{j=3}^4 P_j(\theta) |\chi_j\rangle \langle \chi_j|. \quad (5)$$

The eigenstates $|\psi_1\rangle$ and $|\psi_2\rangle$ represent superpositions of the Bell-like states $|\Phi_1^-\rangle$ and $|\Psi_2^-\rangle$, defined by

$$|\psi_1\rangle = a_1(\theta)[|\Phi_1^-\rangle + c_1(\theta)|\Psi_2^-\rangle], \quad (6a)$$

$$|\psi_2\rangle = a_2(\theta)[|\Phi_1^-\rangle - c_2(\theta)|\Psi_2^-\rangle]. \quad (6b)$$

The normalization factors $a_1(\theta)$, $a_2(\theta)$ and relative phase factors $c_1(\theta)$, $c_2(\theta)$ depend only on the scattering angle θ , whose functional forms are provided in Appendix B.

In contrast, the remaining two eigenstates directly correspond to the Bell-like states and are given by

$$|\chi_3\rangle \equiv |\Phi_3^+\rangle, \quad |\chi_4\rangle \equiv |\Psi_4^+\rangle. \quad (6c)$$

The eigenvalues $P_i(\theta)$ appearing in Eq. (5) represent the measurement probabilities corresponding to the eigenstates defined in Eqs. (6). These eigenstates form a complete




FIG. 2. (a) Eigenstate probabilities $P_i(\theta)$ obtained by projecting the evolved two-photon system onto the eigenbasis defined in Eqs. (6). The state $|\psi_1\rangle$ dominates throughout the entire angular range. (b), (c) Decomposition of eigenstates $|\psi_1\rangle$ and $|\psi_2\rangle$ into intermediate Bell-like basis states $|\Phi_1^-\rangle$ and $|\Psi_2^-\rangle$. (d) Purity, $\text{Tr}[\rho^2]$, of the evolved density matrix as a function of scattering angle θ , showing maximal mixedness at $\theta = 105.6^\circ$.

orthonormal basis, fulfilling the normalization and completeness conditions expressed as

$$\sum_{i=1}^4 P_i(\theta) = 1, \quad \sum_{i=1}^2 |\psi_i\rangle\langle\psi_i| + \sum_{j=3}^4 |\chi_j\rangle\langle\chi_j| = \mathbb{I}_4,$$

where $\mathbb{I}_4$ is the 4×4 identity matrix. Furthermore, the eigenstates satisfy the orthonormal relations such that

$$\begin{aligned} \langle\psi_i|\psi_j\rangle &= \delta_{ij} \quad (i, j = 1, 2), \\ \langle\chi_i|\chi_j\rangle &= \delta_{ij} \quad (i, j = 3, 4), \\ \langle\psi_i|\chi_j\rangle &= 0, \quad (i = 1, 2) \quad (j = 3, 4). \end{aligned}$$

Figure 2(a) illustrates the dependence of the eigenvalues $P_i(\theta)$ on the Compton scattering angle θ .

Furthermore, the probabilities of finding eigenstates $|\psi_1\rangle$ and $|\psi_2\rangle$ in terms of the Bell-like basis states $|\Phi_1^-\rangle$ and $|\Psi_2^-\rangle$ are given by

$$\begin{aligned} p_1(\theta; \Phi_1^-) &= \frac{P_1(\theta)}{1 + c_1^2(\theta)}, & p_1(\theta; \Psi_2^-) &= \frac{P_1(\theta)c_1^2(\theta)}{1 + c_1^2(\theta)}, \\ p_2(\theta; \Phi_1^-) &= \frac{P_2(\theta)}{1 + c_2^2(\theta)}, & p_2(\theta; \Psi_2^-) &= \frac{P_2(\theta)c_2^2(\theta)}{1 + c_2^2(\theta)}, \end{aligned}$$

subject to normalization with respect to the eigenvalues $P_1(\theta)$ and $P_2(\theta)$, such that

$$\begin{aligned} P_1(\theta) &= p_1(\theta; \Phi_1^-) + p_1(\theta; \Psi_2^-), \\ P_2(\theta) &= p_2(\theta; \Phi_1^-) + p_2(\theta; \Psi_2^-). \end{aligned}$$

These relative probabilities and the corresponding eigenvalues (defined in Appendix B) are graphically illustrated in Figs. 2(b) and 2(c).

The spectral decomposition into two distinct terms in Eq. (5) can now be clearly understood. The first term on the right-hand side contains eigenstates $|\psi_1\rangle$ and $|\psi_2\rangle$ whose entanglement continuously varies with the scattering angle θ , ranging from maximally entangled to completely separable. In particular, at $\theta = 90^\circ$, states $|\psi_1\rangle$ and $|\psi_2\rangle$ become fully separable and take the form

$$\begin{aligned} |\psi_1(90^\circ)\rangle &= \frac{1}{2}(|R\rangle + |L\rangle) \otimes (|R\rangle - e^{-2i\phi}|L\rangle), \\ |\psi_2(90^\circ)\rangle &= \frac{1}{2}(|R\rangle + e^{-2i\phi}|L\rangle) \otimes (|R\rangle - |L\rangle). \end{aligned}$$

In contrast, the second term in Eq. (5) involves eigenstates $|\Phi_3^+\rangle$ and $|\Psi_4^+\rangle$ whose maximal entanglement remains invariant across all angles θ .

At forward scattering ($\theta \rightarrow 0^\circ$), a single pure maximally entangled eigenstate emerges where

$$\rho' \rightarrow |\Phi_1^-\rangle\langle\Phi_1^-|, \quad \theta \rightarrow 0^\circ.$$

This implies that signal photons scattering from the electron at $\theta = 0^\circ$ yield a photon state identical to the initial state $|\psi_c^-\rangle$ [Eq. (1)], apart from a relative phase factor $e^{-2i\phi}$, which depends solely by the orientation of the scattering plane.

At backward scattering ($\theta \rightarrow 180^\circ$), the state becomes a classical mixture of Bell-like states such that

$$\rho' \rightarrow 0.8|\Psi_2^-\rangle\langle\Psi_2^-| + 0.2|\Psi_4^+\rangle\langle\Psi_4^+|, \quad \theta \rightarrow 180^\circ.$$

Finally, the reduction in purity for scattering angles between $0 \leq \theta \leq 180^\circ$ is quantified by the purity measure $\text{Tr}(\rho'^2)$, depicted in Fig. 2(d). The purity minimum near $\theta \approx 105^\circ$ follows from the common denominator of the eigenstate probabilities $P_i(\theta)$ (refer to Appendix B) and the sine numerators that drive $P_2 \propto \sin^2(\frac{\theta}{2})\sin^2\theta$ and $P_4 \propto \sin^6(\frac{\theta}{2})$ to nonnegligible values, which dilutes the leading eigenvalue and reduces $\text{Tr}[\rho'^2]$ at that angle.

III. CONCURRENCE OF COMPTON SCATTERING

The degree of entanglement in the evolved two-photon state ρ' , following Compton scattering involving subsystem 2, is quantified using the concurrence $\mathcal{C}(\rho')$. This metric is bounded below by 0 (no entanglement) and above by 1 (maximal entanglement), and is defined as [31,32]

$$\mathcal{C}(\rho') = \max(0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4),$$

where $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4$ are the eigenvalues of the Hermitian matrix

$$R = \sqrt{\sqrt{\rho'} \tilde{\rho}' \sqrt{\rho'}}.$$

The spin-flipped state $\tilde{\rho}'$ is given by

$$\tilde{\rho}' = \sigma_2 \otimes \sigma_2 \rho'^* \sigma_2 \otimes \sigma_2,$$

where ρ'^* denotes the complex conjugate of ρ' in the magic basis. The conjugation is defined as

$$\rho'^* = \sum_{i,j=1}^4 |e_i\rangle\langle e_j| \rho' |e_i\rangle\langle e_j|,$$

with the basis vectors

$$\begin{aligned} |e_1\rangle &= \frac{1}{\sqrt{2}}(|R, R\rangle + |L, L\rangle), \\ |e_2\rangle &= \frac{i}{\sqrt{2}}(|R, R\rangle - |L, L\rangle), \\ |e_3\rangle &= \frac{i}{\sqrt{2}}(|R, L\rangle + |L, R\rangle), \\ |e_4\rangle &= \frac{1}{\sqrt{2}}(|R, L\rangle - |L, R\rangle). \end{aligned}$$

Calculations demonstrate that concurrence depends solely on the polar scattering angles θ , and is independent of the azimuthal angle ϕ . Thus, as illustrated in Fig. 3(a), concurrence is presented as a function of the scattering angle θ and the electron recoil angle φ , following the geometry defined in Fig. 1. Using an incomplete basis set, Tkachev *et al.* [17] concluded that “The [annihilation photon] state never collapses to a separable one.” This conclusion can be directly tested by calculating the concurrence using a complete basis set, as illustrated in Fig. 3(a). The concurrence distinctly vanishes within the angular interval $85.5^\circ \leq \theta \leq 96^\circ$. Although entangled eigenvectors are present in the spectral decomposition, the overall state becomes separable, consistent with the zero concurrence observed. Consequently, the photon pair in this angular range exhibits no entanglement and can be accurately described as a statistical mixture of product states. Outside this region, when $\mathcal{C}(\rho') > 0$, residual entanglement remains.

The concurrence distribution presented in Fig. 3(a) is projected onto the surface of a unit sphere defined in the laboratory coordinate frame x, y, z , as illustrated in Fig. 3(b), with additional viewpoints provided in Fig. 3(c). This representation provides a directional map of entanglement with respect to the signal photon. The point where the scattered photon trajectory intersects the sphere encodes the concurrence value, allowing for a directional interpretation of how the interaction affects entanglement. Similarly, intersections of the scattered electron trajectory can be used to infer the correlations between the idler and signal photons.

IV. QUANTUM STEERING ANALYSIS VIA CHSH VIOLATION AND PERES CONJECTURE

Quantum steering refers to the nonlocal ability of measurements performed locally on one subsystem (the signal) of an entangled photon pair to influence the inferred quantum state of another spatially separated subsystem (the idler). The CHSH inequality [33], a well-established criterion for testing nonlocal correlations, is leveraged to determine whether two-photon states generated by Compton scattering exhibit quantum steering. The application of this inequality clearly delineates polarization correlations into distinct regimes according to correlation strength.

To verify consistency with established formulations, the method of Horodecki *et al.* [34,35] was also applied. In this approach, the correlation tensor $T_{\rho'}$ associated with the




FIG. 3. (a) Concurrence $\mathcal{C}(\rho')$ as a function of scattering angle θ and electron recoil angle φ . The concurrence vanishes within the angular interval $85.5^\circ \leq \theta \leq 96^\circ$, signifying that photon pairs become separable in this region. (b) Representation of concurrence values projected onto a unit sphere in the $\{x_2, y_2, z_2\}$ coordinate frame, illustrating how entanglement persists directionally after scattering. (c) Concurrence distributions viewed from multiple vantage points: an angled view providing depth perception, along with direct views from the $+z$, $+y$, and $-z$ axes, highlighting angular symmetry and areas of entanglement loss.

evolved two-photon density matrix ρ' , Eq. (3), is defined as

$$T_{\rho'} = \begin{bmatrix} \tau_{11} & \tau_{12} & \tau_{13} \\ \tau_{21} & \tau_{22} & \tau_{23} \\ \tau_{31} & \tau_{32} & \tau_{33} \end{bmatrix} \quad (7)$$

with individual elements obtained through

$$\tau_{ij} = \text{Tr}[\rho'(\sigma_i \otimes \sigma_j)], \quad i, j = 1, 2, 3, \quad (8)$$

where σ_i denote the Pauli matrices.

Evaluating the matrix elements in Eq. (7), using Eq. (8), yields

$$T_{\rho'} = \frac{1}{t_{11}} \begin{bmatrix} -(2 - t_{12}) \cos 2\phi & t_{33} \sin 2\phi & 0 \\ (2 - t_{12}) \sin 2\phi & t_{33} \cos 2\phi & 0 \\ 0 & 0 & t_{44} \end{bmatrix}. \quad (9)$$

A direct comparison between the entries τ_{ij} and the elements $S'_{ij}(\rho)$ of the Stokes density matrix $S'(\rho)$, Eq. (2), establishes

$$\tau_{ij} = \frac{S'_{ij}(\rho)}{S'_{00}(\rho)}, \quad i, j = 1, 2, 3,$$

with the labeling convention given in Eq. (15) of Ref. [16].

Therefore, the tensor $T_{\rho'}$ can be directly extracted from the Stokes density matrix $S'(\rho)$ by removing the first column and row, which individually represent the conditional Stokes vectors of the idler and signal photons, respectively, (see Ref. [16]), and then normalizing the remaining 3×3 submatrix by the top-left diagonal entry $S'_{00}(\rho)$.

Next, using Eq. (9), a symmetric correlation matrix is constructed as

$$T_{\rho'}^\top T_{\rho'}, \quad (10)$$

where $T_{\rho'}^\top$ is the transpose of $T_{\rho'}$. The eigenvalues $\lambda_1, \lambda_2, \lambda_3$ of the symmetric matrix given in Eq. (10) are computed and sorted in descending order as

$$\lambda_1 \geq \lambda_2 \geq \lambda_3.$$

From these eigenvalues, the steering measure $M(\rho') = \lambda_1 + \lambda_2$ is defined as the sum of the two largest eigenvalues,




FIG. 4. (a) Quantity $M(\rho')$, defined as the sum of the two largest eigenvalues of the correlation matrix $T_{\rho'}^T T_{\rho'}$, plotted as a function of scattering angle θ . Regions with $M(\rho') > 1$ (shaded magenta) identify angular ranges ($0^\circ \leq \theta \leq 69.3^\circ$ and $131.6^\circ \leq \theta \leq 180^\circ$) exhibiting quantum steering correlations and CHSH inequality violation. The intermediate region ($69.3^\circ < \theta < 131.6^\circ$, shaded yellow) marks the absence of quantum steering correlations ($M(\rho') \leq 1$). Dashed lines indicate the classical-quantum boundary at $M(\rho') = 1$ and critical transition angles at $\theta = 69.3^\circ$ and $\theta = 131.6^\circ$. (b) Directional visualization of steering correlations through projection onto a unit sphere defined in the $\{x_2, y_2, z_2\}$ reference system, illustrating regions where quantum steering remains robust postscreening. (c) Quantum steering correlation maps observed from an angled perspective and three additional viewpoints (+z, +y, and -z axes), emphasizing angular symmetry and delineating regions supporting CHSH inequality violations.

which evaluates to

$$M(\rho') = 2 - \frac{64 \sin^4 \frac{\theta}{2} (13 - 4 \cos \theta - \cos 2\theta)}{[6(3 + \cos 2\theta) - 15 \cos \theta - \cos 3\theta]^2}. \quad (11)$$

The CHSH criterion quantifies quantum steering correlations and distinguishes their strength based on the value of $M(\rho')$, given in Eq. (11), according to

$$M(\rho') = \begin{cases} > 1 & \text{(quantum steering),} \\ \leq 1 & \text{(no quantum steering).} \end{cases} \quad (12)$$

The results from the analysis given in Eq. (12) are presented in Fig. 4(a). Additionally, the directional dependence of quantum steering correlations is illustrated in Figs. 4(b) and 4(c), which display the computed values of $M(\rho')$ projected onto the surface of a unit sphere in the scattering coordinate system. This spherical representation visually emphasizes angular symmetries and how quantum steering vary spatially.

To further interpret these results, the quantum steering correlations identified through the CHSH inequality are com-

pared with the concurrence $\mathcal{C}(\rho')$, a measure of entanglement strength, as summarized in Table I. While concurrence quantifies the presence and degree of entanglement, the CHSH inequality clarifies whether this entanglement can manifest as experimentally accessible nonlocal correlations useful for practical quantum information protocols.

This comparison categorizes the angular scattering ranges into three distinct regimes of quantum correlations. The first regime, distillable entanglement, occurs at angles exhibiting CHSH inequality violations ($M(\rho') > 1$), indicating photon-pair entanglement suitable for practical extraction and application in quantum protocols. The second regime, potentially undistillable entanglement, emerges at angles characterized by nonzero concurrence but without CHSH violation [$M(\rho') \leq 1$]. This classification explicitly follows from the Peres conjecture [35–37], which posits that entangled two-qubit states not violating the CHSH inequality represent bound or undistillable entanglement, that is, entanglement that cannot be converted into practically usable quantum

TABLE I. Summary of quantum correlation regimes identified by concurrence and CHSH analyses. The classification of entanglement distillation explicitly follows from the Peres conjecture, implying that all entangled two-qubit states not violating the CHSH inequality may represent bound (undistillable) entanglement.

Angular range (θ)	Quantum entangled		Quantum steering (CHSH)		Distillation type
	$\mathcal{C}(\rho') > 0$ (Yes)	$\mathcal{C}(\rho') = 0$ (No)	$M(\rho') > 1$ (Yes)	$M(\rho') \not> 1$ (No)	
$0^\circ \leq \theta < 69.3^\circ$	Yes		Yes		Distillable
$69.3^\circ \leq \theta < 85.5^\circ$	Yes		No		Potentially undistillable
$85.5^\circ \leq \theta \leq 96^\circ$	No		No		Not distillable (separable)
$96^\circ < \theta \leq 131.6^\circ$	Yes		No		Potentially undistillable
$131.6^\circ < \theta \leq 180^\circ$	Yes		Yes		Distillable

resources. Finally, the third regime corresponds to angles at which concurrence vanishes ($\mathcal{C}(\rho') = 0$), representing separable states without quantum correlations, where photon pairs behave classically.

V. DISCUSSION

The analysis presented assumes that Compton scattering occurs between an annihilation photon and a stationary, isolated electron in a vacuum. Under these ideal conditions, the combined photon-electron system evolves unitarily and remains globally pure throughout the interaction.

The absence of environmental interactions precludes interpreting the observed loss of entanglement as conventional decoherence, which typically results from interactions with thermal baths, optical losses, or stochastic noise. Instead, the intrinsic unpredictability of Compton scattering outcomes originates entirely from the probabilistic structure inherent in the classical Mueller scattering matrix, independent of external classical noise. Employing a spin-averaged Mueller matrix mathematically captures the information loss associated with tracing over electron spin degrees of freedom, representing a purely mathematical reduction rather than a physical decoherence process.

Any apparent loss of entanglement observed in the reduced photon subsystem must therefore be interpreted from this partial observational perspective. Although the global state remains pure, tracing out the electron leads to a mixed and potentially separable two-photon state.

This result highlights a fundamental quantum mechanical principle: while global states evolve unitarily and remain pure, partial observations (e.g., restricting to photons only) yield effective dynamics that appear nonunitary. Hence, separability of photon states in certain angular regions reflects this effective description rather than genuine dissipation or nonunitary processes at the global scale.

From the perspective of quantum-information applications, the analysis identifies three distinct angular regimes of photon-pair correlations after Compton scattering: (i) angles exhibiting strong entanglement and quantum steering, useful for quantum-information tasks; (ii) intermediate angles with measurable entanglement but no CHSH violation, potentially representing bound entanglement; and (iii) angles where photon pairs become fully separable, excluding their use in entanglement-based protocols.

A physical interpretation of these angular regimes can be given by considering how Compton scattering redistributes polarization correlations between the photons. In the forward and backward directions ($0^\circ \leq \theta < 69.3^\circ$ and $131.6^\circ < \theta \leq 180^\circ$), the scattering amplitudes preserve strong polarization correlations between the annihilation photons, which supports both entanglement (Fig. 3) and the CHSH inequality violation (Fig. 4). These regions therefore correspond to distillable entanglement as summarized in Table I. At intermediate angles ($69.3^\circ \leq \theta < 85.5^\circ$ and $96^\circ < \theta \leq 131.6^\circ$), interference between different polarization-dependent scattering channels leads to partial cancellation of nonlocal correlations. As a result, concurrence remains nonzero (Fig. 3), showing that the photons are still entangled, but the steering measure $M(\rho')$ drops below unity (Fig. 4). According to the Peres conjecture, this entanglement is not distillable since it cannot be converted into nonlocal correlations useful for quantum information tasks. This classification is reflected in the *potentially undistillable* entries of Table I. Near orthogonal scattering ($85.5^\circ \leq \theta \leq 96^\circ$), the reduced two-photon density matrix approaches maximal mixedness, as already indicated by the purity minimum in Fig. 2(d). In this central angular window, both concurrence (Fig. 3) and $M(\rho')$ (Fig. 4) vanish, demonstrating that photon pairs become fully separable, consistent with the *not distillable (separable)* regime in Table I. These three regimes therefore follow directly from the interplay between scattering geometry and the polarization structure of the Compton process, clarifying why entanglement, steering, or neither are supported in the respective angular ranges.

VI. CONCLUSION

The quantum-information potential of these photons depends on the Compton scattering angle. Regions demonstrating significant entanglement and quantum steering (CHSH inequality violation) theoretically offer quantum-information resources applicable to quantum protocols. Conversely, intermediate angular regions exhibiting entanglement without steering (possibly bound entanglement) may have a relatively limited practical utility, conditional on the validity of the Peres conjecture. Photon pairs in angular ranges showing complete separability are unsuitable for quantum-information applications.

Future work will investigate how electron spin polarization influences these quantum correlations, with the aim of identifying conditions under which Compton-scattered

photons can serve reliably and robustly in quantum-information applications. In addition, a dedicated analysis of individual scattering channels will be required to determine the origin of the systematic probability difference between $|\psi_1\rangle$ and $|\psi_2\rangle$, which differ only by a relative phase in their eigenbasis. This will involve obtaining the matrix elements prior to any spin averaging, inserting them into the transfer matrix, and then applying the method developed in the present work. Future work will also examine the angular interval where the concurrence vanishes despite the presence of entangled eigenvectors in the spectral decomposition, in order to determine whether this feature arises solely from spin averaging or reflects a deeper symmetry of the Compton transfer.

ACKNOWLEDGMENTS

This work was supported by the Science and Technology Facilities Council, Grant No. ST/W006383/1, the Japan Society for the Promotion of Science (JSPS), Grant No. 22H05022 (KAKENHI), and additional private funding from P. Caradonna.

DATA AVAILABILITY

The data that support the findings of this article are not publicly available. The data are available from the authors upon reasonable request.

APPENDIX A: DENSITY MATRIX OF THE EVOLVED STATE

The functional form of the ρ'_{mn} matrix elements, given in Eq. (3), where $0 \leq \theta \leq \pi$ and $0 \leq \phi \leq 2\pi$ are given by

$$\begin{aligned}\rho'_{11}(\theta) &= \frac{2 \cos^2 \frac{\theta}{2} (3 - \cos \theta)}{6(3 + \cos 2\theta) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{12}(\theta) &= \frac{\sin^2 \theta (2 - \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{13}(\theta, \phi) &= -\frac{e^{2i\phi} \sin^2 \theta (2 - \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{14}(\theta, \phi) &= -\frac{4e^{2i\phi} \cos^4 \frac{\theta}{2} (2 - \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{22}(\theta) &= \frac{4 \sin^4 \frac{\theta}{2} (3 - 2 \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{23}(\theta, \phi) &= -\frac{4e^{2i\phi} \sin^4 \frac{\theta}{2} (2 - \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{24}(\theta, \phi) &= -\frac{e^{2i\phi} \sin^2 \theta (2 - \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{33}(\theta) &= \frac{4 \sin^4 \frac{\theta}{2} (3 - 2 \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{34}(\theta, \phi) &= \frac{\sin^2 \theta (2 - \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}, \\ \rho'_{44}(\theta) &= \frac{2 \cos^2 \frac{\theta}{2} (3 - \cos \theta)}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta}.\end{aligned}$$

APPENDIX B: STATE PROBABILITY FUNCTIONS

The analytic expressions for the state probabilities shown in Fig. 2(a) are given below.

Let

$$\begin{aligned}f(\theta) &= 8 \\ &\times \sqrt{\cos \theta (1 - \cos \theta)^2 (21 \cos \theta - 18 \cos^2 \theta + 5 \cos^3 \theta - 16) + 16}.\end{aligned}\quad (\text{B1})$$

Then

$$P_1(\theta) = \frac{60 - 44 \cos \theta + 20 \cos 2\theta - 4 \cos 3\theta + f(\theta)}{8[6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta]},$$

and

$$P_2(\theta) = \frac{60 - 44 \cos \theta + 20 \cos 2\theta - 4 \cos 3\theta - f(\theta)}{8[6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta]}.$$

The eigenvalues $P_3(\theta)$ and $P_4(\theta)$ are given by

$$P_3(\theta) = \frac{2 \sin^2 \frac{\theta}{2} \sin^2 \theta}{6(\cos 2\theta + 3) - 15 \cos \theta - \cos 3\theta},$$

and

$$P_4(\theta) = \frac{8 \sin^6 \frac{\theta}{2}}{6 \cos 2\theta - 15 \cos \theta - \cos 3\theta + 18}.$$

APPENDIX C: FUNCTIONAL FORMS OF NORMALIZATION AND PHASE FACTORS

The eigenstates $|\psi_1\rangle$ and $|\psi_2\rangle$ given by Eqs. (6a) and (6b) depend on normalization factors, $a_1(\theta)$ and $a_2(\theta)$, as well as relative phase factors, $c_1(\theta)$ and $c_2(\theta)$. This appendix provides their functional forms, derived from the spectral decomposition of the density matrix in Eq. (5).

The normalization factors $a_1(\theta)$ and $a_2(\theta)$ ensure that the eigenstates $|\psi_1\rangle$ and $|\psi_2\rangle$ have unit norm. They are defined as

$$\begin{aligned}a_1(\theta) &= \frac{16(2 - \cos \theta)}{\sqrt{\csc^4 \theta [f(\theta) - 8 \cos \theta (3 - \cos \theta)^2]^2 + 256(2 - \cos \theta)^2}}, \\ a_2(\theta) &= \frac{16(2 - \cos \theta)}{\sqrt{\csc^4 \theta [f(\theta) + 8 \cos \theta (3 - \cos \theta)^2]^2 + 256(2 - \cos \theta)^2}}.\end{aligned}$$

The factor $c_1(\theta)$ quantifies the relative phase relationship between the Bell-like basis states $|\Phi_1^-\rangle$ and $|\Psi_2^-\rangle$ comprising the eigenstate $|\psi_1\rangle$. It is expressed as

$$c_1(\theta) = \frac{f(\theta) - 8 \cos \theta (3 - \cos \theta)^2}{16 \sin^2 \theta (2 - \cos \theta)}.$$

Likewise, the relative phase factor $c_2(\theta)$ characterizes the corresponding phase relationship within the eigenstate $|\psi_2\rangle$ and is given by

$$c_2(\theta) = \frac{f(\theta) + 8 \cos \theta (3 - \cos \theta)^2}{16 \sin^2 \theta (2 - \cos \theta)}.$$

In these expressions, $f(\theta)$ is a shorthand notation representing the angular-dependent function defined in Eq. (B1).

These amplitude and phase factors fully determine how the eigenstates $|\psi_1\rangle$ and $|\psi_2\rangle$ evolve continuously from maxi-

mally entangled to separable states as the Compton scattering angle θ varies.

- [1] N. Prantzos, C. Boehm, A. M. Bykov, R. Diehl, K. Ferrière, N. Guessoum, P. Jean, J. Knoedlseder, A. Marcowith, I. V. Moskalenko, A. Strong, and G. Weidenspointner, The 511 keV emission from positron annihilation in the galaxy, *Rev. Mod. Phys.* **83**, 1001 (2011).
- [2] A. Berera, Quantum coherence to interstellar distances, *Phys. Rev. D* **102**, 063005 (2020).
- [3] A. Berera and J. Calderón-Figueroa, Viability of quantum communication across interstellar distances, *Phys. Rev. D* **105**, 123033 (2022).
- [4] X.-M. Hu, Y. Guo, B.-H. Liu, C.-F. Li, and G.-C. Guo, Progress in quantum teleportation, *Nat. Rev. Phys.* **5**, 339 (2023).
- [5] I. Gauthier and M. Hawton, Photon correlations in positron annihilation, *Phys. Rev. A* **81**, 062121 (2010).
- [6] G. J. Rozing, P. E. Mijnders, and R. Benedek, Fully relativistic calculation of the two-photon momentum distribution for positron annihilation in tungsten, *Phys. Rev. B* **43**, 6996 (1991).
- [7] S. D. Bass, S. Mariuzzi, P. Moskal, and E. Stepień, *Colloquium: Positronium physics and biomedical applications*, *Rev. Mod. Phys.* **95**, 021002 (2023).
- [8] S. Basu, T. C. Kwee, S. Surti, E. A. Akin, D. Yoo, and A. Alavi, Fundamentals of PET and PET/CT imaging, *Ann. N.Y. Acad. Sci.* **1228**, 1 (2011).
- [9] E. Nevo, P. Kamvosoulis, and G. Currie, PET/MRI, part 3: Protocols and procedures, *J. Nucl. Med. Technol.* **50**, 17 (2022).
- [10] P. Caradonna, Kinematic analysis of multiple Compton scattering in quantum-entangled two-photon systems, *Ann. Phys. (NY)* **470**, 169779 (2024).
- [11] A. Zatcepin and S. I. Ziegler, Detectors in positron emission tomography, *Z. Med. Phys.* **33**, 4 (2023), Special Issue: Recent Developments in Nuclear Medicine Imaging and Therapy.
- [12] C. S. Levin, New imaging technologies to enhance the molecular sensitivity of positron emission tomography, *Proc. IEEE* **96**, 439 (2008).
- [13] H. Zaidi, M. L. Montandon, and S. Meikle, Strategies for attenuation compensation in neurological PET studies, *NeuroImage* **34**, 518 (2007).
- [14] D. Bohm and Y. Aharonov, Discussion of experimental proof for the paradox of Einstein, Rosen, and Podolsky, *Phys. Rev.* **108**, 1070 (1957).
- [15] P. Caradonna, I. D'Amico, D. G. Jenkins, and D. P. Watts, Stokes-parameter representation for Compton scattering of entangled and classically correlated two-photon systems, *Phys. Rev. A* **109**, 033719 (2024).
- [16] P. Caradonna, Compton scattering mediated by quantum entanglement, *Phys. Rev. A* **111**, 053708 (2025).
- [17] I. Tkachev, S. Musin, D. Abdurashitov, A. Baranov, F. Guber, A. Ivashkin, and A. Strizhak, Measuring the evolution of entanglement in Compton scattering, *Sci. Rep.* **15**, 6064 (2025).
- [18] J. Bordes, J. R. Brown, D. P. Watts, M. Bashkanov, K. Gibson, R. Newton, and N. Zachariou, First detailed study of the quantum decoherence of entangled gamma photons, *Phys. Rev. Lett.* **133**, 132502 (2024).
- [19] W. H. McMaster, Matrix representation of polarization, *Rev. Mod. Phys.* **33**, 8 (1961).
- [20] S. Shivashankara, Entanglement entropy of Compton scattering with a witness, *Can. J. Phys.* **101**, 757 (2023).
- [21] A. L. McNamara, M. Toghyani, J. E. Gillam, K. Wu, and Z. Kuncic, Towards optimal imaging with PET: An *in silico* feasibility study, *Phys. Med. Biol.* **59**, 7587 (2014).
- [22] H. Eslami and M. Mohamadian, Optimization of the positron emission tomography image resolution by using quantum entanglement concept, *Eur. Phys. J. Plus* **139**, 976 (2024).
- [23] D. Kim, A. N. Rachman, U. Taisei, M. Uenomachi, K. Shimazoe, and H. Takahashi, Background reduction in PET by double Compton scattering of quantum entangled annihilation photons, *J. Instrum.* **18**, P07007 (2023).
- [24] P. Moskal, Towards total-body modular PET for positronium and quantum entanglement imaging, in *2018 IEEE Nuclear Science Symposium and Medical Imaging Conference Proceedings (NSS/MIC)* (IEEE, New York, 2018), pp. 1–4.
- [25] A. M. Kožuljević, T. Bokulić, D. Grošev, Z. Kuncic, S. Parashari, L. Pavelić, and M. Makek, Investigation of the spatial resolution of PET imaging system measuring polarization-correlated Compton events, *Nucl. Instrum. Methods Phys. Res. Sect. A* **1068**, 169795 (2024).
- [26] S. Watanabe, S.-n. Ishikawa, H. Aono, S. Takeda, H. Odaka, M. Kokubun, T. Takahashi, K. Nakazawa, H. Tajima, M. Onishi, and Y. Kuroda, High energy resolution hard x-ray and gamma-ray imagers using CdTe diode devices, *IEEE Trans. Nucl. Sci.* **56**, 777 (2009).
- [27] M. Frandes, A. Zoglauer, V. Maxim, and R. Prost, A tracking Compton-scattering imaging system for hadron therapy monitoring, *IEEE Trans. Nucl. Sci.* **57**, 144 (2010).
- [28] M. Uenomachi, M. Takahashi, K. Shimazoe, H. Takahashi, K. Kamada, T. Orita, K. Ogane, and A. B. Tsuji, Simultaneous *in vivo* imaging with PET and SPECT tracers using a Compton-PET hybrid camera, *Sci. Rep.* **11**, 17933 (2021).
- [29] A. Wightman, Note on polarization effects in Compton scattering, *Phys. Rev.* **74**, 1813 (1948).
- [30] A. F. Abouraddy, A. V. Sergienko, B. E. Saleh, and M. C. Teich, Quantum entanglement and the two-photon Stokes parameters, *Opt. Commun.* **201**, 93 (2002).
- [31] W. K. Wootters, Entanglement of formation of an arbitrary state of two qubits, *Phys. Rev. Lett.* **80**, 2245 (1998).
- [32] S. A. Hill and W. K. Wootters, Entanglement of a pair of quantum bits, *Phys. Rev. Lett.* **78**, 5022 (1997).
- [33] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Proposed experiment to test local hidden-variable theories, *Phys. Rev. Lett.* **23**, 880 (1969).
- [34] R. Horodecki, P. Horodecki, and M. Horodecki, Violating Bell inequality by mixed spin-1/2 states: Necessary and sufficient condition, *Phys. Lett. A* **200**, 340 (1995).
- [35] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, *Rev. Mod. Phys.* **81**, 865 (2009).
- [36] A. Peres, Collective tests for quantum nonlocality, *Phys. Rev. A* **54**, 2685 (1996).
- [37] A. Peres, Separability criterion for density matrices, *Phys. Rev. Lett.* **77**, 1413 (1996).