PHYSICAL REVIEW B THIRD SERIES, VOLUME 64, NUMBER 12 15 SEPTEMBER 2001-II

Ferromagnetism of GaMnAs studied by polarized neutron reflectometry *(4 pages)* . . . . . 121302(R)
H. Kępa, J. Kutner-Pielaszek, A. Twardowski, C. F. Majkrzak, J. Sadowski, T. Story, and T. M. Giebultowicz

Tunneling of zero-dimensional excitons in a single pair of correlated quantum dots *(4 pages)* . . . . . 121303(R)
J. Seufert, M. Obert, G. Bacher, A. Forchel, T. Passow, K. Leonardi, and D. Hommel

Extremely sharp dependence of the exciton oscillator strength on quantum-well width in the $GaN/Al_xGa_{1-x}N$ system: The polarization field effect *(4 pages)* . . . . . 121304(R)
Marian Zamfirescu, Bernard Gil, Nicolas Grandjean, Guilllaume Malpuech, Alexey Kavokin, Pierre Bigenwald, and Jean Massies

Highly anisotropic transport in the integer quantum Hall effect *(4 pages)* . . . . . 121305(R)
W. Pan, H. L. Stormer, D. C. Tsui, L. N. Pfeiffer, K. W. Baldwin, and K. W. West

Enhanced magnetization at integer quantum Hall states *(4 pages)* . . . . . 121306(R)
I. Meinel, D. Grundler, D. Heitmann, A. Manolescu, V. Gudmundsson, W. Wegscheider, and M. Bichler

Effect of surface-energy anisotropy on the kinetics of quantum dot formation *(4 pages)* . . . . . 121307(R)
Fei Long, S. P. A. Gill, and A. C. F. Cocks

Surface physics, low-dimensional systems, and related topics

Conduction mechanisms and magnetotransport in multiwalled carbon nanotubes *(4 pages)* . . . . . 121401(R)
Stephan Roche, François Triozon, Angel Rubio, and Didier Mayou

Conduction channels of Al wires at finite bias *(4 pages)* . . . . . 121402(R)
Nobuhiko Kobayashi, Masakazu Aono, and Masaru Tsukada

Elastic properties of superhard amorphous carbon pressure-synthesized from C_{60} by surface Brillouin scattering *(4 pages)* . . . . . 121403(R)
M. H. Manghnani, S. Tkachev, P. V. Zinin, X. Zhang, V. V. Brazhkin, A. G. Lyapin, and I. A. Trojan

Phase-coherent transport in ropes of single-wall carbon nanotubes *(4 pages)* . . . . . 121404(R)
J. Appenzeller, R. Martel, Ph. Avouris, H. Stahl, U. Th. Hunger, and B. Lengeler

Catalyst-free synthesis of boron nitride single-wall nanotubes with a preferred zig-zag configuration *(4 pages)* . . . . . 121405(R)
R. S. Lee, J. Gavillet, M. Lamy de la Chapelle, A. Loiseau, J.-L. Cochon, D. Pigache, J. Thibault, and F. Willaime

Noncontact atomic force microscope topography dependent on the electrostatic dipole field of individual molecules *(4 pages)* . . . . . 121406(R)
Akira Sasahara, Hiroshi Uetsuka, and Hiroshi Onishi

Chirality-selective Raman scattering of the *D* mode in carbon nanotubes *(4 pages)* . . . . . 121407(R)
J. Maultzsch, S. Reich, and C. Thomsen

ARTICLES

Electronic structure: wide-band, narrow-band, and strongly correlated systems

Boson-fermion model with on-site Coulomb repulsion between fermions *(7 pages)* 125101
Alfonso Romano

Well localized crystalline orbitals obtained from Bloch functions: The case of $KNbO_3$ *(8 pages)* 125102
Ph. Baranek, C. M. Zicovich-Wilson, C. Roetti, R. Orlando, and R. Dovesi

Energy-transfer processes in quasi-bidimensional crystal arrays *(7 pages)* 125103
Sergio O. Vásquez

Nonadiabatic Pauli susceptibility in fullerene compounds *(8 pages)* 125104
E. Cappelluti, C. Grimaldi, and L. Pietronero

(Continued)