

Measurement of π^0 -hadron correlations relative to the event plane in semicentral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV

D. A. H. Abdallah *et al.**
(ALICE Collaboration)



(Received 11 March 2026; accepted 4 June 2026; published 20 August 2026)

Per-trigger yields of π^0 -hadron correlations were measured in semicentral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV in ALICE at the CERN Large Hadron Collider. The reconstructed $\pi^0 \rightarrow \gamma\gamma$, with a transverse momentum of $11 \text{ GeV}/c < p_T(\pi^0) < 14 \text{ GeV}/c$, is used as the trigger particle to calculate yields of associated charged particles on the near- and away-side. The photons are reconstructed using the ALICE Electromagnetic Calorimeter, and the charged particles are measured in the ALICE central barrel within a pseudorapidity range of $|\eta| < 0.8$. The yields are reported relative to the orientation of the π^0 with the second-order event plane and are background subtracted using the reaction-plane fit method. The data give an indication of a suppression of the associated charged-particle yields near $p_T \approx 2 \text{ GeV}/c$ when comparing out-of-plane to in-plane trigger particles. At associated charged-particle $p_T > 3 \text{ GeV}/c$, no significant event-plane dependence is observed within uncertainties. The results are compared with predictions from the JEWEL model, which implements jet energy loss in an expanding medium. JEWEL predicts no significant modification of either the near- or away-side associated yields, independent of whether medium recoils are included. The observed behavior may indicate the presence of additional energy-loss mechanisms beyond those governed by path-length dependence.

DOI: [10.1103/tqp4-kwrt](https://doi.org/10.1103/tqp4-kwrt)

I. INTRODUCTION

Quantum chromodynamics (QCD), the theory of the strong interaction, predicts a transition from ordinary nuclear matter to a deconfined state of quarks and gluons, referred to as the quark-gluon plasma (QGP) [1]. Ultrarelativistic collisions of heavy ions at the BNL Relativistic Heavy Ion Collider (RHIC) and the CERN Large Hadron Collider (LHC) are able to reach the temperatures necessary to create such a state and enable the study of QGP properties [2,3].

During the very early stages of QGP formation, high-energy quarks and gluons are produced from the initial hard scatterings of the colliding nuclei. These partons interact with the medium, fragment, and produce a shower of particles that is referred to as a jet [4–6]. The interactions cause energy loss and angular broadening of partonic showers, which are hypothesized to be proportional to the path length through the medium and fluctuations along the parton's path [7,8]. As the QGP created in semicentral collisions has significant spatial anisotropies, it is expected that jets lose different amounts of energy depending on their emission angle relative to the second-order event plane Ψ_2 [9–13] of final-state particles, which approximates the reaction plane defined by the beam axis and the collision impact-parameter vector [11–13].

The medium-induced modification of jets, or jet quenching, can be studied experimentally by identifying high transverse momentum (p_T) hadrons as a proxy for the jet and measuring the yield of associated particles on the near- and away-side of these trigger particles [14–25]. Here, the near-side corresponds to the region in pseudorapidity η and azimuthal angle φ that is correlated with the direction of the parton shower containing the trigger particle. Conversely, the away-side is the region in (η, φ) where the high- p_T recoil parton is back-to-back in azimuthal angle to the trigger particle. With sufficient energy loss, a suppression of trigger-associated particle yields is expected at the near- and away-side, with a possible dependence on their orientation relative to the event plane that could indicate a path-length dependence. The quenching of jets has been measured at RHIC and the LHC using inclusive and semi-inclusive hadron and jet production [26–29], correlations of dihadrons and dijets [30–36], as well as various substructure measurements and modifications of jet fragmentation functions [37,38].

One of the challenges of these measurements is the fact that the particle multiplicity in jets is small compared with the number of particles produced by hadronization of the medium. Therefore, the measured trigger-hadron correlations will include a significant combinatorial background, especially toward lower particle p_T . Methods to subtract this background, such as the zero-yield-at-minimum (ZYAM) method [39] and the reaction-plane fit (RPF) method [40–42], attempt to subtract the momentum anisotropies due to the flow of the background. In the case of the RPF method, this is done by a simultaneous fit to the in-plane, midplane, and out-of-plane distributions of the trigger-associated particle pairs in $\Delta\varphi$, where $\Delta\varphi$ is the azimuthal angle between the trigger

*Full author list given at the end of the article.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Open access publication funded by CERN.

and the associated particle. Then, after subtraction, the excess yield of associated particles on the near- and away-side can be obtained by integrating over the $\Delta\phi$ distribution.

This paper presents the per-trigger yields of π^0 -hadron correlations in semicentral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV. The yields are obtained for both the away- and near-side, as well as differentially for the trigger orientation relative to the event plane. The results are compared with model predictions from JEWEL [43–45].

II. EXPERIMENTAL SETUP

The ALICE experiment [46] at the LHC is a heavy-ion detector optimized to handle high particle multiplicities over a broad momentum range [47]. The subdetectors used for the analysis presented in this paper are the Inner Tracking System (ITS) [48] and Time Projection Chamber (TPC) [49] for charged-particle reconstruction, the electromagnetic calorimeter (EMCal) [50] for neutral pions ($\pi^0 \rightarrow \gamma\gamma$), and the V0 detectors [51] for event-plane determination. For more information on the ALICE apparatus and its performance, see Refs. [51–53].

The ITS [48] is a six-layer cylindrical silicon detector around the nominal collision point of the experiment. The first two layers are Silicon Pixel Detectors (SPDs), followed by two layers of Silicon Drift Detectors (SDDs) and two layers of Silicon Strip Detectors (SSDs). The TPC [49] is a large-volume cylindrical gaseous detector that covers the full azimuthal angle and $|\eta| < 0.9$ in pseudorapidity. The information from both the ITS and TPC detectors is used for track reconstruction, covering a transverse-momentum range of $0.15 \text{ GeV}/c < p_T < 30 \text{ GeV}/c$ and a pseudorapidity of $|\eta| < 0.9$.

The EMCal [50] is a sampling calorimeter at a radial distance of 4.28 m from the collision point, with an acceptance of $|\eta| < 0.7$ with $80^\circ < \varphi < 187^\circ$ plus $0.22 < |\eta| < 0.7$ with $260^\circ < \varphi < 320^\circ$. It provides an energy resolution of $\sigma(E)/E = a \oplus b/\sqrt{E} \oplus c/E$ with parameters $a = 1.4 \pm 0.1$, $b = 9.5 \pm 0.2 \text{ GeV}^{1/2}$, and $c = 2.9 \pm 0.9 \text{ GeV}$. A clusterization algorithm is used to aggregate the signal of nearby towers into clusters from which the energy and position are calculated. Photon-like clusters were selected above an energy threshold of 2 GeV within 50 ns of the collision time. Additional background was suppressed by removing those clusters that match with a charged-particle track as reconstructed by the ITS and TPC.

The V0A and V0C [51] are forward scintillator detectors located at pseudorapidities $2.8 < \eta < 5.1$ and $-3.7 < \eta < -1.7$, respectively. Each detector has 32 channels, built up from four concentric rings with eight sectors in azimuthal angle. The sum of the amplitudes measured in the V0 detectors is used to define the centrality classes in Pb-Pb collisions. Furthermore, the V0A is used for calculating event-plane angles.

III. ANALYSIS METHODS

The Pb-Pb data used for the analysis were taken during 2015 and 2018, at a center-of-mass energy per nucleon-nucleon pair of $\sqrt{s_{NN}} = 5.02$ TeV. During both runs, the




FIG. 1. Diphoton invariant mass distribution up to $0.8 \text{ GeV}/c^2$ before and after subtraction of the combinatorial background, for pair momenta $11 \text{ GeV}/c < p_T < 14 \text{ GeV}/c$. The orange and green vertical dashed lines indicate the π^0 and selected sideband mass windows, respectively.

level-1 γ -trigger of the EMCal was used to trigger on events that contain a high-momentum photon with a minimum energy threshold of 10 GeV [54]. The analysis uses the data from this trigger as well as from the default minimum-bias trigger, which requires coincident signals in the V0A and V0C. Furthermore, events are required to have 30%–50% collision centrality and a primary vertex within ± 10 cm of the nominal interaction point along the beam line (z direction). Out-of-bunch pile-up was removed by utilizing the correlation between the number of hits in the ITS and TPC detectors. After applying all these event-selection criteria, 1.68 million EMCal level-1 triggered events as well as 60.8 million minimum-bias triggered events are used for the analysis.

A. Neutral pion and charged-particle reconstruction

Neutral pion (π^0) candidates are reconstructed in the two-photon decay channel $\pi^0 \rightarrow \gamma\gamma$ by calculating the invariant mass of EMCal cluster pairs using $M_{\gamma\gamma} = \sqrt{2E_{\gamma 1}E_{\gamma 2}[1 - \cos(\theta_{\gamma 1\gamma 2})]}$. Here, $E_{\gamma 1}$ and $E_{\gamma 2}$ are the energies of the photon-like clusters 1 and 2, respectively, and $\theta_{\gamma 1\gamma 2}$ is the opening angle between the clusters. The minimum energy of the clusters considered is 2 GeV, and clusters too close to the border of the calorimeter are rejected due to possible incorrect energy measurement from losses at the edge. The π^0 's are selected with a transverse momentum of $11 \text{ GeV}/c < p_T < 14 \text{ GeV}/c$. The π^0 signal is an excess yield on top of a combinatorial background, see Fig. 1. The shape of the peak and combinatorial background are estimated using Gaussian

and polynomial fit functions, capturing both the shape of the signal and the background. Note that the η meson is visible around $M_{\gamma\gamma} = 548 \text{ MeV}/c^2$, which is fitted with a Gaussian but otherwise not used in any part of the analysis other than extending the fit range of the background. The mean (μ_{π^0}) and width (σ_{π^0}) of the fit are used to define the integration range of the signal, where $(\mu_{\pi^0} \pm 2\sigma_{\pi^0})$ is the default.

The charged particles, as reconstructed by the ITS and TPC, require at least one hit in the SPD. The yield of charged particles is corrected for finite track reconstruction efficiency using Pb-Pb events that are simulated using the HIJING Monte Carlo generator in combination with the GEANT3 particle-transport code [55,56]. The efficiency is parametrized two-dimensionally as a function of p_T and η and is on the order of 80% at $|\eta| < 0.5$, and decreases to about 70% for larger $|\eta|$.

B. Event-plane determination

A collision of heavy ions with a nonzero impact parameter typically creates an almond-shaped, or anisotropic, overlap region. The plane spanned by connecting the two centroids of the nuclei with the beam axis is commonly referred to as the reaction plane, and is experimentally not directly accessible. However, we can determine the n th-order event-plane angle using the directions of the final-state particles in the V0 detectors from

$$\Psi_n = \frac{1}{n} \arctan \left(\frac{\sum_i w_i \sin(n\varphi_i)}{\sum_i w_i \cos(n\varphi_i)} \right). \quad (1)$$

Here, φ_i and w_i are the azimuthal angle and signal weight, respectively, of the i th segment of the V0A detector [53,57]. For $n = 2$, Ψ_2 , the second-order event plane, is the experimental proxy for the reaction plane. The directions parallel and perpendicular to this plane are referred to as in-plane and out-of-plane angles, respectively, which are the angles of greatest interest for this analysis. Furthermore, the direction of the π^0 trigger with respect to Ψ_2 is used to distinguish the cases for which the trigger is in, mid, or out of plane. Each of these directions is defined by a region of $\pm 30^\circ$ in φ with respect to Ψ_2 , where the midplane is between the in- and out-of-plane windows.

To account for the finite resolution of the Ψ_n [58], we define the experimental resolution of the m th-order event plane for a $v_n = \langle \cos[n(\varphi - \Psi_n)] \rangle$ flow term as

$$R_{n,m} \equiv \langle \cos[n(\Psi_m^{\text{meas}} - \Psi_m^{\text{truth}})] \rangle, \quad (2)$$

with the brackets denoting the average over all events. This can be used to associate the measured values of anisotropic flow coefficients v_n to their true values via

$$v_n^{\text{meas}} = R_{n,m} v_n^{\text{truth}}. \quad (3)$$

The event-plane resolutions can be determined by measuring the correlation between event-plane angles from different subdetectors, specifically over different ranges in η . In this analysis, we determine the event-plane resolution of the V0A event-plane angles using the A-side ($\eta > 0$) and C-side ($\eta < 0$) of the TPC. An average value of $R_{2,2} = 0.7712 \pm 0.0003$

is used to calculate the correction on the ratio of out-of-plane over in-plane yields of π^0 -hadron correlations.

C. π^0 -hadron correlations

The angular correlations between π^0 candidates and charged-particle tracks are measured as a function of the $\Delta\eta$ and $\Delta\varphi$ of each pair, as well as the p_T of the associated charged particle and the angle between the π^0 trigger and the second-order event plane. Furthermore, the acceptance of the ALICE detector geometry is corrected using the event-mixing technique. The mixed-event correlations are calculated between π^0 -hadron candidates, where the π^0 and associated charged hadrons are explicitly from different events. The events used for mixing are matched in terms of event-plane angle, centrality, and position of the event vertex. The mixed-event correlation function $[\text{ME}(\Delta\eta, \Delta\varphi)]$ is normalized such that its maximum value is unity, and any φ dependence in the tracking efficiency cancels in this approach. The correlation between π^0 candidates and charged particles from the same event $[\text{SE}(\Delta\eta, \Delta\varphi)]$ is divided by those from the mixed events, and as such, the correlation function $C(\Delta\eta, \Delta\varphi)$ is defined as

$$C(\Delta\eta, \Delta\varphi) = \frac{\text{SE}(\Delta\eta, \Delta\varphi)}{\text{ME}(\Delta\eta, \Delta\varphi)}. \quad (4)$$

These correlations are projected onto the $\Delta\varphi$ axis in two regions of $\Delta\eta$, namely, the near- $\Delta\eta$ ($|\Delta\eta| < 0.8$) region that includes the near-side jet peak, and the far- $\Delta\eta$ ($0.8 < |\Delta\eta| < 1.35$) region where the impact of jet effects is negligible for $|\Delta\varphi| < \pi/2$.

As the π^0 candidates have contamination from combinatorial pairs of EMCal clusters, there is a background contribution to $C(\Delta\eta, \Delta\varphi)$. Therefore, the measured correlation functions of both signal (C_{sig}) and background (C_{bkg}) are related via the purity $P = N_{\text{sig}}/(N_{\text{sig}} + N_{\text{bkg}})$ of the sample via

$$\frac{1}{N_{\text{trig}}} C = P \frac{1}{N_{\text{sig}}} C_{\text{sig}} + (1 - P) \frac{1}{N_{\text{bkg}}} C_{\text{bkg}}, \quad (5)$$

where N_{trig} , N_{sig} , and N_{bkg} , are the number of π^0 candidates in the total measured, signal, and background region, respectively. As the background cluster pairs may have different correlations with charged hadrons, this effect must be corrected for an accurate measurement of π^0 -hadron correlations. To do this, the correlations between background cluster pairs and hadrons are studied using cluster pairs from a sideband region in the diphoton invariant mass, indicated by the green lines in Fig. 1, wherein the pairs are assumed to be purely combinatorial. For high- p_T cluster pairs, the lower invariant-mass range is kinematically excluded by the effect of cluster merging, which prevents constructing a sideband on the left side of the π^0 , as can be seen in Fig. 1. To accommodate for this, the correlations with the upper-mass sideband are studied as a function of invariant mass and extrapolated to the π^0 -mass region. Having obtained the correlations from the background, the true π^0 -hadron correlations are extracted




FIG. 2. Distributions of π^0 -hadron correlations as a function of $\Delta\phi$ in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, for inclusive, in-, mid-, and out-of-plane orientations of the π^0 -trigger with respect to the event plane. The near- $\Delta\eta$ ($|\Delta\eta| < 0.8$) region that includes the near-side jet peak is shown in blue markers, and the far- $\Delta\eta$ ($0.8 < |\Delta\eta| < 1.35$) region is shown in red markers. The associated charged hadrons have $1.5 \text{ GeV}/c < p_T^{\text{assoc}} < 2.5 \text{ GeV}/c$. The flow background is fitted using the RPF method and shown by the colored solid lines, where the thickness of the line is the statistical uncertainty of the fit.

using the following equation:

$$\frac{1}{N_{\text{trig}}} \frac{d^2N}{d\Delta\eta d\Delta\phi} \equiv \frac{1}{N_{\text{sig}}} C_{\text{sig}} = \frac{1}{P} \left[\frac{1}{N_{\text{trig}}} C - (1 - P) \frac{1}{N_{\text{bkg}}} C_{\text{bkg}} \right]. \quad (6)$$

D. Subtraction of anisotropic flow

In heavy-ion collisions, the spatial anisotropies of the initial state are transformed into momentum-space anisotropies of the final-state particles, which are referred to as anisotropic flow. The particle correlations that arise from this are a dominant background when measuring trigger-hadron correlations in $(\Delta\eta, \Delta\phi)$. In addition, the trigger particle may have correlations with the symmetry planes of the event due to the energy loss being different in and out of plane. When restricting the trigger particles to a specific angular range with respect to the event plane, the subtraction of this background becomes even more important.

The reaction plane fit method (RPF) is used to subtract the contribution from the anisotropic flow of the underlying event [40,41]. This method employs a simultaneous fit of the flow coefficients v_n for both the associated particle (v_n^{assoc}) and the trigger particle (v_n^{trig}) using

$$\frac{dN}{d\Delta\phi} = B \left[1 + \sum_n 2v_n^{\text{assoc}} v_n^{\text{trig}} \cos(n\Delta\phi) \right], \quad (7)$$

where $\Delta\phi$ is the azimuthal angle between the trigger and associated hadron, B is an overall normalization factor, and n is an integer up to $n = 4$. v_n^{assoc} is expected to be close to the charged particle v_n . A similar statement can be made for v_n^{trig} , which is consistent with flow measurements at higher transverse momentum. The v_1 parameter is fixed to the results from an independent measurement [59] to improve the fit quality. For the higher harmonics, instead of fixing the RPF parameters to those of independent measurements, we choose to fit the parameters in the far $\Delta\eta$ region with $|\Delta\phi| < \pi/2$,

where jet-induced trigger-hadron correlations are expected to be negligible.

Figure 2 shows the distributions of π^0 -hadron correlations as a function of $\Delta\phi$ in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, for associated charged hadrons within the transverse-momentum range $1.5 \text{ GeV}/c < p_T^{\text{assoc}} < 2.5 \text{ GeV}/c$. The different panels show the results for inclusive, in-, mid-, and out-of-plane orientations of the π^0 -trigger with respect to the event plane, with the background fitted using the RPF method. It shows a significant $\Delta\phi$ modulation that is expected from the magnitude of anisotropic flow coefficients v_n .

The resulting π^0 -hadron correlations after subtracting the background are shown in Fig. 3, for associated charged hadrons within the transverse-momentum range $1.5 \text{ GeV}/c < p_T^{\text{assoc}} < 2.5 \text{ GeV}/c$ and $|\Delta\eta| < 1.5$. For each of the event-plane orientations of the π^0 -trigger the $\Delta\phi$ distributions have a significant peak on the near-side, while the peaks on the away-side are less significant, especially for midplane and out of plane.

E. Yield of away-side and near-side correlations

In general, a direct hard-scattering component of the π^0 -hadron correlations appears as a back-to-back signal in the azimuthal angle ϕ . We therefore expect this to manifest as an excess of the yield of associated charged hadrons on the away-side and near-side, relative to the direction of the π^0 trigger. These peaks are studied quantitatively by calculating their yields and widths. The yield is calculated using a double integral in $\Delta\eta$ and $\Delta\phi$ as

$$Y = \int_{\Delta\eta} \int_{\Delta\phi} \frac{1}{N_{\text{trig}}} \frac{d^2N}{d(\Delta\eta)d(\Delta\phi)} d(\Delta\phi) d(\Delta\eta). \quad (8)$$

For the near-side peak, the $\Delta\phi$ limits are $-\pi/3 \text{ rad} < \Delta\phi < \pi/3 \text{ rad}$. For the away-side peak, the limits are $2\pi/3 \text{ rad} < \Delta\phi < 4\pi/3 \text{ rad}$. Since the far- $\Delta\eta$ region on the near-side is used to calculate the background with the RPF, that region is not used in order to avoid autocorrelation




FIG. 3. Background-subtracted distributions of π^0 -hadron correlations as a function of $\Delta\phi$ in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV for inclusive, in-, mid-, and out-of-plane orientations of the π^0 trigger with respect to the event plane. The associated charged hadrons have $1.5 \text{ GeV}/c < p_{\text{T}}^{\text{assoc}} < 2.5 \text{ GeV}/c$.

effects. Therefore, the $\Delta\eta$ limits for the near-side are given by $|\Delta\eta| < 0.8$, and on the away-side by $|\Delta\eta| < 1.35$.

Furthermore, uncertainties of the RPF parameters introduce uncertainties on the extracted yield via changes in the background subtraction. This uncertainty on the yield is labeled the RPF uncertainty and is calculated by varying the RPF parameters according to their covariance matrix. The RPF uncertainty is reported as a separate statistical uncertainty on the yield of π^0 -hadron correlations.

IV. SYSTEMATIC UNCERTAINTIES

The systematic uncertainties related to the extracted yield of π^0 -hadron correlations on the near-side and away-side are listed in Table I for the inclusive, in-plane, midplane, and out-of-plane trigger orientations. Table II shows the systematic uncertainties for the yield ratios of out of plane to in plane, which are calculated directly on the ratio, taking into account the correlations and are found to partially cancel. The systematic uncertainties are indicated by a range that corresponds to the minimum and maximum size of the respective uncertainty as a function of $p_{\text{T}}^{\text{assoc}}$. The total systematic uncertainty of the measurement is calculated by treating the separate contributions as uncorrelated and adding them in

quadrature. The sources of systematic uncertainties will now be discussed briefly.

Purity of the π^0 -trigger. As the π^0 candidates are reconstructed from two photon-like clusters, they contain a combinatorial background from random pairings. The systematic uncertainty of the purity is obtained by varying the fit functions and fit range of the $M_{\gamma\gamma}$ distribution of Fig. 1. The yields on the away-side and near-side have been recalculated using the variations in purity, of which the rms is taken as the systematic uncertainty. The relative systematic uncertainty assigned to this source is at most 7%.

π^0 -trigger sideband subtraction. The correlation C_{bkg} is estimated by changing the sidebands in the $M_{\gamma\gamma}$ invariant-mass distribution, shown as green dot-dashed lines in Fig. 1. This sideband has been varied, as well as the methods to extrapolate to the π^0 -mass region. This relative uncertainty ranges from 1%–27%, depending on the event-plane orientation and $p_{\text{T}}^{\text{assoc}}$, and is overall slightly higher on the near-side.

$\Delta\eta$ and $\Delta\phi$ selection. The range in $\Delta\eta$ that defines the near-side is by default chosen to be $|\Delta\eta| < 0.8$ and, for the away-side, $|\Delta\eta| < 1.35$. Since the yields are normalized to $\Delta\eta$, any variation due to this choice is corrected for. However, depending on the range, it is possible to either include or exclude regions of true π^0 -hadron correlations. As such, $\Delta\eta$

TABLE I. Sources of relative systematic uncertainties for the yield of near-side ($|\Delta\eta| < 0.8$) and away-side ($|\Delta\eta| < 1.35$) π^0 -hadron correlations in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, for the inclusive, in-, mid-, and out-of-plane orientations of the π^0 -trigger with respect to the event plane.

	near-side				away-side			
	inclusive	in plane	midplane	out of plane	inclusive	in plane	midplane	out of plane
π^0 purity	1%–5%	1%–7%	1%–6%	1%–5%	0%–5%	0.5%–6%	0.5%–5%	0%–3%
Sideband subtraction	3%–18%	2%–14%	3%–27%	2%–17%	1%–16%	3%–8%	2%–26%	2%–10%
$\Delta\eta$ selection	7%–20%	13%–25%	14%–23%	13%–24%	5%–14%	13%–22%	15%–27%	17%–21%
$\Delta\phi$ selection	0.5%–4%	0.5%–3%	0.5%–6%	0.5%–3%	3%–17%	3%–17%	3%–18%	4%–14%
v_2^{trig} treatment	0%–1%	0%–7%	0%–6%	0%–17%	0%–1%	0%–10%	0%–8%	0%–18%
$v_1^{\text{trig}} v_1^{\text{assoc}}$ calculation	<1%	<1%	<1%	<1%	<1%	<1%	<1%	<1%
Tracking efficiency	4%	4%	4%	4%	4%	4%	4%	4%
Total	13%–25%	18%–27%	19%–34%	15%–31%	9%–23%	18%–26%	18%–42%	15%–28%

TABLE II. Sources of relative systematic uncertainties for the ratio of out-of-plane over in-plane yields of π^0 -hadron correlations on the near-side ($|\Delta\eta| < 0.8$) and away-side ($|\Delta\eta| < 1.35$).

	near-side		away-side	
	$p_T^{\text{assoc}} < 4 \text{ GeV}/c$	$p_T^{\text{assoc}} > 4 \text{ GeV}/c$	$p_T^{\text{assoc}} < 4 \text{ GeV}/c$	$p_T^{\text{assoc}} > 4 \text{ GeV}/c$
π^0 purity	1%–3%	1%–3%	1%–5%	1%–4%
Sideband subtraction	2%–11%	1%–3%	4%–17%	3%–12%
$\Delta\eta$ selection	12%–17%	1%–13%	10%–50%	6%–11%
$\Delta\phi$ selection	4%–6%	1%	6%–10%	1%–4%
v_2^{trig} treatment	1%–11%	<1%	4%–33%	1%
$v_1^{\text{trig}} v_1^{\text{assoc}}$ calculation	<1%	<1%	<1%	<1%
Tracking efficiency	<1%	<1%	<1%	<1%
Event-plane res. corr.	1%–4%	1%	1%	1%
Total	13%–35%	3%–14%	26%–65%	6%–16%

and $\Delta\phi$ have been varied, and the effect on the associated charged-hadron yield has been calculated. At most, these relative uncertainties are 27% for $\Delta\eta$ and 18% for $\Delta\phi$.

RPF method. The fit parameters of the RPF method, which vary as a function of p_T^{assoc} , have uncertainties that directly propagate to an uncertainty in the π^0 -hadron correlations and have been accounted for in the statistical uncertainty of the measurement. However, the choice of fixing the fit parameters using input from independent measurements is a source of systematic uncertainty. The variations include fixing the v_2^{trig} parameters to values found with the RPF at lower v_2^{assoc} , where the v_2^{trig} is the same, but with better statistical precision. The v_1 parameter is estimated from independent measurements [59], whose uncertainty is propagated here. The assigned relative systematic uncertainty for the v_2^{trig} treatment is at most 17% for out-of-plane yields, and the relative $v_1^{\text{trig}} v_1^{\text{assoc}}$ uncertainty is negligible (<1%).

Charged-particle tracking efficiency. The relative uncertainty on the tracking efficiency of charged-particle tracks amounts to 4% and results in a scale uncertainty on the yields of π^0 -hadron correlations. This uncertainty has been obtained by varying the tracking parameters that are used for charged tracks matched between the ITS and TPC. In addition, the event-plane dependence of the tracking efficiency has been studied and found to be negligible.

Event-plane resolution. The event-plane resolution varies slightly with centrality, on the order of $\pm 10\%$ from the mean centrality (40%) to more central and peripheral collisions. A variation on $R_{n,m}$ of 4% was chosen, and the effect on the π^0 -hadron yields was used to calculate the corresponding relative systematic uncertainty. It was found to be at most 4% at low p_T^{assoc} on the near-side, and 1% for higher p_T^{assoc} on the away-side.

V. RESULTS

The yields of π^0 -hadron correlations on the near-side and away-side in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02 \text{ TeV}$ are shown in Fig. 4 as a function of p_T^{assoc} for in-, mid-, and out-of-plane orientations of the π^0 -trigger. The π^0 candidates were reconstructed with a transverse momentum within the range $11 \text{ GeV}/c < p_T < 14 \text{ GeV}/c$. For the near-side, the

difference in pseudorapidity of the pairs is $|\Delta\eta| < 0.8$, while for the away-side, it is $|\Delta\eta| < 1.35$. The RPF method is used to subtract the anisotropic flow background for each interval in p_T^{assoc} . The results for both near- and away-side follow a steeply falling spectrum, with mostly insignificant changes in the yield for the different event plane distinctions at a given p_T^{assoc} . Note that the RPF uncertainty for $p_T^{\text{assoc}} = 1.5\text{--}2.5 \text{ GeV}/c$ on both the near and away side is much larger than for any of the other bins. Here, the variations of the RPF parameters, specifically the flow coefficients (v_n^{assoc}), introduce a large variation on the extracted yield.

The event-plane dependence is studied by comparing the ratio of the yields for out-of-plane trigger orientations to in-plane trigger orientations. This is done for the near-side and the away-side peaks, with the results shown in Fig. 5. Note that the event-plane resolution correction, as calculated by Eq. (3), is applied on these ratios, and systematic uncertainties partially cancel in the ratio. A deviation from unity might indicate an event-plane dependence of trigger-hadron measurements. In particular, suppression at higher p_T^{assoc} could indicate event-plane-dependent jet-energy loss, especially on the away-side. Such energy loss could also produce a ratio above unity at lower p_T^{assoc} due to energy conservation, as seen in the analysis of jet-hadron correlations in Au-Au and pp collisions in Ref. [16]. The results are generally consistent with unity, except at $p_T^{\text{assoc}} = 1.5\text{--}2.5 \text{ GeV}/c$ for both the near-side and away-side peaks, where the ratios are below unity with a χ^2 of 16.65 (4.08σ) and 14.42 (3.80σ), respectively. This possibly indicates path-length-dependent energy loss that has a larger impact on charged particles at lower p_T^{assoc} , because, on average, particles produced out of plane traverse a longer path-length in the medium than particles produced in plane. These observations are consistent with Ref. [20].

In addition, the results are compared with predictions made using the JEWEL 2.2.0 model, for the interaction of jets with the QGP medium [44,45]. The JEWEL model was run to match the experimental conditions and produce the same observable, using the two main modes of JEWEL, which are referred to as JEWEL without recoils and JEWEL with recoils. The difference between the modes is whether the recoiling partons from the jet's interaction with the medium are saved into the event to participate in hadronization with the jet. JEWEL with recoils should properly simulate the




FIG. 4. Yield of near-side ($|\Delta\eta| < 0.8$) and away-side ($|\Delta\eta| < 1.35$) π^0 -hadron correlations as a function of p_T^{assoc} in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV, for the in-, mid-, and out-of-plane orientations of the π^0 -trigger with respect to the event plane. The data points are slightly displaced for visibility.

conservation of the jet's lost energy, but introduces a nontrivial background not present in JEWEL without recoils. For both the near-side and the away-side, JEWEL predicts no significant deviation from unity, consistent with the measurement within the uncertainties. However, the data exhibit a stronger suppression at $p_T^{\text{assoc}} = 1.5$ – 2.5 GeV/c.

VI. CONCLUSIONS

The per-trigger yields of π^0 -hadron correlations are presented in this paper for semicentral Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV. The π^0 -trigger was reconstructed using its decay into two photons and with a transverse momentum of $11 \text{ GeV}/c < p_T < 14 \text{ GeV}/c$. The yields are reported relative




FIG. 5. Ratio of out-of-plane over in-plane yields of π^0 -hadron correlations on the near-side ($|\Delta\eta| < 0.8$) and away-side ($|\Delta\eta| < 1.35$), as a function of p_T^{assoc} in Pb-Pb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV.

to the orientation of the π^0 -trigger with the second-order event plane, and are background subtracted using the reaction-plane fit method.

The out-of-plane and in-plane yields are fully consistent within 1σ for $p_T^{\text{assoc}} > 2.5$ GeV/c, on both the near-side and the away-side. At lower p_T^{assoc} , the correlations exhibit a deviation from unity that indicates an event-plane dependent suppression. However, JEWEL predicts minimal event-plane dependence and does not deviate from unity by more than a few percent. This might indicate there are other significant sources of energy loss besides the component related to its path length. The effects at lower momenta should be explored by using model calculations that propagate a full parton shower in an event-by-event hydrodynamical simulation. Additionally, future measurements using a prompt photon as the trigger particle are expected to be less susceptible to jet-selection biases and could therefore exhibit a stronger event-plane dependence. Such measurements would provide a more precise comparison with increasingly comprehensive theoretical models, thereby offering improved sensitivity to the path-length dependence of parton energy loss.

ACKNOWLEDGMENTS

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centers and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Rede Nacional de Física de Altas Energias (Renafae), Financiadora de Estudos e Projetos (Finep), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and The São Paulo Research Foundation (FAPESP), Brazil; Bulgarian Ministry of Education and Science, within the National Roadmap for Research Infrastructures 2020–2027 (object CERN), Bulgaria; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC) and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the VILLUM FONDEN and Danish National Research Foundation (DNRF), Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l’Energie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Center

National de la Recherche Scientifique (CNRS), France; Bundesministerium für Forschung, Technologie und Raumfahrt (BMFTR) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; National Research and Innovation Agency - BRIN, Indonesia; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Académico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Pontificia Universidad Católica del Perú, Peru; Ministry of Science and Higher Education, National Science Centre and WUT ID-UB, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics, Ministry of Research and Innovation and Institute of Atomic Physics and Universitatea Nationala de Stiinta si Tehnologie Politehnica Bucuresti, Romania; Ministerstvo školstva, vyzkumu, vyvoja a mladeze SR, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSTDA) and National Science, Research and Innovation Fund (NSRF via PMU-B B05F650021), Thailand; Turkish Energy, Nuclear and Mineral Research Agency (TENMAK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and U.S. Department of Energy, Office of Nuclear Physics (DOE NP), United States of America. In addition, individual groups or members have received support from: Czech Science Foundation (Grant No. 23–07499S), Czech Republic; FORTE project, Reg. No. CZ.02.01.01/00/22_008/0004632, Czech Republic, co-funded by the European Union, Czech Republic; European Research Council (Grants No. 101220549 and No. 950692), European Union; Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) “Neutrinos and Dark Matter in Astro- and Particle Physics” (Grant No. SFB 1258), Germany; CONVECS project, CUP C97H23001700002 FESR 2021–2027 program, Italy; FAIR - Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy).

DATA AVAILABILITY

This paper has associated data in an HEPData repository (see Ref. [60]).

- [1] W. Busza, K. Rajagopal, and W. van der Schee, Heavy ion collisions: The big picture, and the big questions, *Annu. Rev. Nucl. Part. Sci.* **68**, 339 (2018).
- [2] A. Bazavov *et al.* (HotQCD Collaboration), Equation of state in $(2 + 1)$ -flavor QCD, *Phys. Rev. D* **90**, 094503 (2014).
- [3] C. Ratti, Lattice QCD and heavy ion collisions: A review of recent progress, *Rep. Prog. Phys.* **81**, 084301 (2018).
- [4] M. Gyulassy and M. Plumer, Jet quenching in dense matter, *Phys. Lett. B* **243**, 432 (1990).
- [5] G.-Y. Qin and X.-N. Wang, Jet quenching in high-energy heavy-ion collisions, *Int. J. Mod. Phys. E* **24**, 1530014 (2015).
- [6] M. Connors, C. Nattrass, R. Reed, and S. Salur, Jet measurements in heavy ion physics, *Rev. Mod. Phys.* **90**, 025005 (2018).
- [7] S. Chatrchyan *et al.* (CMS Collaboration), Jet momentum dependence of jet quenching in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Lett. B* **712**, 176 (2012).
- [8] M. Aaboud *et al.* (ATLAS Collaboration), Measurement of the nuclear modification factor for inclusive jets in Pb + Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector, *Phys. Lett. B* **790**, 108 (2019).
- [9] J. Wu, W. Ke, and X.-N. Wang, Bayesian inference of the path-length dependence of jet energy loss, *Phys. Rev. C* **108**, 034911 (2023).
- [10] A. Ogrodnik, M. Rybář, and M. Spousta, Flavor and path-length dependence of jet quenching from inclusive jet and γ -jet suppression, *Eur. Phys. J. C* **85**, 899 (2025).
- [11] K. Aamodt *et al.* (ALICE Collaboration), Elliptic flow of charged particles in Pb-Pb collisions at 2.76 TeV, *Phys. Rev. Lett.* **105**, 252302 (2010).
- [12] C. Beattie, G. Nijs, M. Sas, and W. van der Schee, Hard probe path lengths and event-shape engineering of the quark-gluon plasma, *Phys. Lett. B* **836**, 137596 (2023).
- [13] S. Acharya *et al.* (ALICE Collaboration), Modification of charged-particle jets in event-shape engineered PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Lett. B* **851**, 138584 (2024).
- [14] C. Adler *et al.* (STAR Collaboration), Disappearance of back-to-back high p_T hadron correlations in central Au + Au collisions at $\sqrt{s_{NN}} = 200$ -GeV, *Phys. Rev. Lett.* **90**, 082302 (2003).
- [15] S. S. Adler *et al.* (PHENIX Collaboration), Jet structure from dihadron correlations in $d + Au$ collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **73**, 054903 (2006).
- [16] L. Adamczyk *et al.* (STAR Collaboration), Jet-hadron correlations in $\sqrt{s_{NN}} = 200$ GeV $p + p$ and central Au + Au collisions, *Phys. Rev. Lett.* **112**, 122301 (2014).
- [17] A. Adare *et al.* (PHENIX Collaboration), Suppression of away-side jet fragments with respect to the reaction plane in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **84**, 024904 (2011).
- [18] K. Aamodt *et al.* (ALICE Collaboration), Particle-yield modification in jet-like azimuthal di-hadron correlations in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. Lett.* **108**, 092301 (2012).
- [19] H. Agakishiev *et al.* (STAR Collaboration), Event-plane-dependent dihadron correlations with harmonic v_n subtraction in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **89**, 041901 (2014).
- [20] S. Acharya *et al.* (ALICE Collaboration), Jet-hadron correlations measured relative to the second order event plane in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. C* **101**, 064901 (2020).
- [21] H. Agakishiev *et al.* (STAR Collaboration), Measurements of dihadron correlations relative to the event plane in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Chin. Phys. C* **45**, 044002 (2021).
- [22] A. Adare *et al.* (PHENIX Collaboration), Measurement of two-particle correlations with respect to second- and third-order event planes in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **99**, 054903 (2019).
- [23] M. Abdulhamid *et al.* (STAR Collaboration), Jet-hadron correlations with respect to the event plane in $\sqrt{s_{NN}} = 200$ GeV Au + Au collisions in STAR, *Phys. Rev. C* **109**, 044909 (2024).
- [24] N. J. Abdulameer *et al.* (PHENIX Collaboration), Jet modification via π^0 -hadron correlations in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **110**, 044901 (2024).
- [25] S. Acharya *et al.* (ALICE Collaboration), Observation of medium-induced yield enhancement and acoplanarity broadening of low- p_T jets from measurements in pp and central Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. Lett.* **133**, 022301 (2024).
- [26] S. Acharya *et al.* (ALICE Collaboration), Measurements of inclusive jet spectra in pp and central Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. C* **101**, 034911 (2020).
- [27] J. Adam *et al.* (STAR Collaboration), Measurement of inclusive charged-particle jet production in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **102**, 054913 (2020).
- [28] V. Khachatryan *et al.* (CMS Collaboration), Measurement of inclusive jet cross sections in pp and PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. C* **96**, 015202 (2017).
- [29] G. Aad *et al.* (ATLAS Collaboration), Measurements of the nuclear modification factor for jets in Pb + Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector, *Phys. Rev. Lett.* **114**, 072302 (2015).
- [30] G. Aad *et al.* (ATLAS Collaboration), Observation of a centrality-dependent dijet asymmetry in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector at the LHC, *Phys. Rev. Lett.* **105**, 252303 (2010).
- [31] G. Aad *et al.* (ATLAS Collaboration), Measurements of the suppression and correlations of dijets in Pb + Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. C* **107**, 054908 (2023).
- [32] S. Chatrchyan *et al.*, (CMS Collaboration), Observation and studies of jet quenching in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. C* **84**, 024906 (2011).
- [33] S. Afanasiev *et al.* (PHENIX Collaboration), High- p_T π^0 production with respect to the reaction plane in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **80**, 054907 (2009).
- [34] A. Adare *et al.* (PHENIX Collaboration), Neutral pion production with respect to centrality and reaction plane in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. C* **87**, 034911 (2013).
- [35] J. Adam *et al.* (ALICE Collaboration), Jet-like correlations with neutral pion triggers in pp and central Pb-Pb collisions at 2.76 TeV, *Phys. Lett. B* **763**, 238 (2016).
- [36] S. Acharya *et al.* (ALICE Collaboration), Measurement of isolated photon-hadron correlations in $\sqrt{s_{NN}} = 5.02$ TeV pp and p -Pb collisions, *Phys. Rev. C* **102**, 044908 (2020).
- [37] G. Aad *et al.* (ATLAS Collaboration), Measurement of substructure-dependent jet suppression in Pb-Pb collisions at 5.02 TeV with the ATLAS detector, *Phys. Rev. C* **107**, 054909 (2023).

- [38] S. Acharya *et al.* (ALICE Collaboration), Medium-induced modification of groomed and ungroomed jet mass and angularities in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Lett. B* **864**, 139409 (2025).
- [39] J. Adams *et al.* (STAR Collaboration), Distributions of charged hadrons associated with high transverse momentum particles in pp and Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV, *Phys. Rev. Lett.* **95**, 152301 (2005).
- [40] N. Sharma, J. Mazer, M. Stuart, and C. Nattrass, Background subtraction methods for precision measurements of di-hadron and jet-hadron correlations in heavy ion collisions, *Phys. Rev. C* **93**, 044915 (2016).
- [41] C. Nattrass and T. Todoroki, Event plane dependence of the flow modulated background in dihadron and jet-hadron correlations in heavy ion collisions, *Phys. Rev. C* **97**, 054911 (2018).
- [42] C. Nattrass, N. Sharma, J. Mazer, M. Stuart, and A. Bejnood, Disappearance of the Mach cone in heavy ion collisions, *Phys. Rev. C* **94**, 011901(R) (2016).
- [43] K. C. Zapp, F. Krauss, and U. A. Wiedemann, A perturbative framework for jet quenching, *J. High Energy Phys.* **03** (2013) 080.
- [44] J. G. Milhano and K. C. Zapp, Origins of the di-jet asymmetry in heavy ion collisions, *Eur. Phys. J. C* **76**, 288 (2016).
- [45] K. C. Zapp, JEWEL 2.0.0: Directions for use, *Eur. Phys. J. C* **74**, 2762 (2014).
- [46] K. Aamodt *et al.* (ALICE Collaboration), The ALICE experiment at the CERN LHC, *JINST* **3**, S08002 (2008).
- [47] S. Acharya (ALICE Collaboration), The ALICE experiment—A journey through QCD, *Eur. Phys. J. C* **84**, 813 (2024).
- [48] K. Aamodt *et al.* (ALICE Collaboration), Alignment of the ALICE Inner Tracking System with cosmic-ray tracks, *JINST* **5**, P03003 (2010).
- [49] J. Alme *et al.*, The ALICE TPC, a large 3-dimensional tracking device with fast readout for ultra-high multiplicity events, *Nucl. Instrum. Methods Phys. Res., Sect. A* **622**, 316 (2010).
- [50] S. Acharya *et al.* (ALICE Collaboration), Performance of the ALICE Electromagnetic Calorimeter, *J. Instrum.* **18**, P08007 (2023).
- [51] P. Cortese *et al.* (ALICE Collaboration), ALICE technical design report on forward detectors: FMD, T0 and V0 (2004), <https://cds.cern.ch/record/781854>.
- [52] E. Abbas *et al.* (ALICE Collaboration), Performance of the ALICE VZERO system, *J. Instrum.* **8**, P10016 (2013).
- [53] B. Abelev *et al.* (ALICE Collaboration), Performance of the ALICE experiment at the CERN LHC, *Int. J. Mod. Phys. A* **29**, 1430044 (2014).
- [54] O. Bourrion, N. Arbor, G. Conesa-Balbastre, C. Furget, R. Guernane, and G. Marcotte, The ALICE EMCAL L1 trigger first year of operation experience, *JINST* **8**, C01013 (2013).
- [55] M. Gyulassy and X.-N. Wang, HIJING 1.0: A Monte Carlo program for parton and particle production in high-energy hadronic and nuclear collisions, *Comput. Phys. Commun.* **83**, 307 (1994).
- [56] R. Brun, F. Bruyant, F. Carminati, S. Giani, M. Maire, A. McPherson, G. Patrick, and L. Urban, GEANT detector description and simulation tool, *CERN Program Library Long Write-up* W5013, (CERN, Geneva, 1993) (Edition 1994) (10, 1994).
- [57] S. Voloshin and Y. Zhang, Flow study in relativistic nuclear collisions by Fourier expansion of azimuthal particle distributions, *Z. Phys. C: Part. Fields* **70**, 665 (1996).
- [58] A. M. Poskanzer and S. A. Voloshin, Methods for analyzing anisotropic flow in relativistic nuclear collisions, *Phys. Rev. C* **58**, 1671 (1998).
- [59] B. Abelev *et al.* (ALICE Collaboration), Directed flow of charged particles at midrapidity relative to the spectator plane in Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, *Phys. Rev. Lett.* **111**, 232302 (2013).
- [60] D. Ali Hassan Abdallah *et al.* (ALICE Collaboration), Measurement of π^0 -hadron correlations relative to the event plane in semicentral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, HEPData (collection) (2026), doi:10.17182/hepdata.180693.

D. A. H. Abdallah¹³⁴, I. J. Abualrob¹¹², S. Acharya⁴⁹, K. Agarwal^{23,*}, G. Aglieri Rinella³², L. Aglietta²⁴, N. Agrawal²⁵, Z. Ahammed¹³², S. Ahmad¹⁵, I. Ahuja³⁶, Z. Akbar⁷⁹, V. Akishina³⁸, M. Al-Turany⁹⁴, B. Alessandro⁵⁵, R. Alfaro Molina⁶⁶, B. Ali¹⁵, A. Alici^{25,†}, J. Alme²⁰, G. Alocco²⁴, T. Alt⁶³, I. Altsybeev⁹², C. Andrei⁴⁴, N. Andreou¹¹¹, A. Andronic¹²³, M. Angeletti³², V. Anguelov⁹¹, F. Antinori⁵³, P. Antonioli⁵⁰, N. Apadula⁷¹, H. Appelshäuser⁶³, S. Arcelli^{25,†}, R. Arnaldi⁵⁵, I. C. Arsene¹⁹, M. Arslandok¹³⁵, A. Augustinus³², R. Averbeck⁹⁴, M. D. Azmi¹⁵, H. Baba¹²¹, A. R. J. Babu¹³⁴, A. Badalà⁵², J. Bae¹⁰⁰, Y. Bae¹⁰⁰, Y. W. Baek¹⁰⁰, X. Bai¹¹⁶, R. Bailhache⁶³, Y. Bailung¹²⁵, R. Bala⁸⁸, A. Baldissieri¹²⁷, B. Balis², S. Bangalia¹¹⁴, Z. Banoo⁸⁸, V. Barbasova³⁶, F. Barile³¹, L. Barioglio⁵⁵, M. Barlou²⁴, B. Barman⁴⁰, G. G. Barnaföldi⁴⁵, L. S. Barnby¹¹¹, E. Barreau⁹⁹, V. Barret¹²⁴, L. Barreto¹⁰⁶, K. Barth³², E. Bartsch⁶³, N. Bastid¹²⁴, G. Batigne⁹⁹, D. Battistini^{34,92}, B. Batyunya¹³⁹, L. Baudino^{24,‡}, D. Bauri⁴⁶, J. L. Bazo Alba⁹⁸, I. G. Bearden⁸⁰, P. Becht⁹⁴, D. Behera^{47,77}, S. Behera⁴⁶, M. A. C. Behling⁶³, I. Belikov¹²⁶, V. D. Bella¹²⁶, F. Bellini²⁵, R. Bellwied¹¹², L. G. E. Beltran¹⁰⁵, Y. A. V. Beltran⁴³, G. Bencedi⁴⁵, O. Benchikhi⁷³, A. Bensaoula¹¹², S. Beole²⁴, A. Berdnikova⁹¹, L. Bergmann⁷¹, L. Bernardinis²³, L. Betev³², P. P. Bhaduri¹³², T. Bhalla⁸⁷, A. Bhasin⁸⁸, B. Bhattacharjee⁴⁰, L. Bianchi²⁴, J. Bielčák³⁴, J. Bielčiková⁸³, A. Bilandzic⁹², A. Binoy¹¹⁴, G. Biro⁴⁵, S. Biswas⁴, M. B. Blidaru⁹⁴, N. Bluhme³⁸, C. Blume⁶³, F. Bock⁸⁴, T. Bodova²⁰, L. Boldizsár⁴⁵, M. Bombara³⁶, P. M. Bond³², G. Bonomi^{131,54}, H. Borel¹²⁷, A. Borissov¹³⁹, A. G. Borquez Carcamo⁹¹, E. Botta²⁴, N. Bouchhar¹⁷, Y. E. M. Bouziani⁶³, D. C. Brandibur⁶², L. Bratrud⁶³, P. Braun-Munzinger⁹⁴, M. Bregant¹⁰⁶, M. Broz³⁴, G. E. Bruno^{31,93}, V. D. Buchakchiev³⁵, M. D. Buckland⁸², H. Buesching⁶³, S. Bufalino²⁹, P. Buhler⁷³, N. Burmasov¹³⁹, Z. Buthelezi^{67,120}, A. Bylinkin²⁰, J. C. Cabanillas Noris¹⁰⁵, M. F. T. Cabrera¹¹², H. Caines¹³⁵, A. Caliva²⁸, E. Calvo Villar⁹⁸, J. M. M. Camacho¹⁰⁵, P. Camerini²³, M. T. Camerlingo⁴⁹, F. D. M. Canedo¹⁰⁶, S. Cannito²³

- S. L. Cantway¹³⁵, M. Carabas¹⁰⁹, F. Carnesecchi³², C. Carr⁹⁷, L. A. D. Carvalho¹⁰⁶, J. Castillo Castellanos¹²⁷, M. Castoldi³², F. Catalano¹¹², S. Cattaruzzi²³, R. Cerri²⁴, I. Chakaberia⁷¹, P. Chakraborty¹³³, J. W. O. Chan¹¹², S. Chandra¹³², S. Chapeland³², M. Chartier¹¹⁵, S. Chattopadhyay¹³², M. Chen³⁹, T. Cheng⁶, M. I. Cherciu⁶², C. Cheshkov¹²⁵, D. Chiappara²⁷, V. Chibante Barroso³², D. D. Chinellato⁷³, F. Chinu²⁴, E. S. Chizzali^{92,8}, J. Cho⁵⁷, S. Cho⁵⁷, P. Chochula³², Z. A. Chochulska^{133,11}, P. Christakoglou⁸¹, P. Christiansen⁷², T. Chujo¹²², B. Chytla¹³³, M. Ciacco²⁴, C. Cicalo⁵¹, G. Cimador^{32,24}, F. Cindolo⁵⁰, F. Colamaria⁴⁹, D. Colella³¹, A. Colelli³¹, M. Colocci²⁵, M. Concas³², G. Conesa Balbastre⁷⁰, Z. Conesa del Valle¹²⁸, G. Contin²³, J. G. Contreras³⁴, M. L. Coquet⁹⁹, P. Cortese^{130,55}, M. R. Cosentino¹⁰⁸, F. Costa³², S. Costanza²¹, P. Crochet¹²⁴, M. M. Czarnynoga¹³³, A. Dainese⁵³, E. Dall'occo³², G. Dange³⁸, M. C. Danisch¹⁶, A. Danu⁶², A. Daribayeva³⁸, P. Das³², S. Das⁴, A. R. Dash¹²³, S. Dash⁴⁶, A. De Caro²⁸, G. de Cataldo⁴⁹, J. de Cuveland³⁸, A. De Falco²², D. De Gruttola²⁸, N. De Marco⁵⁵, C. De Martin²³, S. De Pasquale²⁸, R. Deb¹³¹, R. Del Grande³⁴, L. Dello Stritto³², G. G. A. de Souza^{106,1}, P. Dhankher¹⁸, D. Di Bari³¹, M. Di Costanzo²⁹, A. Di Mauro³², B. Di Ruzza^{129,49,†}, B. Diab³², Y. Ding⁶, J. Ditzel⁶³, R. Divià³², U. Dmitrieva⁵⁵, A. Dobrin⁶², B. Dönigus⁶³, L. Döpper⁴¹, L. Drzensla², J. M. Dubinski¹³³, A. Dubla⁹⁴, P. Dupieux¹²⁴, N. Dzalaiova¹³, T. M. Eder¹²³, E. C. Ege⁶³, R. J. Ehlers⁷¹, F. Eisenhut⁶³, R. Ejima⁸⁹, D. Elia⁴⁹, B. Erasmus⁹⁹, F. Ercolessi²⁵, B. Espagnon¹²⁸, G. Eulisse³², D. Evans⁹⁷, L. Fabbietti⁹², G. Fabbri⁵⁰, M. Faggin³², J. Faivre⁷⁰, W. Fan¹¹², T. Fang⁶, A. Fantoni⁴⁸, A. Feliciello⁵⁵, W. Feng⁶, A. Fernández Téllez⁴³, B. Fernando¹³⁴, L. Ferrandi¹⁰⁶, A. Ferrero¹²⁷, C. Ferrero^{55,**}, A. Ferretti²⁴, D. Finogeev¹³⁹, F. M. Fionda⁵¹, A. N. Flores¹⁰⁴, S. Foertsch⁶⁷, I. Fokin⁹¹, U. Follo^{55,**}, R. Forynski¹¹¹, E. Fragiaco⁵⁶, H. Friberg⁹², U. Fuchs³², D. Fuligno²³, N. Funicello²⁸, C. Furget⁷⁰, A. Furs¹³⁹, T. Fusayasu⁹⁵, J. J. Gaardhøje⁸⁰, M. Gagliardi²⁴, A. M. Gago⁹⁸, T. Gahlaut⁴⁶, C. D. Galvan¹⁰⁵, S. Gami⁷⁷, C. Garabatos⁹⁴, J. M. Garcia⁴³, E. Garcia-Solis⁹, S. Garetti¹²⁸, C. Gargiulo³², P. Gasik⁹⁴, A. Gautam¹¹⁴, M. B. Gay Ducati⁶⁵, M. Germain⁹⁹, R. A. Gernhauser⁹², M. Giacalone³², G. Gioachin²⁹, S. K. Giri¹³², P. Giubellino⁵⁵, P. Giubileo²⁷, P. Glässel⁹¹, E. Glimos¹¹⁹, M. G. F. S. A. Gomes⁹¹, L. Gonella²³, V. Gonzalez¹³⁴, M. Gorgon², K. Goswami⁴⁷, S. Gotovac³³, V. Grabski⁶⁶, L. K. Graczykowski¹³³, E. Grecka⁸³, A. Grelli⁵⁸, C. Grigoras³², S. Grigoryan^{1,139}, O. S. Groettvik³², M. Gronbeck⁴¹, F. Grosa³², S. Gross-Börling⁹⁴, J. F. Grosse-Oetringhaus³², R. Grosso⁹⁴, D. Grund³⁴, N. A. Grunwald⁹¹, R. Guernane⁷⁰, M. Guibaud⁹⁹, K. Gulbrandsen⁸⁰, J. K. Gumprecht⁷³, T. Gündem⁶³, T. Gunji¹²¹, J. Guo¹⁰, W. Guo⁶, A. Gupta⁸⁸, R. Gupta⁸⁸, R. Gupta⁴⁷, K. Gwizdzial¹³³, L. Gyulai⁴⁵, T. Hachiya⁷⁵, C. Hadjidakis¹²⁸, F. U. Haider⁸⁸, S. Haidlova³⁴, M. Haldar⁴, W. Ham¹⁰⁰, H. Hamagaki⁷⁴, Y. Han¹³⁷, R. Hannigan¹⁰⁴, J. Hansen⁷², J. W. Harris¹³⁵, A. Harton⁹, M. V. Hartung⁶³, A. Hasan¹¹⁸, H. Hassan¹¹³, D. Hatzifotiadou⁵⁰, P. Hauer⁴¹, L. B. Havener¹³⁵, E. Hellbär³², H. Helstrup³⁷, M. Hemmer⁶³, S. G. Hernandez¹¹², G. Herrera Corral⁸, K. F. Hetland³⁷, B. Heybeck⁶³, H. Hillemanns³², B. Hippolyte¹²⁶, I. P. M. Hobus⁸¹, F. W. Hoffmann³⁸, B. Hofman⁵⁸, Y. Hong⁵⁷, A. Horzyk², Y. Hou^{11,94}, P. Hristov³², L. M. Huhta¹¹³, T. J. Humanic⁸⁵, V. Humlova³⁴, M. Husar⁸⁶, A. Hutson¹¹², D. Hutter³⁸, M. C. Hwang¹⁸, M. Inaba¹²², A. Isakov⁸¹, T. Isidori¹¹⁴, M. S. Islam⁴⁶, M. Ivanov⁹⁴, M. Ivanov¹³, K. E. Iversen⁷², M. Jablonski², B. Jacak^{18,71}, N. Jacazio²⁵, P. M. Jacobs⁷¹, A. Jadlovská¹⁰², S. Jadlovská¹⁰², S. Jaelani⁷⁹, J. N. Jager⁶³, C. Jahnke¹⁰⁷, M. J. Jakubowska¹³³, E. P. Jamro², D. M. Janik³⁴, M. A. Janik¹³³, C. A. Jauch⁹⁴, S. Ji¹⁶, Y. Ji⁹⁴, S. Jia⁸⁰, T. Jiang¹⁰, A. A. P. Jimenez⁶⁴, S. Jin¹⁰, F. Jonas⁷¹, D. M. Jones¹¹⁵, J. M. Jowett^{32,94}, J. Jung⁶³, M. Jung⁶³, A. Junique³², J. Juračka³⁴, J. Kaewjai^{101,115}, A. Kaiser^{32,94}, P. Kalinak⁵⁹, A. Kalweit³², H. Kang¹², A. Karasu Uysal¹³⁶, N. Karatzenis⁹⁷, T. Karavicheva¹³⁹, M. J. Karwowska¹³³, V. Kashyap⁷⁷, M. Keil³², B. Ketzer⁴¹, J. Keul⁶³, S. S. Khade⁴⁷, A. Khuntia⁵⁰, Z. Khuranova⁶³, B. Kileng³⁷, B. Kim¹⁰⁰, D. J. Kim¹¹³, D. Kim¹⁰⁰, E. J. Kim⁶⁸, G. Kim⁵⁷, H. Kim⁵⁷, J. Kim¹³⁷, J. Kim⁵⁷, J. Kim¹³⁷, J. Kim³², M. Kim¹⁸, S. Kim¹⁷, T. Kim¹³⁷, J. T. Kinner¹²³, I. Kisel³⁸, A. Kisiel¹³³, J. L. Klay⁵, J. Klein³², S. Klein⁷¹, C. Klein-Bösing¹²³, M. Kleiner⁶³, A. Kluge³², M. B. Knuesel¹³⁵, C. Kobdaj¹⁰¹, R. Kohara¹²¹, A. Kondratyev¹³⁹, J. Konig⁶³, P. J. Konopka³², G. Kornakov¹³³, M. Korwieser⁹², C. Koster⁸¹, A. Kotliarov⁸³, N. Kovacic⁸⁶, M. Kowalski¹⁰³, V. Kozuharov³⁵, G. Kozlov³⁸, I. Králik⁵⁹, A. Kravčáková³⁶, M. A. Krawczyk³², L. Krcal³², F. Krizek⁸³, K. Krizkova Gajdosova³⁴, C. Krug⁶⁵, M. Krüger⁶³, E. Kryshen¹³⁹, V. Kučera⁵⁷, C. Kuhn¹²⁶, D. M. Kumar¹³², L. Kumar⁸⁷, N. Kumar⁸⁷, S. Kumar⁴⁹, S. Kundu³², M. Kuo¹²², P. Kurashvili⁷⁶, S. Kurita⁸⁹, S. Kushpil⁸³, A. Kuznetsov¹³⁹, M. J. Kweon⁵⁷, Y. Kwon¹³⁷, S. L. La Pointe³⁸, P. La Rocca²⁶, A. Lakrathok¹⁰¹, S. Lambert⁹⁹, A. R. Landou⁷⁰, R. Langoy¹¹⁸, P. Larionov³², E. Laudi³², L. Lautner⁹², R. A. N. Laveaga¹⁰⁵, R. Lavicka⁷³, R. Lea^{131,54}, J. B. Lebert³⁸, H. Lee¹⁰⁰, S. Lee⁵⁷, I. Legrand⁴⁴, G. Legras¹²³, A. M. Lejeune³⁴, T. M. Lelek², I. León Monzón¹⁰⁵, M. M. Lesch⁹², P. Lévai⁴⁵, M. Li⁶, P. Li¹⁰, X. Li¹⁰, B. E. Liang-Gilman¹⁸, J. Lien¹¹⁸, R. Lietava⁹⁷, I. Likmeta¹¹², B. Lim⁵⁵, H. Lim¹⁶, S. H. Lim¹⁶, Y. N. Lima¹⁰⁶, S. Lin¹⁰, V. Lindenstruth³⁸, C. Lippmann⁹⁴, D. Liskova¹⁰², D. H. Liu⁶, J. Liu¹¹⁵, Y. Liu⁶, G. S. S. Liveraro¹⁰⁷, I. M. Lofnes^{20,37}, C. Loizides²⁰, S. Lokos¹⁰³, J. Lömker⁵⁸, X. Lopez¹²⁴, E. López Torres⁷, C. Lotteau¹²⁵, P. Lu¹¹⁶, W. Lu⁶, Z. Lu¹⁰, O. Lubyets⁹⁴, G. A. Lucia²⁹, F. V. Lugo⁶⁶, J. Luo³⁹, G. Luparello⁵⁶, J. M. Friedrich⁹², Y. G. Ma³⁹, V. Machacek⁸⁰, M. Mager³², M. Mahlein⁹², A. Maire¹²⁶, E. Majerz², M. V. Makariev³⁵, G. Malfattore⁵⁰, N. M. Malik⁸⁸, N. Malik¹⁵, D. Mallick¹²⁸, N. Mallick¹¹³, G. Mandaglio^{30,52}, S. Mandal⁷⁷, S. K. Mandal⁷⁶, A. Manea⁶², R. Manhart⁹², A. K. Manna⁴⁷, F. Manso¹²⁴, G. Mantzaridis⁹²

- V. Manzari⁴⁹, Y. Mao⁶, R. W. Marcjan², G. V. Margagliotti²³, A. Margotti⁵⁰, A. Marín⁹⁴, C. Markert¹⁰⁴, P. Martinengo³², M. I. Martínez⁴³, M. P. P. Martins^{32,106}, S. Masciocchi⁹⁴, M. Maserà²⁴, A. Masoni⁵¹, L. Massacrier¹²⁸, O. Massen⁵⁸, A. Mastroserio^{129,49}, L. Mattei^{24,124}, S. Mattiazzo²⁷, A. Matyja¹⁰³, J. L. Mayo¹⁰⁴, F. Mazzaschi³², M. Mazzilli³¹, Y. Melikyan⁴², M. Melo¹⁰⁶, A. Menchaca-Rocha⁶⁶, J. E. M. Mendez⁶⁴, E. Meninno⁷³, M. W. Menzel^{32,91}, M. Meres¹³, L. Micheletti⁵⁵, D. Mihai¹⁰⁹, D. L. Mihaylov⁹², A. U. Mikalsen²⁰, K. Mikhaylov¹³⁹, L. Millot⁷⁰, N. Minafra^{114,†}, D. Miśkowiec⁹⁴, A. Modak⁵⁶, B. Mohanty⁷⁷, M. Mohisin Khan^{15,‡}, M. A. Molander⁴², M. M. Mondal⁷⁷, S. Monira¹³³, D. A. Moreira De Godoy¹²³, A. Morsch³², C. Moscatelli²³, T. Mrnjavac³², S. Mrozinski⁶³, V. Muccifora⁴⁸, S. Muhuri¹³², A. Mulliri²², M. G. Munhoz¹⁰⁶, R. H. Munzer⁶³, L. Musa³², J. Musinsky⁵⁹, J. W. Myrcha¹³³, B. Naik¹²⁰, A. I. Nambrath¹⁸, B. K. Nandi⁴⁶, R. Nania⁵⁰, E. Nappi⁴⁹, A. F. Nassirpour¹⁷, V. Nastase¹⁰⁹, A. Nath⁹¹, N. F. Nathanson⁸⁰, C. Nattrass¹¹⁹, A. Neagu¹⁹, L. Nellen⁶⁴, R. Nepeivoda⁷², S. Nese¹⁹, N. Nicassio³¹, B. S. Nielsen⁸⁰, E. G. Nielsen⁸⁰, F. Noferini⁵⁰, S. Noh¹², P. Nomokonov¹³⁹, J. Norman¹¹⁵, N. Novitzky⁸⁴, J. Nystrand²⁰, M. R. Ockleton¹¹⁵, M. Ogino⁷⁴, J. Oh¹⁶, S. Oh¹⁷, A. Ohlson⁷², M. Oida⁸⁹, L. A. D. Oliveira¹⁰⁷, M. H. Oliver¹³⁵, C. Oppedisano⁵⁵, A. Ortiz Velasquez⁶⁴, H. Osanai⁷⁴, J. Otwinowski¹⁰³, M. Oya⁸⁹, K. Oyama⁷⁴, S. Padhan^{131,46}, D. Pagano^{131,54}, V. Pagliarino⁵⁵, G. Pačić⁶⁴, A. Palasciano^{93,49}, I. Panasenko⁷², P. Panigrahi⁴⁶, C. Pantouvakis²⁷, H. Park¹²², J. Park¹²², S. Park¹⁰⁰, T. Y. Park¹³⁷, J. E. Parkkila¹³³, P. B. Pati⁸⁰, Y. Patley⁴⁶, R. N. Patra⁴⁹, J. Patter⁴⁷, B. Paul¹³², F. Pazdic⁹⁷, H. Pei⁶, T. Peitzmann⁵⁸, X. Peng^{53,11}, S. Perciballi²⁴, G. M. Perez⁷, M. Petrovici⁴⁴, S. Piano⁵⁶, M. Pikna¹³, P. Pillot⁹⁹, O. Pinazza^{32,50}, C. Pinto³², S. Pisano⁴⁸, M. Płoskoń⁷¹, A. Plachta¹³³, M. Planinic⁸⁶, D. K. Plociennik², S. Politano³², N. Poljak⁸⁶, A. Pop⁴⁴, S. Porteboeuf-Houssais¹²⁴, J. S. Potgieter¹¹⁰, I. Y. Pozos⁴³, K. K. Pradhan⁴⁷, S. K. Prasad⁴, S. Prasad^{45,47}, R. Preghenella⁵⁰, F. Prino⁵⁵, C. A. Pruneau¹³⁴, M. Puccio³², S. Pucillo²⁸, S. Pulawski¹¹⁷, L. Quaglia²⁴, A. M. K. Radhakrishnan⁴⁷, S. Ragoni¹⁴, A. Rai¹³⁵, A. Rakotozafindrabe¹²⁷, N. Ramasubramanian¹²⁵, L. Ramello^{130,55}, C. O. Ramírez-Álvarez⁴³, M. Rasa²⁶, S. S. Räsänen⁴², R. Rath⁹⁴, M. P. Rauch²⁰, I. Ravasenga³², M. Razza²⁵, K. F. Read^{84,119}, C. Reckziegel¹⁰⁸, A. R. Redelbach³⁸, K. Redlich^{76,88}, H. D. Regules-Medel⁴³, A. Rehman²⁰, F. Reidt³², H. A. Reme-Ness³⁷, K. Reygers⁹¹, M. Richter²⁰, A. A. Riedel⁹², W. Riegler³², A. G. Riffero²⁴, M. Rignanese²⁷, C. Ripoli²⁸, C. Ristea⁶², M. V. Rodriguez³², M. Rodríguez Cahuatzin⁴³, K. Røed¹⁹, E. Rogochaya¹³⁹, D. Rohr³², D. Röhrlich²⁰, S. Rojas Torres³⁴, P. S. Rokita¹³³, G. Romanenko²⁵, F. Ronchetti³², D. Rosales Herrera⁴³, K. Roslon¹³³, A. Rossi⁵³, A. Roy⁴⁷, A. Roy¹¹⁸, S. Roy⁴⁶, N. Rubini⁵⁰, O. Rubza¹⁵, J. A. Rudolph⁸¹, D. Ruggiano¹³³, R. Rui²³, P. G. Russek², A. Rustamov⁷⁸, A. Rybicki¹⁰³, L. C. V. Ryder¹¹⁴, G. Ryu⁶⁹, J. Ryu¹⁶, W. Rzeska⁹², B. Sabiu⁵⁰, R. Sadek⁷¹, S. Sadhu⁴¹, A. Saha³¹, S. Saha⁷⁷, B. Sahoo⁴⁷, R. Sahoo⁴⁷, D. Sahu⁶⁴, P. K. Sahu⁶⁰, J. Saini¹³², S. Sakai¹²², S. Sambyal⁸⁸, D. Samitz⁷³, I. Sanna³², D. Sarkar⁸⁰, V. Sarritzu²², V. M. Sarti⁹², M. H. P. Sas⁸¹, U. Savino²⁴, S. Sawan⁷⁷, E. Scapparone⁵⁰, J. Schambach⁸⁴, H. S. Scheid³², C. Schiaua⁴⁴, R. Schicker⁹¹, F. Schlepper^{32,91}, A. Schmah⁹⁴, C. Schmidt⁹⁴, M. Schmidt⁹⁰, J. Schoengarth⁶³, R. Schotter⁷³, A. Schröter³⁸, J. Schukraft³², K. Schweda⁹⁴, G. Scioli²⁵, E. Scomparin⁵⁵, J. E. Seger¹⁴, D. Sekihata¹²¹, M. Selina⁸¹, I. Selyuzhenkov⁹⁴, S. Senyukov¹²⁶, J. J. Seo⁹¹, L. Serkin^{64,||}, L. Šerkšnytė³², A. Sevcenco⁶², T. J. Shaba⁶⁷, A. Shabetai⁹⁹, R. Shahoyan³², B. Sharma⁸⁸, D. Sharma⁴⁶, H. Sharma⁵³, M. Sharma⁸⁸, S. Sharma⁸⁸, T. Sharma⁴⁰, U. Sharma⁸⁸, O. Sheibani¹³⁴, K. Shigaki⁸⁹, M. Shimomura⁷⁵, Q. Shou³⁹, S. Siddhanta⁵¹, T. Siemiarz⁷⁶, T. F. Silva¹⁰⁶, W. D. Silva¹⁰⁶, D. Silvermyr⁷², T. Simantathammakul¹⁰¹, R. Simeonov³⁵, B. Singh⁴⁶, B. Singh⁸⁸, B. Singh⁹², K. Singh⁴⁷, R. Singh⁷⁷, R. Singh⁵³, S. Singh¹⁵, T. Sinha⁹⁶, B. Sitar¹³, M. Sitta^{130,55}, T. B. Skaali¹⁹, G. Skorodumovs⁹¹, N. Smirnov¹³⁵, K. L. Smith¹⁶, R. J. M. Snellings⁵⁸, E. H. Solheim¹⁹, S. Solokhin⁸¹, C. Sonnabend^{32,94}, J. M. Sonneveld⁸¹, F. Soramel²⁷, A. B. Soto-Hernandez⁸⁵, R. Spijkers⁸¹, C. Sporleder¹¹³, I. Sputowska¹⁰³, J. Staa⁷², J. Stachel⁹¹, L. L. Stahl¹⁰⁶, I. Stan⁶², A. G. Stejskal¹¹⁴, T. Stellhorn¹²³, S. F. Stiefelmaier⁹¹, D. Stocco⁹⁹, I. Storehaug¹⁹, N. J. Strangmann⁶³, P. Stratmann¹²³, S. Strazzi²⁵, A. Sturniolo^{115,30,52}, Y. Su⁶, A. A. P. Suaide¹⁰⁶, C. Suire¹²⁸, A. Suiu¹⁰⁹, M. Sukhanov¹³⁹, M. Suljic³², V. Sumberia⁸⁸, S. Sumowidagdo⁷⁹, P. Sun¹⁰, N. B. Sundstrom⁵⁸, L. H. Tabares⁷, A. Tabikh⁷⁰, S. F. Taghavi⁹², J. Takahashi¹⁰⁷, M. A. Talamantes Johnson⁴³, G. J. Tambave⁷⁷, Z. Tang¹¹⁶, J. Tanwar⁸⁷, J. D. Tapia Takaki¹¹⁴, N. Tapus¹⁰⁹, L. A. Tarasovicova³⁶, M. G. Tazila⁴⁴, A. Tauro³², A. Tavira García^{104,128}, G. Tejeda Muñoz⁴³, L. Terlizzi²⁴, C. Terrevoli⁴⁹, D. Thakur⁵⁵, S. Thakur⁴, M. Thogersen¹⁹, D. Thomas¹⁰⁴, A. M. Tiekoetter¹²³, N. Tiltmann^{32,123}, A. R. Timmins¹¹², A. Toia⁶³, R. Tokumoto⁸⁹, S. Tomassini²⁵, K. Tomohiro⁸⁹, Q. Tong⁶, V. V. Torres⁹⁹, A. Trifiró^{30,52}, T. Triloki⁹³, A. S. Triolo³², S. Tripathy³², T. Tripathy¹²⁴, S. Trogolo²⁴, V. Trubnikov³, W. H. Trzaska¹¹³, T. P. Trzcinski¹³³, C. Tsolanta¹⁹, R. Tu³⁹, R. Turrisi⁵³, T. S. Tveter¹⁹, K. Ullaland²⁰, B. Ulukutlu⁹², S. Upadhyaya¹⁰³, A. Uras¹²⁵, M. Urioni²³, G. L. Usai²², M. Vaid⁸⁸, M. Vala³⁶, N. Valle⁵⁴, L. V. R. van Doremalen⁵⁸, M. van Leeuwen⁸¹, C. A. van Veen⁹¹, R. J. G. van Weelden⁸¹, D. Varga⁴⁵, Z. Varga¹³⁵, P. Vargas Torres⁶⁴, O. Vázquez Doce⁴⁸, O. Vazquez Rueda¹¹², G. Vecil^{23,‡}, P. Veen¹²⁷, E. Vercellin²⁴, R. Verma⁴⁶, R. Vértesi⁴⁵, M. Verweij⁵⁸, L. Vickovic³³, Z. Vilakazi¹²⁰, A. Villani²³, C. J. D. Villiers⁶⁷, T. Virgili²⁸, M. M. O. Virta⁴², A. Vodopyanov¹³⁹, M. A. Völkl⁹⁷, S. A. Voloshin¹³⁴, G. Volpe³¹, B. von Haller³², I. Vorobyev³², N. Vozniuk¹³⁹, J. Vrláková³⁶, J. Wan³⁹, C. Wang³⁹, D. Wang³⁹

Y. Wang¹¹⁶, Y. Wang³⁹, Y. Wang⁶, Z. Wang³⁹, F. Weiglhofer³², S. C. Wenzel³², J. P. Wessels¹²³, P. K. Wiacek², J. Wiechula⁶³, J. Wikne¹⁹, G. Wilk⁷⁶, J. Wilkinson⁹⁴, G. A. Willems¹²³, N. Wilson¹¹⁵, B. Windelband⁹¹, J. Witte⁹¹, M. Wojnar², C. I. Worek², J. R. Wright¹⁰⁴, C.-T. Wu^{6,27}, W. Wu^{92,39}, Y. Wu¹¹⁶, K. Xiong³⁹, Z. Xiong¹¹⁶, L. Xu^{125,6}, R. Xu⁶, Z. Xue⁷¹, A. Yadav⁴¹, A. K. Yadav¹³², Y. Yamaguchi⁸⁹, S. Yang⁵⁷, S. Yang²⁰, S. Yano⁸⁹, Z. Ye⁷¹, E. R. Yeats¹⁸, J. Yi⁶, R. Yin³⁹, Z. Yin⁶, I.-K. Yoo¹⁶, J. H. Yoon⁵⁷, H. Yu¹², S. Yuan²⁰, A. Yuncu⁹¹, V. Zaccolo²³, C. Zampolli³², F. Zanone⁹¹, N. Zardoshti³², P. Závada⁶¹, B. Zhang⁹¹, C. Zhang¹²⁷, M. Zhang^{124,6}, M. Zhang^{27,6}, S. Zhang³⁹, X. Zhang⁶, Y. Zhang¹¹⁶, Y. Zhang¹¹⁶, Z. Zhang⁶, D. Zhou⁶, Y. Zhou⁸⁰, Z. Zhou³⁹, J. Zhu³⁹, S. Zhu^{94,116}, Y. Zhu⁶, A. Zingaretti²⁷, S. C. Zugravel⁵⁵ and N. Zurlo^{131,54}

(ALICE Collaboration)

¹*A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia*

²*AGH University of Krakow, Cracow, Poland*

³*Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kyiv, Ukraine*

⁴*Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India*

⁵*California Polytechnic State University, San Luis Obispo, California, USA*

⁶*Central China Normal University, Wuhan, China*

⁷*Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba*

⁸*Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico*

⁹*Chicago State University, Chicago, Illinois, USA*

¹⁰*China Nuclear Data Center, China Institute of Atomic Energy, Beijing, China*

¹¹*China University of Geosciences, Wuhan, China*

¹²*Chungbuk National University, Cheongju, Republic of Korea*

¹³*Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovak Republic*

¹⁴*Creighton University, Omaha, Nebraska, USA*

¹⁵*Department of Physics, Aligarh Muslim University, Aligarh, India*

¹⁶*Department of Physics, Pusan National University, Pusan, Republic of Korea*

¹⁷*Department of Physics, Sejong University, Seoul, Republic of Korea*

¹⁸*Department of Physics, University of California, Berkeley, California, USA*

¹⁹*Department of Physics, University of Oslo, Oslo, Norway*

²⁰*Department of Physics and Technology, University of Bergen, Bergen, Norway*

²¹*Dipartimento di Fisica, Università di Pavia, Pavia, Italy*

²²*Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy*

²³*Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy*

²⁴*Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy*

²⁵*Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy*

²⁶*Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy*

²⁷*Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy*

²⁸*Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy*

²⁹*Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy*

³⁰*Dipartimento di Scienze MIFT, Università di Messina, Messina, Italy*

³¹*Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy*

³²*European Organization for Nuclear Research (CERN), Geneva, Switzerland*

³³*Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia*

³⁴*Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic*

³⁵*Faculty of Physics, Sofia University, Sofia, Bulgaria*

³⁶*Faculty of Science, P.J. Šafárik University, Košice, Slovak Republic*

³⁷*Faculty of Technology, Environmental and Social Sciences, Bergen, Norway*

³⁸*Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany*

³⁹*Fudan University, Shanghai, China*

⁴⁰*Gauhati University, Department of Physics, Guwahati, India*

⁴¹*Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany*

⁴²*Helsinki Institute of Physics (HIP), Helsinki, Finland*

⁴³*High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico*

⁴⁴*Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania*

⁴⁵*HUN-REN Wigner Research Centre for Physics, Budapest, Hungary*

⁴⁶*Indian Institute of Technology Bombay (IIT), Mumbai, India*

⁴⁷*Indian Institute of Technology Indore, Indore, India*

- ⁴⁸*INFN, Laboratori Nazionali di Frascati, Frascati, Italy*
- ⁴⁹*INFN, Sezione di Bari, Bari, Italy*
- ⁵⁰*INFN, Sezione di Bologna, Bologna, Italy*
- ⁵¹*INFN, Sezione di Cagliari, Cagliari, Italy*
- ⁵²*INFN, Sezione di Catania, Catania, Italy*
- ⁵³*INFN, Sezione di Padova, Padova, Italy*
- ⁵⁴*INFN, Sezione di Pavia, Pavia, Italy*
- ⁵⁵*INFN, Sezione di Torino, Turin, Italy*
- ⁵⁶*INFN, Sezione di Trieste, Trieste, Italy*
- ⁵⁷*Inha University, Incheon, Republic of Korea*
- ⁵⁸*Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University/Nikhef, Utrecht, Netherlands*
- ⁵⁹*Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovak Republic*
- ⁶⁰*Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India*
- ⁶¹*Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic*
- ⁶²*Institute of Space Science (ISS), Bucharest, Romania*
- ⁶³*Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany*
- ⁶⁴*Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico*
- ⁶⁵*Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil*
- ⁶⁶*Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico*
- ⁶⁷*iThemba LABS, National Research Foundation, Somerset West, South Africa*
- ⁶⁸*Jeonbuk National University, Jeonju, Republic of Korea*
- ⁶⁹*Korea Institute of Science and Technology Information, Daejeon, Republic of Korea*
- ⁷⁰*Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France*
- ⁷¹*Lawrence Berkeley National Laboratory, Berkeley, California, USA*
- ⁷²*Lund University Department of Physics, Division of Particle Physics, Lund, Sweden*
- ⁷³*Marietta Blau Institute, Vienna, Austria*
- ⁷⁴*Nagasaki Institute of Applied Science, Nagasaki, Japan*
- ⁷⁵*Nara Women's University (NWU), Nara, Japan*
- ⁷⁶*National Centre for Nuclear Research, Warsaw, Poland*
- ⁷⁷*National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India*
- ⁷⁸*National Nuclear Research Center, Baku, Azerbaijan*
- ⁷⁹*National Research and Innovation Agency - BRIN, Jakarta, Indonesia*
- ⁸⁰*Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark*
- ⁸¹*Nikhef, National institute for subatomic physics, Amsterdam, Netherlands*
- ⁸²*Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom*
- ⁸³*Nuclear Physics Institute of the Czech Academy of Sciences, Husinec-Řež, Czech Republic*
- ⁸⁴*Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA*
- ⁸⁵*Ohio State University, Columbus, Ohio, USA*
- ⁸⁶*Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia*
- ⁸⁷*Physics Department, Panjab University, Chandigarh, India*
- ⁸⁸*Physics Department, University of Jammu, Jammu, India*
- ⁸⁹*Physics Program and International Institute for Sustainability with Knotted Chiral Meta Matter (WPI-SKCM²), Hiroshima University, Hiroshima, Japan*
- ⁹⁰*Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany*
- ⁹¹*Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- ⁹²*Physik Department, Technische Universität München, Munich, Germany*
- ⁹³*Politecnico di Bari and Sezione INFN, Bari, Italy*
- ⁹⁴*Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany*
- ⁹⁵*Saga University, Saga, Japan*
- ⁹⁶*Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India*
- ⁹⁷*School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom*
- ⁹⁸*Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru*
- ⁹⁹*SUBATECH, IMT Atlantique, Nantes Université, CNRS-IN2P3, Nantes, France*
- ¹⁰⁰*Sungkyunkwan University, Suwon City, Republic of Korea*
- ¹⁰¹*Suranaree University of Technology, Nakhon Ratchasima, Thailand*
- ¹⁰²*Technical University of Košice, Košice, Slovak Republic*
- ¹⁰³*The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland*
- ¹⁰⁴*The University of Texas at Austin, Austin, Texas, USA*
- ¹⁰⁵*Universidad Autónoma de Sinaloa, Culiacán, Mexico*

- ¹⁰⁶*Universidade de São Paulo (USP), São Paulo, Brazil*
¹⁰⁷*Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil*
¹⁰⁸*Universidade Federal do ABC, Santo Andre, Brazil*
¹⁰⁹*Universitatea Nationala de Stiinta si Tehnologie Politehnica Bucuresti, Bucharest, Romania*
¹¹⁰*University of Cape Town, Cape Town, South Africa*
¹¹¹*University of Derby, Derby, United Kingdom*
¹¹²*University of Houston, Houston, Texas, USA*
¹¹³*University of Jyväskylä, Jyväskylä, Finland*
¹¹⁴*University of Kansas, Lawrence, Kansas, USA*
¹¹⁵*University of Liverpool, Liverpool, United Kingdom*
¹¹⁶*University of Science and Technology of China, Hefei, China*
¹¹⁷*University of Silesia in Katowice, Katowice, Poland*
¹¹⁸*University of South-Eastern Norway, Kongsberg, Norway*
¹¹⁹*University of Tennessee, Knoxville, Tennessee, USA*
¹²⁰*University of the Witwatersrand, Johannesburg, South Africa*
¹²¹*University of Tokyo, Tokyo, Japan*
¹²²*University of Tsukuba, Tsukuba, Japan*
¹²³*Universität Münster, Institut für Kernphysik, Münster, Germany*
¹²⁴*Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*
¹²⁵*Université de Lyon, CNRS/IN2P3, Institut de Physique des 2 Infinis de Lyon, Lyon, France*
¹²⁶*Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France, Strasbourg, France*
¹²⁷*Université Paris-Saclay, Centre d'Etudes de Saclay (CEA), IRFU, Département de Physique Nucléaire (DPhN), Saclay, France*
¹²⁸*Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France*
¹²⁹*Università degli Studi di Foggia, Foggia, Italy*
¹³⁰*Università del Piemonte Orientale, Vercelli, Italy*
¹³¹*Università di Brescia, Brescia, Italy*
¹³²*Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India*
¹³³*Warsaw University of Technology, Warsaw, Poland*
¹³⁴*Wayne State University, Detroit, Michigan, USA*
¹³⁵*Yale University, New Haven, Connecticut, USA*
¹³⁶*Yildiz Technical University, Istanbul, Turkey*
¹³⁷*Yonsei University, Seoul, Republic of Korea*
¹³⁸*Affiliated with an institute formerly covered by a cooperation agreement with CERN*
¹³⁹*Affiliated with an international laboratory covered by a cooperation agreement with CERN*

* Also at INFN Trieste, Trieste, Italy.

† Deceased.

‡ Also at Fondazione Bruno Kessler (FBK), Trento, Italy.

§ Also at Max-Planck-Institut für Physik, Munich, Germany.

|| Also at Czech Technical University in Prague, Prague, Czech Republic.

¶ Also at Instituto de Física da Universidade de São Paulo.

** Also at Dipartimento DET del Politecnico di Torino, Turin, Italy.

†† Also at University College of Dublin, Dublin, Ireland.

‡‡ Also at Department of Applied Physics, Aligarh Muslim University, Aligarh, India.

§§ Also at Institute of Theoretical Physics, University of Wrocław, Wrocław, Poland.

||| Also at Facultad de Ciencias, Universidad Nacional Autónoma de México, Mexico City, Mexico.