

Disentangling isospin violation in charmonia decaying into $\Xi\bar{\Xi}$ pairs

Huan-Ran Wen^{1,2}, Chun-Qiu Zhao^{1,3,*}, Xu Cao^{1,2,4,†} and Jian-Ping Dai^{3,‡}

¹State Key Laboratory of Heavy Ion Science and Technology, *Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

²School of Nuclear Sciences and Technology, *University of Chinese Academy of Sciences, Beijing 100049, China*

³Department of Physics, *Yunnan University, Kunming 650091, China*

⁴Research Center for Hadron and CSR Physics, *Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China*



(Received 24 August 2026; accepted 8 September 2026; published 28 September 2026)

Based on a model-independent isospin decomposition, we perform a data-driven analysis of J/ψ and $\psi(2S)$ decays into $\Xi\bar{\Xi}$ pairs by leveraging the full set of observables from high-statistics BESIII data. We find that the isospin-violating amplitudes are predominantly driven by the electric couplings, with magnitudes reaching approximately 10%, contrasting with the smaller magnetic contributions of about 3%. A partial-wave analysis further reveals that S -wave isospin violation is comparable to the D -wave contribution.

DOI: [10.1103/1jj9-jzz7](https://doi.org/10.1103/1jj9-jzz7)

I. INTRODUCTION

The decays of charmonium states into baryon and their antiparticles are of particular interest from both experimental and theoretical perspectives. A systematic description of those exclusive processes can be achieved using a perturbative approach within the QCD factorization framework [1–3]. However, final-state rescattering interactions are essential to accurately account for the experimental data [4,5]. A large relative phase, nearly orthogonal, is found between the strong and electromagnetic amplitudes for both J/ψ and $\psi(2S)$ decays [6]. These processes can also be used to test the gluon spin [7], as well as to investigate constituent quark mass effects [8]. Regarding the testing of $SU(3)$ flavor symmetry [9–12], the breaking effect is found to be less significant for $\psi(2S)$ than for J/ψ [13,14]. Even more intriguing is the mechanism of isospin violation stemming from up and down quark mass differences and electromagnetic effects, which remains a key issue in understanding these decays. The isospin violations due to quark masses is expected to be much smaller than $SU(3)$ violations predominantly due to the strange-quark mass. Early on, it was pointed out that isospin violations from

electromagnetic corrections could be significant and depend sensitively on the baryon magnetic form factors [15].

Noticeable isospin violations are not unique to baryonic final states but are also manifest in other exclusive channels. A notable example is $J/\psi \rightarrow K^*(892)\bar{K}$, where a very pronounced isospin breaking from the electromagnetic interaction was reported [16]. The BESIII Collaboration has measured precisely the branching fraction for the isospin-violating decay $\psi(3770) \rightarrow J/\psi\pi^0$ [17,18], thereby enabling a reliable extraction of the light-quark mass ratio. Further searches were conducted for the isospin-violating transitions $\chi_{c0,2} \rightarrow \pi^0\eta_c$ [19] and doubly Okubo-Zweig-Iizuka-suppressed decay $J/\psi \rightarrow \phi\pi^0$ [20]. The G -parity violating decay $J/\psi \rightarrow \pi^+\pi^-$ attracts interest as the one-photon exchange process is expected to dominate over other amplitudes that involve gluons and are hence suppressed [21].

The study of hyperons is especially advantageous because they act as self-analyzing polarimeters through their weak decays [22]. By measuring the polarization and correlations of the hyperon-antihyperon system, one can unambiguously determine the relative phase between the electric and magnetic form factors [23–28]. The isospin-violating decay $\psi \rightarrow \Sigma^0\bar{\Lambda} + c.c.$, which is mediated by a purely electromagnetic mechanism and results in a branching fraction about two orders of magnitude lower than those of isospin-allowed channels [29–32], provides a reference scale for the magnitude of isospin-violating amplitudes as well as a unique opportunity to study two-photon exchange effects in the timelike region [33]. This work presents a

* Contact author: zhaochunqiu@stu.ynu.edu.cn

† Contact author: caoxu@impcas.ac.cn

‡ Contact author: daijianping@ynu.edu.cn

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

model-independent separation of the electric and magnetic isospin-violating amplitudes in Sec. II, and equivalently their S - and D -wave components in Sec. III, based on the full experimental information from the sequential decays of J/ψ and $\psi(2S)$ into $\Xi\bar{\Xi}$ pairs.

II. ISOSPIN DECOMPOSITION OF AMPLITUDES

It is feasible to perform a model-independent isospin decomposition of the timelike electromagnetic form factors for baryon isospin multiplets by transforming from the physical basis to the isospin basis [34,35]. Similarly, the electromagnetic couplings in charmonium decays into baryon-antibaryon pairs can be decomposed into isospin components, where the isovector component accounts for isospin violation. The isospin decomposition for an isospin-doublet baryon pairs involves four independent complex amplitudes (2 isospin states $\times$ 2 electromagnetic couplings), yielding seven relevant real parameters after fixing a global phase. These parameters are solely determined by eight independent observables available at each energy point: the decay rates and angular distributions, along with the relative phases between electric and magnetic couplings derived from polarization observables for both members of the isospin doublet. While applicable to both the nucleon and Ξ isospin doublets, the latter is particularly advantageous as all relevant observables are accessible with high precision through self-analyzing weak decays. This facilitates a purely data-driven investigation of isospin violated amplitudes of charmonium, as detailed below.

We first recall the basic formula for the decay of a vector charmonium into an octet baryon-antibaryon pair $Y\bar{Y}$, where the angular distribution is given as

$$\frac{d\Gamma_Y}{d\cos\theta} = \frac{M_\psi\beta}{32\pi} \left(|G_M|^2 + \frac{|G_E|^2}{\tau} \right) (1 + \alpha_Y \cos^2\theta), \quad (1)$$

where θ denotes the angle between the baryon momentum and the positron beam direction in the center-of-mass (c.m.) frame. $\beta = \sqrt{1 - \tau^{-1}}$ is the velocity of the baryon with $\tau = M_\psi^2/4M_Y^2$. The angular distribution parameter α_Y is given by $\alpha_Y = (\tau - R_Y^2)/(\tau + R_Y^2)$ with $R = |G_E/G_M|$. The partial decay branching ratio is

$$\begin{aligned} \mathcal{B}_{\psi \rightarrow Y\bar{Y}} &= \frac{M_\psi\beta}{12\pi\Gamma_\psi} \left(|G_M|^2 + \frac{|G_E|^2}{2\tau} \right) \\ &\equiv \frac{M_\psi\beta}{12\pi\Gamma_\psi} \frac{2\tau + 1}{2\tau} |G_{\text{eff}}|^2 \end{aligned} \quad (2)$$

with Γ_ψ being the total width of charmonium. Thus, simultaneous measurements of the decay rates and angular distributions unambiguously separate the moduli of the electric and magnetic couplings. In the case of unpolarized electron and positron beams, the transverse polarization of

the hyperon and the spin correlation of hyperon-antihyperon are

$$\begin{aligned} P_y &= \frac{\sin 2\theta}{\sqrt{\tau}D} \Im m(G_M G_E^*) \\ &= \frac{\sqrt{1 - \alpha_\psi^2} \sin\theta \cos\theta}{1 + \alpha_\psi \cos^2\theta} \sin\Delta\Phi, \end{aligned} \quad (3)$$

$$\begin{aligned} C_{xz} &= \frac{\sin 2\theta}{\sqrt{\tau}D} \Re e(G_M G_E^*) \\ &= \frac{\sqrt{1 - \alpha_\psi^2} \sin\theta \cos\theta}{1 + \alpha_\psi \cos^2\theta} \cos\Delta\Phi, \end{aligned} \quad (4)$$

with $D = (1 + \cos^2\theta)|G_M|^2 + \frac{1}{\tau}\sin^2\theta|G_E|^2$. Those polarized observables can be reconstructed via the sequential decay chains of the hyperons [23,25,36]. Consequently, the relative phases $\Delta\Phi$ between the electromagnetic couplings can be unambiguously determined, providing the necessary input for the isospin analysis.

We then perform an isospin decomposition of the electromagnetic couplings into their isoscalar ($I = 0$) and isovector ($I = 1$) components [35]:

$$G_{E,M}^{+,-} = \frac{G_{E,M}^1 \pm G_{E,M}^0}{2} \quad (5)$$

where $+$ and $-$ correspond to $\psi \rightarrow \Xi^0\bar{\Xi}^0$ and $\Xi^-\bar{\Xi}^+$ channels, respectively. The complex amplitudes are parametrized, without loss of generality, as $G_{E,M}^I = |G_{E,M}^I|e^{i\phi_{E,M}^I}$. The strength and relative phase of the isospin-violating effect is characterized by

$$\delta_{E,M} = \left| \frac{G_{E,M}^1}{G_{E,M}^0} \right|, \quad \phi_{E,M} = \arg \left[\frac{G_{E,M}^1}{G_{E,M}^0} \right], \quad (6)$$

for electric and magnetic couplings, respectively. Here the relative phases between the isovector and isoscalar components are defined as $\phi_{E,M} = \phi_{E,M}^1 - \phi_{E,M}^0$. Since charmonium states are isospin singlets, any nonvanishing value of $\delta_{E,M}$ quantifies the degree of isospin violation in the decay process, with $\delta_{E,M} \ll 1$ being generally expected. Eq. (5) yields a practical ratio to bound isospin violation [34]:

$$R_I^{E,M} = \frac{|G_{E,M}^+|^2 - |G_{E,M}^-|^2}{|G_{E,M}^+|^2 + |G_{E,M}^-|^2} = \frac{2\delta_{E,M} \cos\phi_{E,M}}{1 + \delta_{E,M}^2} \quad (7)$$

if the electric and magnetic couplings $|G_{E,M}^\pm|$ for both $\psi \rightarrow \Xi^0\bar{\Xi}^0$ and $\Xi^-\bar{\Xi}^+$ channels are individually measured.

Using Eqs. (2) and (5), the isoscalar and isovector components can be conveniently expressed in terms of the unpolarized observables, $|G_{\text{eff}}|$ and $R_\pm$ [34]:

TABLE I. The isospin violation relevant ratio, R_I^M , and R_I^E extracted from the branching ratios and the α_Y . Earlier measurements are listed for reference but are now regarded as superseded.

Decay process	Branching ratio	α_Y	$\Delta\Phi/\text{rad}$	$ R_I^M $	$ R_I^E $
$J/\psi \rightarrow \Xi^0 \bar{\Xi}^0$	$(1.17 \pm 0.04) \times 10^{-3}$ [37]	$0.514 \pm 0.006 \pm .0015$ [38] ^a	$1.168 \pm 0.019 \pm 0.018$ [38]	0.08 ± 0.04	0.18 ± 0.05
$J/\psi \rightarrow \Xi^- \bar{\Xi}^+$	$(0.97 \pm 0.08) \times 10^{-3}$ [37]	$0.586 \pm 0.012 \pm 0.010$ [27] ^b	$1.213 \pm 0.046 \pm 0.016$ [27]		
$\psi(2S) \rightarrow \Xi^0 \bar{\Xi}^0$	$(2.59 \pm 0.02) \times 10^{-4}$ [39] ^c	$0.768 \pm 0.029 \pm 0.025$ [39] ^d	$0.257 \pm 0.061 \pm 0.009$ [39] ^e	0.040 ± 0.032	0.19 ± 0.16
$\psi(2S) \rightarrow \Xi^- \bar{\Xi}^+$	$(2.87 \pm 0.11) \times 10^{-4}$ [37]	$0.693 \pm 0.048 \pm 0.049$ [40] ^f	$0.667 \pm 0.111 \pm 0.058$ [40]		

^a $0.66 \pm 0.03 \pm 0.05$ [41] is superseded.

^b $0.58 \pm 0.04 \pm 0.08$ [42] is superseded.

^c $(2.3 \pm 0.4) \times 10^{-4}$ [37] is superseded.

^d $0.65 \pm 0.09 \pm 0.14$ [41] and $0.665 \pm 0.086 \pm 0.081$ [43] are superseded.

^e $-0.05 \pm 0.150 \pm 0.020$ [43] is superseded.

^f $0.91 \pm 0.13 \pm 0.14$ [42] is superseded.

$$|G_M^0|^2 + |G_M^1|^2 = \frac{2(2\tau_+ + 1)|G_{\text{eff}}^+|^2}{2\tau_+ + R_+^2} + \frac{2(2\tau_- + 1)|G_{\text{eff}}^-|^2}{2\tau_- + R_-^2}, \quad (8)$$

$$|G_E^0|^2 + |G_E^1|^2 = \frac{2(2\tau_+ + 1)R_+^2|G_{\text{eff}}^+|^2}{2\tau_+ + R_+^2} + \frac{2(2\tau_- + 1)R_-^2|G_{\text{eff}}^-|^2}{2\tau_- + R_-^2}, \quad (9)$$

$$\Re[G_M^1 G_M^{0\dagger}] = \frac{(2\tau_+ + 1)|G_{\text{eff}}^+|^2}{2\tau_+ + R_+^2} - \frac{(2\tau_- + 1)|G_{\text{eff}}^-|^2}{2\tau_- + R_-^2}, \quad (10)$$

$$\Re[G_E^1 G_E^{0\dagger}] = \frac{(2\tau_+ + 1)R_+^2|G_{\text{eff}}^+|^2}{2\tau_+ + R_+^2} - \frac{(2\tau_- + 1)R_-^2|G_{\text{eff}}^-|^2}{2\tau_- + R_-^2}, \quad (11)$$

Here, $\tau_{\pm}$ takes into account the small isospin-breaking effects arising from the $\Xi^{0,-}$ mass difference. Achieving a complete separation of the electric and magnetic couplings in the isospin basis requires the knowledge of the relative phases $\Delta\Phi$ of both channels. This is achieved by substituting Eq. (5) into the polarization observables in Eqs. (3) and (4), yielding

$$\Re[G_M^1 G_E^{1*} + G_M^0 G_E^{0*}] = 2|G_M^+||G_E^+| \cos \Delta\Phi^+ + 2|G_M^-||G_E^-| \cos \Delta\Phi^- \quad (12)$$

$$\Re[G_M^1 G_E^{0*} + G_M^0 G_E^{1*}] = 2|G_M^+||G_E^+| \cos \Delta\Phi^+ - 2|G_M^-||G_E^-| \cos \Delta\Phi^- \quad (13)$$

$$\Im[G_M^1 G_E^{1*} + G_M^0 G_E^{0*}] = 2|G_M^+||G_E^+| \sin \Delta\Phi^+ + 2|G_M^-||G_E^-| \sin \Delta\Phi^- \quad (14)$$

$$\Im[G_M^1 G_E^{0*} + G_M^0 G_E^{1*}] = 2|G_M^+||G_E^+| \sin \Delta\Phi^+ - 2|G_M^-||G_E^-| \sin \Delta\Phi^- \quad (15)$$

An alternative expression in terms of the polarization observables P_y and C_{xz} can be found in Appendix A of Ref. [34].¹ This complete set of relations in Eqs. (8)–(15) enables the separation of the electric and magnetic couplings in the isospin basis ($I = 0, 1$), transformed from the physical basis (Ξ^0, Ξ^-).

Two subtleties should be noted when applying them to practical use. First, the global phase is arbitrary, and we fix $\varphi_M^0 = 0$ by convention. Consequently, all other phases are

defined relative to φ_M^0 . Second, although the equations are analytical, finding a numerical solution is not straightforward, especially when considering the uncertainties associated with the measured observables. We therefore use the ratio $R_I^{E,M}$ in Eq. (7) to determine the allowed boundaries for the relative magnitudes and phases in the isospin basis. The extracted values of $R_I^{E,M}$ and their uncertainties, as listed in Table I, determine the allowed regions for the relative magnitudes $\delta_{E,M}$ and phases $\phi_{E,M}$, which are illustrated in Figs. 1 and 2. Consistent with the expectation that isospin violation is small, we restrict the plot to $\delta_{E,M} \leq 1$. The blue band representing R_I^E lies above the orange band for R_I^M , reflecting the observed hierarchy

¹We note that there are typographical errors in Eqs. (A.6) and (A.7) of Ref. [34], where $\Re$ and $\Im$ should be interchanged.




FIG. 1. Allowed regions for $\delta_{E,M}$ and $\phi_{E,M}$ in $J/\psi \rightarrow \Xi\bar{\Xi}$ decays. The blue and orange bands represent constraints derived from the R_I^E and R_I^M values listed in Table I, respectively. The points with error bars denote the numerical solutions obtained from the χ^2 minimization.




FIG. 2. Same as Fig. 1 but for $\psi(2S) \rightarrow \Xi\bar{\Xi}$ decay.

$R_I^E > R_I^M$.² For the $J/\psi \rightarrow \Xi\bar{\Xi}$ decay, the data provide unambiguous significance for this relation. However, for $\psi(2S) \rightarrow \Xi\bar{\Xi}$, the two bands exhibit some overlap, providing only marginal evidence. Future updated measurements of the branching ratios would be essential to clarify this situation. Within the constraints of the allowed parameter space, a χ^2 minimization is subsequently employed to obtain numerical solutions to Eqs. (8)–(15) within the allowed parameter space. Bootstrap resampling with 2000

²Previous BESIII data with lower precision suggested the opposite hierarchy, $R_I^E < R_I^M$, in both channels (see Table I and Ref. [34]), underscoring the crucial importance of high-precision data for such investigations.

TABLE II. The isoscalar $G_{E,M}^0$ and isovector $G_{E,M}^1$ couplings and their phases (in rad) for J/ψ and $\psi(2S)$ decaying into $\Xi\bar{\Xi}$.

channel	$J/\psi \rightarrow \Xi\bar{\Xi}$	$\psi(2S) \rightarrow \Xi\bar{\Xi}$
$ G_M^0 $	$(2.77 \pm 0.066) \times 10^{-3}$	$(2.06 \pm 0.043) \times 10^{-3}$
$ G_M^1 $	$(7.91 \pm 6.01) \times 10^{-5}$	$(6.21 \pm 3.05) \times 10^{-5}$
$ G_E^0 $	$(1.39 \pm 0.038) \times 10^{-3}$	$(8.09 \pm 1.01) \times 10^{-4}$
$ G_E^1 $	$(1.17 \pm 0.41) \times 10^{-4}$	$(8.66 \pm 3.02) \times 10^{-5}$
$\phi_M^1 - \phi_M^0$	-0.0053 ± 0.29	-0.0015 ± 0.0008
$\phi_E^0 - \phi_M^0$	-1.19 ± 0.027	-0.36 ± 0.10
$\phi_E^1 - \phi_M^0$	-0.95 ± 0.55	0.0025 ± 0.0012
δ_E	$0.084^{+0.030}_{-0.013}$	0.11 ± 0.040
δ_M	$0.029^{+0.022}_{-0.013}$	$0.030^{+0.006}_{-0.015}$

iterations is used to propagate the uncertainties of the observables. The resulting numerical solutions, including their uncertainties, are listed in Table II and presented as colored points with error bars in Figs. 1 and 2. As our primary conclusion, evidence for $\delta_E > \delta_M$ is observed in the J/ψ channel at the 2σ significance level, with similar indications found in the $\psi(2S)$ channel. The isospin-violating magnitudes for the electric and magnetic couplings are approximately 10% and 3%, respectively, for both charmonium states. The relative phase ϕ_M is found to be consistent with zero, while the relative phase ϕ_E shows a significant deviation from zero in both decay channels. Interestingly, the value of $\phi_E^1 - \phi_M^0$ for $\psi(2S)$ remains consistent with zero, whereas that for J/ψ deviates substantially from zero, suggesting distinct decay mechanisms for the two charmonium states.

III. S- AND D- PARTIAL WAVES

The studied amplitudes can be alternatively described using various formalisms, such as helicity amplitudes [44,45], covariant tensor formalisms [46], or by considering Dirac and Pauli couplings [47]. In this section, we focus on an alternative decomposition in terms of S- and D-partial waves, which is equivalent to the electromagnetic form factor representation [46,48–51]

$$G_M = g_s + \frac{1}{\sqrt{2}}g_d \quad (16)$$

$$\frac{G_E}{\sqrt{\tau}} = g_s - \sqrt{2}g_d \quad (17)$$

Note that we adopt a normalization convention different from that in the literature, chosen to express the partial branching fractions in a more transparent manner:

$$\mathcal{B}_{\psi \rightarrow Y\bar{Y}} = \frac{M_\psi \beta}{8\pi\Gamma_\psi} (|g_s|^2 + |g_d|^2), \quad (18)$$




FIG. 3. Allowed regions for $\delta_{s,d}$ and $\phi_{s,d}$ in $J/\psi \rightarrow \Xi\bar{\Xi}$ decays. The blue and orange bands represent constraints derived from the R_I^S and R_I^D , respectively. The points with error bars denote the numerical solutions obtained from the χ^2 minimization.

In the isospin partial-wave basis, the couplings are typically written as

$$g_{s,d}^{1,0} = \frac{g_{s,d}^+ \pm g_{s,d}^-}{2} \quad (19)$$

where the upper sign 1 (or 0) applies to the isovector (or isoscalar) case. $g_{s,d}^+$ (or $g_{s,d}^-$) are S -wave and D -wave coupling constants of Ξ^0 (or Ξ^-), respectively. Consequently, the left-hand sides of Eqs. (8)–(15) can be rewritten using $g_{s,d}^\pm$. The complex amplitudes are further parametrized as $g_{s,d}^I = |g_{s,d}^I|e^{i\phi_{s,d}^I}$.

Following the same procedure as in Sec. II, a partial-wave relevant ratio is introduced to bound the extent of isospin violation:




FIG. 4. Same as Fig. 3 but for $\psi(2S) \rightarrow \Xi\bar{\Xi}$ decay.

TABLE III. The coupling constant and phases (in rad) of S -wave and D -wave for J/ψ and $\psi(2S)$ decaying into $\Xi\bar{\Xi}$.

Channel	$J/\psi \rightarrow \Xi\bar{\Xi}$	$\psi(2S) \rightarrow \Xi\bar{\Xi}$
$ g_s^0 $	$(1.97 \pm 0.15) \times 10^{-3}$	$(15.04 \pm 0.28) \times 10^{-4}$
$ g_s^1 $	$(1.06 \pm 3.99) \times 10^{-4}$	$(6.72 \pm 1.27) \times 10^{-5}$
$ g_d^0 $	$(12.07 \pm 0.64) \times 10^{-4}$	$(7.99 \pm 0.28) \times 10^{-4}$
$ g_d^1 $	$(2.14 \pm 18.21) \times 10^{-5}$	$(2.77 \pm 1.14) \times 10^{-5}$
$\phi_s^1 - \phi_s^0$	0.71 ± 1.03	-1.24 ± 0.34
$\phi_d^0 - \phi_s^0$	-1.78 ± 1.19	-1.18 ± 0.05
$\phi_d^1 - \phi_s^0$	0.98 ± 1.12	-1.90 ± 1.21
δ_s	$0.054^{+0.034}_{-0.014}$	0.045 ± 0.010
δ_d	$0.018^{+0.046}_{-0.017}$	0.035 ± 0.018

$$R_I^{S,D} = \frac{|g_{s,d}^+|^2 - |g_{s,d}^-|^2}{|g_{s,d}^+|^2 + |g_{s,d}^-|^2} = \frac{2\delta_{s,d} \cos \phi_{s,d}}{1 + \delta_{s,d}^2} \quad (20)$$

where $\delta_{s,d} = |g_{s,d}^1/g_{s,d}^0|$ is relative magnitude and $\phi_{s,d} = \phi_{s,d}^1 - \phi_{s,d}^0$ is relative phase between isovector and isoscalar couplings. The allowed regions for these parameters are shown in Figs. 3 and 4, where the final solutions from the χ^2 minimization are indicated by colored points with error bars. The corresponding numerical results are summarized in Table III. To account for the arbitrary global phase, ϕ_s^0 is fixed at zero during the numerical search.

At current precision, the data suggest that isospin violation in the S -wave is of a similar magnitude to that in the D -wave. The large errors in the extracted phases preclude any definitive conclusions at this stage, highlighting the need for more precise data in future experiments.

IV. SUMMARY AND PERSPECTIVE

As established in earlier work [34], the relative magnitudes of the isoscalar and isovector components in baryon-antibaryon pairs produced via electron-positron annihilation can be unambiguously determined using a complete set of observables from the relevant isospin channels. Although this framework has been applied to nucleon and hyperon electromagnetic form factors through differential cross-section analysis, and to charmonium decays by incorporating isospin violation, previous analyses were limited by large uncertainties and the absence of polarization observables. With the accumulation of high-statistics data and reduced uncertainties, it is now possible to perform a more accurate, model-independent analysis based on the isospin decomposition of vector charmonium decays into hyperon-antihyperon pairs. This paper is devoted to outlining the full theoretical framework for isospin decomposition in terms of electromagnetic couplings and applying it to the available experimental data for J/ψ and $\psi(2S)$ decays into $\Xi\bar{\Xi}$ pairs.

Our analysis demonstrates that the isospin violations are predominantly driven by the electric couplings (approximately 10%), whereas the magnetic contributions are significantly smaller (about 3%). This finding is particularly interesting, as the QCD factorization framework previously predicted that isospin violations would be more sensitive to the baryon magnetic form factors [15]. The isospin violations in the S - and D -wave components are of comparable magnitude, each at the level of several percent. Our results contribute to the understanding of the decay mechanisms of charmonia into baryon-antibaryon pairs, while potentially offering insights into the quantum entanglement inherent in these systems [52–54].

Our framework is well-suited for the isospin decomposition of nucleon-antinucleon system [55–57]. For vector charmonia decay, preliminary analysis indicates that isospin violation in the electric coupling is more pronounced [34]. By utilizing recently developed techniques [58,59] to measure proton polarization, one can effectively validate the current conclusions and provide more stringent constraints on the isospin-violating amplitudes. In contrast, measuring $\Delta\Phi$ for nucleon electromagnetic form factors at any energies with high precision is more challenging due to the high statistics required by polarization measurements. Alternatively, dispersion-theoretic approaches that globally

fit both spacelike and timelike data provide valuable information on these phases [60,61].

Despite the increased complexity of the decomposition for isospin triplets, our framework is applicable to the $J/\psi, \psi(2S) \rightarrow \Sigma^\pm \bar{\Sigma}^\mp$ final states [62–67]. Nevertheless, a complete isospin analysis is currently hindered by the lack of experimental data for the $\Sigma^- \bar{\Sigma}^+$ channel [68]. The potentially more pronounced isospin-violating effects in these channels, estimated at approximately 30% [34], merit further attention.

ACKNOWLEDGMENTS

We are grateful to Xiongfei Wang for useful discussions. This work is supported by the National Natural Science Foundation of China (Grants No. 12547111 and No. 12165022), the National Key R&D Program of China under Grant No. 2023YFA1606703, and Yunnan Fundamental Research Project under Contract No. 202301AT070162.

DATA AVAILABILITY

There are no publicly available research data or software supporting this manuscript. Requests for further information or data should be sent to the authors.

-
- [1] J. Bolz and P. Kroll, *Eur. Phys. J. C* **2**, 545 (1998).
 - [2] N. Kivel, *Eur. Phys. J. A* **58**, 26 (2022).
 - [3] N. Kivel, *Eur. Phys. J. A* **58**, 138 (2022).
 - [4] H. Chen and R. G. Ping, *Phys. Lett. B* **644**, 54 (2007).
 - [5] C.-Q. Geng, X.-N. Jin, and C.-W. Liu, *arXiv:2607.00590*.
 - [6] K. Zhu, X.-H. Mo, and C.-Z. Yuan, *Int. J. Mod. Phys. A* **30**, 1550148 (2015).
 - [7] S. J. Brodsky and G. P. Lepage, *Phys. Rev. D* **24**, 2848 (1981).
 - [8] C. Carimalo, *Int. J. Mod. Phys. A* **02**, 249 (1987).
 - [9] G. López Castro, J. L. Lucio M., and J. Pestieau, *AIP Conf. Proc.* **342**, 441 (1995).
 - [10] D.-H. Wei, *J. Phys. G* **36**, 115006 (2009).
 - [11] X. H. Mo and J. Y. Zhang, *Phys. Lett. B* **826**, 136927 (2022).
 - [12] X. H. Mo, P. Wang, and J. Y. Zhang, *Phys. Rev. D* **107**, 094009 (2023).
 - [13] R. Baldini Ferroli, A. Mangoni, S. Pacetti, and K. Zhu, *Phys. Lett. B* **799**, 135041 (2019).
 - [14] R. B. Ferroli, A. Mangoni, S. Pacetti, and K. Zhu, *Phys. Rev. D* **103**, 016005 (2021).
 - [15] M. Claudson, S. L. Glashow, and M. B. Wise, *Phys. Rev. D* **25**, 1345 (1982).
 - [16] M. Ablikim *et al.* (BESIII Collaboration), *arXiv:2606.12002*.
 - [17] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **86**, 092008 (2012).
 - [18] M. Ablikim *et al.* (BESIII Collaboration), *arXiv:2606.14105*.
 - [19] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **91**, 112018 (2015).
 - [20] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **91**, 112001 (2015).
 - [21] R. Baldini Ferroli, A. Mangoni, and S. Pacetti, *Phys. Rev. C* **98**, 045210 (2018).
 - [22] T. D. Lee and C.-N. Yang, *Phys. Rev.* **108**, 1645 (1957).
 - [23] G. Fäldt and A. Kupsc, *Phys. Lett. B* **772**, 16 (2017).
 - [24] N. Salone, P. Adlarson, V. Batozskaya, A. Kupsc, S. Leupold, and J. Tandean, *Phys. Rev. D* **105**, 116022 (2022).
 - [25] X. Cao, Y.-T. Liang, and R.-G. Ping, *Phys. Rev. D* **110**, 014035 (2024).
 - [26] M. Ablikim *et al.* (BESIII Collaboration), *Nat. Phys.* **15**, 631 (2019).
 - [27] M. Ablikim *et al.* (BESIII Collaboration), *Nature (London)* **606**, 64 (2022).
 - [28] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **133**, 101902 (2024).
 - [29] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **86**, 032008 (2012).
 - [30] M. Ablikim *et al.* (BESIII Collaboration), *Nat. Commun.* **15**, 8812 (2024).

- [31] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **103**, 112004 (2021).
- [32] F. Takahashi *et al.* (Particle Data Group Collaboration), *Int. J. Mod. Phys. A* **41**, 2630011 (2026).
- [33] X. Cao, Y.-H. Lin, and H.-Q. Zhou, *arXiv:2609.19652*.
- [34] J.-P. Dai, X. Cao, and H. Lenske, *Phys. Lett. B* **846**, 138192 (2023).
- [35] X. Cao, J.-P. Dai, and H. Lenske, *Phys. Rev. D* **105**, L071503 (2022).
- [36] E. Tomasi-Gustafsson, F. Lacroix, C. Duterte, and G.I. Gakh, *Eur. Phys. J. A* **24**, 419 (2005).
- [37] R. L. Workman *et al.* (Particle Data Group Collaboration), *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [38] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **108**, L031106 (2023).
- [39] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **113**, L091501 (2026).
- [40] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **106**, L091101 (2022).
- [41] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Lett. B* **770**, 217 (2017).
- [42] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **93**, 072003 (2016).
- [43] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **108**, L011101 (2023).
- [44] N. H. Buttimore and E. Jennings, *Eur. Phys. J. A* **31**, 9 (2007).
- [45] N. H. Buttimore and E. Jennings, *Eur. Phys. J. A* **33**, 21 (2007).
- [46] S.-M. Wu, J.-J. Wu, and B.-S. Zou, *Phys. Rev. D* **104**, 054018 (2021).
- [47] T. Barnes, X. Li, and W. Roberts, *Phys. Rev. D* **77**, 056001 (2008).
- [48] J. Haidenbauer, U.-G. Meißner, and L.-Y. Dai, *Phys. Rev. D* **103**, 014028 (2021).
- [49] Q.-H. Yang, D. Guo, L.-Y. Dai, J. Haidenbauer, X.-W. Kang, and U.-G. Meißner, *Sci. Bull.* **68**, 2729 (2023).
- [50] Q.-H. Yang, D. Guo, M.-Y. Li, L.-Y. Dai, J. Haidenbauer, and U.-G. Meißner, *J. High Energy Phys.* **08** (2024) 208.
- [51] L.-Y. Dai, J. Haidenbauer, and U.-G. Meißner, *Chin. Phys. Lett.* **42**, 030202 (2025).
- [52] C. Li, X. Cao, A.-Q. Guo, C.-X. Yu, H.-W. Zhang, and Z. Zhang, *Chin. Phys. C* **50**, 093106 (2026).
- [53] H.-W. Zhang, X. Cao, and T.-F. Feng, *arXiv:2605.19642*.
- [54] H.-W. Zhang, X. Cao, and T.-F. Feng, *Phys. Rev. D* **113**, 114022 (2026).
- [55] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **86**, 032014 (2012).
- [56] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **98**, 032006 (2018).
- [57] M. Ablikim *et al.* (BESIII Collaboration), *arXiv:2608.15074*.
- [58] M. Ablikim *et al.* (BESIII Collaboration), *arXiv:2607.19927*.
- [59] M. Ablikim *et al.* (BESIII Collaboration), *arXiv:2607.19948*.
- [60] Y.-H. Lin, H.-W. Hammer, and U.-G. Meißner, *Eur. Phys. J. A* **57**, 255 (2021).
- [61] Y.-H. Lin, H.-W. Hammer, and U.-G. Meißner, *Phys. Rev. Lett.* **128**, 052002 (2022).
- [62] M. Ablikim *et al.* (BES Collaboration), *Phys. Rev. D* **78**, 092005 (2008).
- [63] M. Ablikim *et al.* (BESIII Collaboration), *J. High Energy Phys.* **11** (2021) 226.
- [64] M. Ablikim *et al.* (BESIII Collaboration), *J. High Energy Phys.* **12** (2024) 186.
- [65] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **135**, 141804 (2025).
- [66] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **125**, 052004 (2020).
- [67] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. Lett.* **131**, 191802 (2023).
- [68] M. Ablikim *et al.* (BESIII Collaboration), *J. High Energy Phys.* **12** (2022) 016.