

Three-gluon decays of radially excited quarkonia $\psi(2S)$ and $\Upsilon(2S)$ with both relativistic and QCD radiative corrections

Chao-Jie Fan[✉] and Jun-Kang He^{✉*}

*College of Physics and Electronic Science, Hubei Normal University, Huangshi 435002, China
and Key Laboratory of Quark and Lepton Physics (MOE), Central China Normal University,
Wuhan 430079, China*



(Received 6 April 2026; accepted 12 June 2026; published 6 July 2026)

For the radially excited heavy quarkonia $V = \psi(2S)$ and $\Upsilon(2S)$, the nodal structure of the wave function renders the three-gluon decay $V \rightarrow ggg$ acutely sensitive to relativistic corrections, a longstanding challenge for reliable theoretical predictions. Within the Bethe-Salpeter formalism under the covariant instantaneous ansatz, we construct analytic harmonic-oscillator wave functions incorporating the $2S$ node and derive model-independent relations among the polarized decay widths from helicity-flip and phase-space symmetries. Motivated by the strikingly slow $\hat{q}^2$ -order convergence driven by destructive interference at the node, we introduce a concise phenomenological treatment of the higher-order contributions that preserves the correct low-momentum limit. Including both relativistic and QCD radiative corrections, our predictions for $\Gamma(V \rightarrow ggg)$, $\Gamma(V \rightarrow e^+e^-)$, and R_V agree well with experiment, and the extracted β_V lies at the lower end of typical phenomenological ranges, reflecting a more localized momentum-space wave function.

DOI: [10.1103/5hm-yvk9](https://doi.org/10.1103/5hm-yvk9)

I. INTRODUCTION

Charmonium states provide a unique laboratory for testing the interplay between perturbative and nonperturbative QCD, as well as the role of relativistic effects in heavy-quark bound systems. With $\langle v^2 \rangle \approx 0.3$ for charmonium, relativistic corrections of relative order v^2 are no longer negligible and must be systematically included for precision phenomenology [1,2]. The radially excited vector state $\psi(2S)$ is of particular interest; its larger radius and the node in the radial wave function lead to enhanced relativistic dynamics and characteristic cancellations in both production and decay amplitudes, offering a sensitive probe beyond the ground state J/ψ .

Over the past two decades, relativistic corrections for $\psi(2S)$ have been systematically investigated in both production and decay processes, progressively revealing their indispensable role. Studies in diffractive photoproduction at HERA revealed that the $\sigma(\psi(2S))/\sigma(J/\psi)$ ratio is suppressed by the radial node [3,4]. The nonrelativistic QCD (NRQCD) approach has also been applied to $\psi(2S)$ production, where predictions for yields and especially

polarization observables for $\psi(2S)$ still exhibit some discrepancies with experimental data at hadron colliders [5]. In the same period, the instantaneous Salpeter method was applied to annihilation decays, demonstrating that the node structures in the $\psi(2S)$ radial wave function generate large relativistic corrections even for bottomonium, leading to a slower falloff of three-gluon and leptonic widths with radial quantum number [6]. By the late 2010s, full Bethe-Salpeter (B-S) calculations for semileptonic decays to orbitally and radially excited charmonia [including $\psi(2S)$] showed that relativistic effects can exceed 50% for excited states [7]. Lattice QCD determinations of B-S wave functions for $\psi(2S)$ further provided nonperturbative input, confirming the nodal structure [8].

In the 2020s, the field has advanced to high-precision NRQCD applications. For hadroproduction, leading-power fragmentation supplemented by v^2 corrections has been applied to high- p_T prompt $\psi(2S)$ at the LHC. Most recently, Bertone *et al.* [9] performed the first dedicated study and found that v^2 corrections increase the gluon-fragmentation cross section by factors of 3.3–6.6 (depending on $\langle v^2 \rangle = 0.25$ –0.50) while affecting the charm channel only moderately; the resulting predictions achieve excellent agreement with ATLAS and CMS data at $\sqrt{s} = 13$ TeV for $p_T \gtrsim 60$ GeV and also improve the description of polar anisotropy λ_θ . Similarly, Copeland *et al.* [10] investigated $\psi(2S)$ production inside jets at LHCb using NRQCD combined with fragmenting jet functions and a gluon fragmentation improved Pythia framework, demonstrating

*Contact author: hejk@hbnu.edu.cn

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

high sensitivity of the jet substructure variable $z = p_T(\psi(2S))/p_T(\text{jet})$ to the long-distance matrix elements and providing new constraints on the excited-state wave-function parameters. These latest works, together with earlier double-charmonium studies [11], confirm that relativistic corrections and wave-function details (including implicit nodal effects through LDMEs) are indispensable for accurate $\psi(2S)$ production phenomenology across a wide kinematic range. On the decay side, v^2 corrections to exclusive hadronic channels $\psi(nS) \rightarrow \rho\pi$ and $\gamma\pi$ were shown to play a key role in resolving the $\rho\pi$ puzzle for $\psi(2S)$ [12], while in the exclusive hadronic decay $\psi(2S) \rightarrow p\bar{p}$, the v^2 corrections with $\langle v^2 \rangle \approx 0.64$ dominate the amplitude, leading to their crucial role in understanding the decay process and a worsened convergence of the v^2 expansion [13]. Most recently, the B-S formalism has yielded complete analytic results for vector quarkonium $J/\psi \rightarrow ggg$, with relativistic corrections reaching $\sim 77\%$ and paving the way for excited-state extensions [14].

The radially excited quarkonium decay $V \rightarrow ggg$ [with $V = \psi(2S)$ or $\Upsilon(2S)$] stands out as an ideal channel. The purely gluonic final state eliminates hadronization uncertainties, allowing the nodal structure of the long-distance B-S wave function to be probed through its convolution with the hard-scattering amplitude. This process is therefore well-suited to probe v^2 -order relativistic corrections originating primarily from the initial-state quarkonium structure. Although the destructive interference induced by the radial node has been repeatedly highlighted in production studies, it has not yet been isolated using analytic B-S wave functions in the decay $V \rightarrow ggg$. A relativistic B-S calculation incorporating the node in this channel, with all eight Dirac structures of the covariant amplitude kept consistently, is therefore still missing, posing an ongoing challenge in providing reliable and precise predictions.

In this work we compute the complete q^2 -order relativistic corrections to the three-gluon decays $\psi(2S) \rightarrow ggg$ and $\Upsilon(2S) \rightarrow ggg$, by employing the analytic form of the B-S wave function for the radially excited states $\psi(2S)$ and $\Upsilon(2S)$ given in our latest work [14]. We find that the q^2 -order relativistic corrections in the $\psi(2S) \rightarrow ggg$ process are significantly larger than those in the $J/\psi \rightarrow ggg$ process, with the correction factor $\kappa\beta_V^2/M^2$ approximately equal to 0.77 for J/ψ and 2.7 for $\psi(2S)$, resulting in an unphysical negative decay width for the $\psi(2S)$ channel. To obtain a physically meaningful prediction, we adopt a phenomenological treatment of the hard kernel in the $V \rightarrow ggg$ process, as proposed in Ref. [15], which effectively incorporates higher-order (beyond q^2 -order) relativistic corrections and the nodal structure effects of the radially excited state V . And our theoretical predictions for the branching ratios $\mathcal{B}(V \rightarrow ggg)$ and $\mathcal{B}(V \rightarrow e^+e^-)$ are consistent with experimental data. Moreover, we extract the harmonic oscillator parameter β_V , which provides

valuable input for calibrating nonperturbative models of quarkonium structure.

The paper is organized as follows. The theoretical framework and the analytic calculation for the decay process $V \rightarrow ggg$ are presented in detail in Sec. II. In Sec. III we show our results and phenomenological discussions. The last section is our summary.

II. THEORETICAL FRAMEWORK

A. Bethe-Salpeter equation

The B-S equation [16,17] provides a covariant, relativistic framework for describing bound states. In this work it is implemented under the covariant instantaneous ansatz (CIA) [18–20], in which the interaction kernel is reduced to an instantaneous potential while the B-S amplitude is kept fully covariant. This implementation is particularly well-suited to the systematic treatment of radially excited quarkonia, where the nodal structure of the wave function is crucial. In this formalism, the B-S wave function $\Psi(K, q)$ encodes the complete bound-state information and is determined by the dynamical equation [14,19–24],

$$S_F^{-1}(f)\Psi(K, q)S_F^{-1}(-\bar{f}) = \int \frac{d^4 q'}{(2\pi)^4} [-i\mathcal{K}(K, q, q')\Psi(K, q')]. \quad (2.1)$$

Applying this formalism to heavy quarkonium V [with $V = \psi(2S)$ or $\Upsilon(2S)$], $\mathcal{K}(K, q, q')$ represents the interaction kernel between the internal quark and antiquark, and $S_F(p) = i/(\not{p} - \hat{m} + i\epsilon)$ is the propagator with the effective mass $\hat{m}$ of the c or b quark. The momenta of the quark and antiquark are given by

$$f = \frac{K}{2} + q, \quad \bar{f} = \frac{K}{2} - q, \quad (2.2)$$

where q and K represent the internal momentum and the total momentum of the quarkonium, respectively.

For convenience, the internal momentum q is decomposed into a transverse component $\hat{q}$ (with $\hat{q} \cdot K = 0$) and a longitudinal component $q_{\parallel}$ (parallel to the total momentum K),

$$q^\mu = q_{\parallel}^\mu + \hat{q}^\mu, \quad q_{\parallel}^\mu = \frac{q \cdot K}{M} K^\mu. \quad (2.3)$$

Here both $q_K = q \cdot K/M$ and $\hat{q}^2 = q^2 - q_K^2$ are Lorentz invariant variables and M is the mass of the heavy quarkonium.

The CIA [18,20,23,25,26] posits that the interaction is instantaneous, so that the interaction kernel $\mathcal{K}(K, q, q')$ is independent of the timelike component of the internal momentum and reduces to a function of the transverse relative momenta only. In this framework the B-S kernel is taken, in the spin-color sector, to be of one-gluon-exchange

type [19,20],

$$\mathcal{K}(K, q, q') = \left(\frac{1}{2} \vec{\lambda}_1 \cdot \frac{1}{2} \vec{\lambda}_2 \right) (\gamma_\mu \otimes \gamma^\mu) V(\hat{q} - \hat{q}'), \quad (2.4)$$

while its scalar spatial part $V(\hat{q}, \hat{q}')$ contains, in a single structure, a long-distance harmonic confinement that smoothly interpolates towards a linear confinement for heavy quarkonia [18,19],

$$\begin{aligned} V(\hat{q}, \hat{q}') &= \frac{3}{4} \omega_{q\bar{q}}^2 \int d^3\vec{r} \left[r^2 (1 + 4\hat{m}_1 \hat{m}_2 A_0 M_s^2 r^2)^{-1/2} - \frac{C_0}{\omega_0^2} \right] e^{i(\hat{q}-\hat{q}') \cdot \vec{r}}, \\ \omega_{q\bar{q}}^2 &= 4M_s \hat{m}_1 \hat{m}_2 \omega_0^2 \alpha_s(M_s^2), \\ M_s &= \max(M, m_1 + m_2), \end{aligned} \quad (2.5)$$

where $\hat{m}_i = m_i/(m_1 + m_2)$, ω_0 is the basic harmonic spring constant, C_0/ω_0^2 is a constant that takes account of the correct zero-point energies, and A_0 controls the harmonic-to-linear interpolation. The running of $\omega_{q\bar{q}}^2$ through $\alpha_s(M_s^2)$ encodes the asymptotic-freedom dependence of the short-range interaction, while the polynomial-in- r structure encodes confinement. In the heavy-quarkonium limit ($\omega \simeq m$) this kernel can be cast, in momentum space, in the compact local form [19],

$$\begin{aligned} V(\hat{q}, \hat{q}') &= \bar{V}(\hat{q}) \delta^3(\hat{q} - \hat{q}'), \\ \bar{V}(\hat{q}) &= \omega_{q\bar{q}}^2 \left[\kappa \bar{\nabla}_{\hat{q}}^2 + \frac{C_0}{\omega_0^2} \right] (2\pi)^3, \end{aligned} \quad (2.6)$$

so that the coupled Salpeter equations for the scalar coefficients of the B-S amplitude reduce to an approximate three-dimensional harmonic-oscillator equation; the harmonic-oscillator form in Eq. (2.10) below is the corresponding $2S$ analytic solution, with the parameter β_V determined in Sec. III from the experimental ratio R_V . The corresponding Salpeter wave function is

$$\psi(\hat{q}) = \frac{i}{2\pi} \int dq_K \Psi(K, q). \quad (2.7)$$

For a vector quarkonium V , the complete decomposition of the Salpeter wave function in terms of various Dirac structures is [14,19,26–29]

$$\begin{aligned} \psi(\hat{q}) &= M \not{\epsilon} f_1(\hat{q}) + \not{\epsilon} \not{K} f_2(\hat{q}) + [\not{\epsilon} \not{q} - \hat{q} \cdot \epsilon] f_3(\hat{q}) \\ &+ [\not{K} \not{q} - (\hat{q} \cdot \epsilon) \not{K}] \frac{f_4(\hat{q})}{M} + (\hat{q} \cdot \epsilon) f_5(\hat{q}) \\ &+ (\hat{q} \cdot \epsilon) \not{K} \frac{f_6(\hat{q})}{M} + (\hat{q} \cdot \epsilon) \not{q} f_7(\hat{q}) \\ &+ (\hat{q} \cdot \epsilon) \not{K} \not{q} \frac{f_8(\hat{q})}{M}, \end{aligned} \quad (2.8)$$

where each scalar function $f_i(\hat{q})$ ($i = 1, 2, \dots, 8$) depends on $\hat{q}^2$, and ϵ is the polarization vector of V . Equation (2.8) is the most general Lorentz-covariant decomposition of the Salpeter amplitude allowed by the $J^{PC} = 1^{--}$ quantum numbers [19,26,27]. Within the CIA framework of Refs. [18–20], the eight scalar coefficients $f_i(\hat{q})$ satisfy a coupled set of Salpeter integral equations whose interaction kernel is the one specified in Eqs. (2.4)–(2.6). In the heavy-quarkonium limit those coupled equations reduce to an approximate three-dimensional harmonic-oscillator equation, and the $f_i(\hat{q})$ become proportional to a single common radial function $f(\hat{q})$ of harmonic-oscillator form, with calculable relative weights ordered by the power-counting rule of Refs. [26,30] in powers of $\hat{q}/M$. In the present work the coupled Salpeter equations are not solved numerically; we instead take over this analytic solution and, to capture the radial node of the $2S$ states, supplement it with the standard first-radial-excitation factor $(1 - 2\hat{q}^2/3\beta_V^2)$ in the oscillator basis. The only phenomenological input left is the harmonic-oscillator parameter β_V , fixed in Sec. III from the experimental ratio R_V . In our recent study of ground-state decays [14] only the Dirac structures up to subleading order, i.e., $\mathcal{O}(\hat{q})$, were retained. In this work, we extend the framework by including all independent structures, i.e., those up to $\mathcal{O}(\hat{q}^2)$. Following Refs. [26,30], all eight scalar coefficients $f_i(\hat{q})$ in Eq. (2.8) are kept and approximated in terms of one common scalar radial function $f(\hat{q})$ with relative weights fixed by the power-counting rule up to $\mathcal{O}(\hat{q}^2/M^2)$ and by matching to the NRQCD spin projector in the limit $m \simeq M/2$ [31]. This procedure provides an approximate closed form of the complete Salpeter amplitude that is consistent with the order at which the $\hat{q}^2$ -order relativistic corrections to the decay amplitude are computed; it is not a truncation that drops Dirac components. The resulting compact expression for $\psi(\hat{q})$ reads,

$$\begin{aligned} \psi(\hat{q}) &= \left(M \not{\epsilon} - \not{\epsilon} \not{K} + 2(\hat{q} \cdot \epsilon) - \frac{2\not{K}(\hat{q} \cdot \epsilon)}{M} + \frac{2\not{K} \not{q}}{M} \right. \\ &\quad \left. + \frac{2(\hat{q} \cdot \epsilon) \not{q}}{M} - \frac{2(\hat{q} \cdot \epsilon) \not{K} \not{q}}{M^2} \right) f(\hat{q}), \end{aligned} \quad (2.9)$$

where the Dirac structures of the wave function are consistent with the spin projector commonly adopted in the NRQCD approach with the limit $m \approx M/2$ [31]. The scalar function is taken as

$$f(\hat{q}) = N_V \left(\frac{3}{2} \right)^{1/2} \frac{1}{\pi^{3/4}} \frac{1}{\beta_V^{3/2}} \left(1 - \frac{2\hat{q}^2}{3\beta_V^2} \right) e^{-\frac{\hat{q}^2}{2\beta_V^2}}, \quad (2.10)$$

where N_V is a constant and β_V is the harmonic oscillator parameter. To retain the radial node structure of the excited state V , we adopt the above harmonic-oscillator form for the scalar function $f(\hat{q})$. We stress that within the CIA framework of Refs. [18–20] this is not an *ad hoc* ansatz competing

with the OGE-type kernel; the spatial part $V(\hat{q}, \hat{q}')$ specified in Eqs. (2.4)–(2.6) is constructed so that, in the heavy-quark limit, the coupled Salpeter equations for the scalar coefficients of Eq. (2.8) reduce to a three-dimensional harmonic-oscillator equation [19]; Eq. (2.10) is the corresponding 2S analytic solution and the nodal factor $(1 - 2\hat{q}^2/3\beta_V^2)$ is its first radial excitation. The instantaneous reduction of the kernel neglects the retardation of the exchanged gluon, whose leading effect is of order v^2 relative to the static potential and generates Breit-Fermi-type terms (spin-orbit, tensor, Darwin) in the bound-state equation. Such retardation and Breit-Fermi corrections enter our calculation of $V \rightarrow ggg$ only through the bound-state wave function, and their net effect is absorbed into the fitted value of β_V . The v^2 relativistic corrections that enter the decay amplitude dynamically—through the small Dirac components of Eq. (2.8) and through the expansion of the hard kernel in $\hat{q}$ —are kept explicitly, and are precisely those that are resummed to all orders in the phenomenological improvement introduced later in Sec. II B following Ref. [15].

This extended form of $\psi(\hat{q})$ systematically includes relativistic spin-orbit couplings, providing a consistent and improved parametrization of the Salpeter wave function. For convenience, we write it as $\psi(\hat{q}) = \mathcal{P}(\hat{q})f(\hat{q})$, with $\mathcal{P}(\hat{q})$ denoting the Dirac-structure factor in parentheses of Eq. (2.9).

B. The decay $V \rightarrow ggg$ in perturbative QCD

Within the B-S formalism, the decay amplitude for $V \rightarrow ggg$ is obtained through the convolution of the bound-state wave function with a perturbative hard-scattering kernel. Since the wave function $\Psi(K, q)$ encodes the full information about the relativistic internal motion, this framework allows a systematic treatment of the higher-order corrections induced by the nodal structure. In the rest frame of V , the amplitude can be written explicitly as

$$\mathcal{A} = \sqrt{N_c} \int \frac{d^4 q}{(2\pi)^4} \text{Tr}[\Psi(K, q) \mathcal{O}(f, \bar{f})], \quad (2.11)$$

where the factor $\sqrt{N_c}$ reflects the color-singlet nature of the quark-antiquark pair, $\mathcal{O}(f, \bar{f})$ denotes the hard-scattering amplitude for $q\bar{q} \rightarrow ggg$, and the momenta of the constituent quark and antiquark are

$$\begin{aligned} f^\mu &= \frac{K^\mu}{2} + q^\mu = \left(\frac{M}{2} + q^0, \mathbf{q} \right), \\ \bar{f}^\mu &= \frac{K^\mu}{2} - q^\mu = \left(\frac{M}{2} - q^0, -\mathbf{q} \right). \end{aligned} \quad (2.12)$$

Under the CIA [18,20,23,25,26], it is consistent to treat $q^0 \ll M$. Under this approximation, the momenta reduce to

$$\begin{aligned} f^\mu &\approx \left(\frac{M}{2}, \mathbf{q} \right) = \frac{K^\mu}{2} + \hat{q}^\mu, \\ \bar{f}^\mu &\approx \left(\frac{M}{2}, -\mathbf{q} \right) = \frac{K^\mu}{2} - \hat{q}^\mu, \end{aligned} \quad (2.13)$$

and the hard-scattering amplitude simplifies to

$$\mathcal{O}(f, \bar{f}) \approx \mathcal{O}(\hat{q}). \quad (2.14)$$

From another point of view [15], this treatment can be connected with the on shell condition, which maintains the gauge invariance of the hard-scattering amplitude.

With the above approximations, the decay amplitude of $V \rightarrow ggg$ takes the compact form

$$\mathcal{A} = -i\sqrt{N_c} \int \frac{d^3 \hat{q}}{(2\pi)^3} f(\hat{q}) \text{Tr}[\mathcal{P}(\hat{q}) \mathcal{O}(\hat{q})]. \quad (2.15)$$

The explicit expression for the hard-scattering amplitude $\mathcal{O}(\hat{q})$ is

$$\begin{aligned} \mathcal{O}(\hat{q}) &= \frac{-ig_s^3}{4N_c} (\text{Tr}[T^{a_1} T^{a_2} T^{a_3}] I(k_1, k_2, k_3, \epsilon_1, \epsilon_2, \epsilon_3) + \text{Tr}[T^{a_2} T^{a_1} T^{a_3}] I(k_2, k_1, k_3, \epsilon_2, \epsilon_1, \epsilon_3) \\ &\quad + \text{Tr}[T^{a_2} T^{a_3} T^{a_1}] I(k_2, k_3, k_1, \epsilon_2, \epsilon_3, \epsilon_1) + \text{Tr}[T^{a_3} T^{a_2} T^{a_1}] I(k_3, k_2, k_1, \epsilon_3, \epsilon_2, \epsilon_1) \\ &\quad + \text{Tr}[T^{a_3} T^{a_1} T^{a_2}] I(k_3, k_1, k_2, \epsilon_3, \epsilon_1, \epsilon_2) + \text{Tr}[T^{a_1} T^{a_3} T^{a_2}] I(k_1, k_3, k_2, \epsilon_1, \epsilon_3, \epsilon_2)) \end{aligned} \quad (2.16)$$

with

$$\begin{aligned} I(k_1, k_2, k_3, \epsilon_1, \epsilon_2, \epsilon_3) &= \frac{\not{\epsilon}_1^* \cdot (\not{k}_1 - \not{k}_2 - \not{k}_3 + 2\hat{q} + M) \cdot \not{\epsilon}_2^* \cdot (\not{k}_1 + \not{k}_2 - \not{k}_3 + 2\hat{q} + M) \cdot \not{\epsilon}_3^*}{[-k_1 \cdot k_2 - k_1 \cdot k_3 + 2(k_1 \cdot \hat{q}) + \hat{q}^2 + i\epsilon][-k_1 \cdot k_3 - k_2 \cdot k_3 - 2(k_3 \cdot \hat{q}) + \hat{q}^2 + i\epsilon]}, \end{aligned} \quad (2.17)$$

where k_i and ϵ_i ($i = 1, 2, 3$) are the momenta and polarization vectors of the outgoing gluons. Using the color identity $\text{Tr}[T^{a_1} T^{a_2} T^{a_3}] = (d^{a_1 a_2 a_3} + i f^{a_1 a_2 a_3})/4$, the amplitude decomposes into a part proportional to the symmetric structure constant $d^{a_1 a_2 a_3}$ and another proportional to the antisymmetric constant $f^{a_1 a_2 a_3}$. The latter contribution vanishes identically after integration over the internal momentum $\hat{q}$. Thus the hard-scattering amplitude reduces to

$$\mathcal{O}(\hat{q}) = \frac{-i g_s^3 d^{a_1 a_2 a_3}}{16 N_c} (I(k_1, k_2, k_3, \epsilon_1, \epsilon_2, \epsilon_3) + 5 \text{ permutations of } 1, 2, \text{ and } 3.). \quad (2.18)$$

The decay width is obtained from

$$\Gamma^{ggg} = \frac{1}{2M} \int \frac{1}{3} \frac{1}{3!} \sum_{\text{color, pol}} |\mathcal{A}|^2 d\Phi_3, \quad (2.19)$$

where $\int d\Phi_3$ denotes the integration over the three-body phase space, the factor $1/3$ averages over the initial quarkonium polarizations, $1/3!$ accounts for the identical gluons in the final state, and the sum runs over the colors and polarizations of all external particles.

The decay $V \rightarrow ggg$ possesses a soft-gluon singularity, which gives rise to a potential IR divergence. When the full $\hat{q}$ -dependence of the hard amplitude is retained, we find the integration over the internal momentum $\hat{q}$ becomes divergent starting at $\mathcal{O}(\hat{q}^4)$. Physically, this divergence corresponds to the process in which a soft gluon converts the incoming S -wave color-singlet quarkonium into a P -wave color-octet intermediate state [31]. Within the established NRQCD factorization formalism, such IR divergences can be systematically absorbed into the nonperturbative matrix elements of P -wave color-octet operators [31]. Phenomenological determinations of these higher-order matrix elements remain uncertain, but power-counting arguments suggest that $\mathcal{O}(v^4)$ contributions are generally suppressed compared with the leading $\mathcal{O}(v^2)$ terms [14,31].

Up to $\hat{q}^2$ -order relativistic corrections, the modulus squared of the decay amplitude is obtained as

$$|\mathcal{A}|^2 = \left| \sqrt{N_c} \int \frac{d^3 \hat{q}}{(2\pi)^3} f(\hat{q}) \mathcal{M}^0(0) \right|^2 + 2\text{Re} \left[N_c \int \frac{d^3 \hat{q}}{(2\pi)^3} \frac{d^3 \hat{q}'}{(2\pi)^3} f(\hat{q}) f^*(\hat{q}') \mathcal{M}^{0*}(0) \mathcal{M}^1(0) \right] \quad (2.20)$$

with

$$\mathcal{M}^0(0) = \text{Tr}[\mathcal{P}(0) \mathcal{O}(0)] \quad (2.21)$$

and

$$\mathcal{M}^1(0) = \frac{\hat{q}^\alpha \hat{q}^\beta}{2!} \left[\frac{\partial \text{Tr}[\mathcal{P}(\hat{q}) \mathcal{O}(\hat{q})]}{\partial \hat{q}^\alpha \partial \hat{q}^\beta} \right]_{\hat{q}=0}. \quad (2.22)$$

Here we expand the modulus squared $|\mathcal{A}|^2$ to quadratic order in the internal momentum $\hat{q}$, and make the following substitution to project out the S -wave state:

$$\hat{q}^\alpha \hat{q}^\beta \rightarrow \frac{\hat{\mathbf{q}}^2}{3} \left(-g^{\alpha\beta} + \frac{K^\alpha K^\beta}{M^2} \right), \quad (2.23)$$

so that $\mathcal{M}^1(0)$ becomes

$$\mathcal{M}^1(0) = \frac{1}{2!} \frac{\hat{\mathbf{q}}^2}{3} \left[\left(-g^{\alpha\beta} + \frac{K^\alpha K^\beta}{M^2} \right) \frac{\partial \text{Tr}[\mathcal{P}(\hat{q}) \mathcal{O}(\hat{q})]}{\partial \hat{q}^\alpha \partial \hat{q}^\beta} \right]_{\hat{q}=0}. \quad (2.24)$$

Performing the integration over the three-body phase space in Eq. (2.19) yields the decay width including $\hat{q}^2$ -order relativistic corrections,

$$\Gamma^{ggg} = \frac{10(\pi^2 - 9)\alpha_s^3}{81\pi^6 M} \int d^3 \hat{q} d^3 \hat{q}' f(\hat{q}) f(\hat{q}') \times \left(1 - \frac{48 + 121\pi^2}{48(\pi^2 - 9)} \frac{\hat{\mathbf{q}}^2}{M^2} \right). \quad (2.25)$$

However, our subsequent numerical evaluation for the radially excited state $\psi(2S)$ (presented later in this work) reveals that this expression produces an unphysical negative decay width. Inspired by the phenomenological improvement proposed in Ref. [15], we therefore introduce the following improved expression:

$$\Gamma^{ggg} = \frac{10(\pi^2 - 9)\alpha_s^3}{81\pi^6 M} \times \left| \int d^3 \hat{q} f(\hat{q}) \left[1 + \frac{48 + 121\pi^2}{96(\pi^2 - 9)} \frac{\hat{\mathbf{q}}^2}{M^2} \right]^{-1} \right|^2. \quad (2.26)$$

This form preserves the $\hat{q}^2$ -order relativistic correction in the small- $\hat{q}$ limit while effectively incorporating partial higher-order contributions at large momentum. It guarantees a positive definite decay width for $\psi(2S)$ and ensures better consistency with the full $\hat{q}$ -dependent hard kernel across both nonrelativistic and highly relativistic regimes. This improved treatment significantly enhances the convergence of the integral of the hard kernel convolved with the scalar wave function $f(\hat{q})$ across both low- and high-momentum regions; it exactly reproduces the $\hat{q}^2$ -order result at small $\hat{q}$ (owing to the rapid convergence of the Taylor expansion), while the denominator provides natural suppression at large $\hat{q}$ and prevents any pathological behavior from the wave-function tail. It is worth noting that a strict perturbative expansion to $\mathcal{O}(\hat{q}^4)$ would induce an IR divergence. Although within the NRQCD factorization formalism such

divergences can be absorbed into nonperturbative P -wave color-octet matrix elements, the phenomenological determination of these additional parameters introduces significant uncertainty for the decays of radially excited states. In this context, while our phenomenological treatment is not a first-principles derivation, its remarkably concise and physically transparent form offers a practical yet intuitive way to incorporate the dominant higher-order relativistic corrections beyond $\hat{q}^2$, together with the nodal structure of the radial wave function that is characteristic of the $2S$ state.

We can now exploit these expressions to study the polarization characteristics and symmetries of the decay.

To discuss the symmetry in the decay process $V \rightarrow ggg$, the corresponding polarized decay widths are given by

$$\Gamma_{[\lambda_V, \lambda_1, \lambda_2, \lambda_3]}^{ggg} = \frac{1}{2M} \int \frac{1}{3!} \sum_{\text{color}} |\mathcal{A}|^2 d\Phi_3. \quad (2.27)$$

Here, λ_V denotes the helicity of the heavy quarkonium, while λ_i ($i = 1, 2, 3$) represents the gluon helicity. We adopt the following conventions:

$$\begin{aligned} K &= M(1, 0, 0, 0), & \varepsilon(\pm 1) &= \frac{1}{\sqrt{2}}(0, \mp 1, -i, 0), & \varepsilon(0) &= (0, 0, 0, 1) \\ k_1 &= \frac{Mx_1}{2}(1, 0, 0, 1), & \epsilon_1(\pm 1) &= \frac{1}{\sqrt{2}}(0, \mp 1, -i, 0), \\ k_2 &= \frac{Mx_2}{2}(1, \sin \theta_{12}, 0, \cos \theta_{12}), & \epsilon_2(\pm 1) &= \frac{1}{\sqrt{2}}(0, \mp \cos \theta_{12}, -i, \pm \sin \theta_{12}), \\ k_3 &= \frac{Mx_3}{2}(1, -\sin \theta_{13}, 0, \cos \theta_{13}), & \epsilon_3(\pm 1) &= \frac{1}{\sqrt{2}}(0, \mp \cos \theta_{13}, -i, \mp \sin \theta_{13}), \end{aligned} \quad (2.28)$$

where θ_{ij} is the angle between gluons i and j .

III. RESULTS AND DISCUSSIONS

A. Polarized decay widths

The set-up of polarized decay widths $\Gamma_{[\lambda_V, \lambda_1, \lambda_2, \lambda_3]}^{ggg}$, the kinematic conventions for the helicity vectors and the gluon momenta, the helicity-flip and gluon-permutation symmetries, the resulting equivalence classes Γ_1 – Γ_4 of non-vanishing configurations, and the analytic $\hat{q}^2$ -order expressions for these polarized widths are obtained exactly as in our previous work on the ground-state decays $J/\psi \rightarrow ggg$ and $\Upsilon(1S) \rightarrow ggg$ [14], now with the Dirac structures extended to $\mathcal{O}(\hat{q}^2)$ and the $1S$ scalar radial function replaced

by the $2S$ one in Eq. (2.10). To avoid a lengthy repetition, we summarize here only the symmetry-derived results that are directly needed for the present discussion, and collect the corresponding explicit expressions in the Appendix.

In particular, the nonvanishing polarized decay widths fall into four equivalence classes $\Gamma_1, \Gamma_2, \Gamma_3$, and Γ_4 defined in Eqs. (A4)–(A7) of the Appendix; $\Gamma_{1,2,3}$ each collect four equivalent helicity configurations, while Γ_4 collects only two. The corresponding analytic $\hat{q}^2$ -order expressions are given in Eqs. (A8)–(A11) of the Appendix. Employing the phenomenological treatment of higher-order relativistic corrections introduced in Sec. II B leads to the improved expressions,

$$\Gamma_1 = \frac{10\sqrt{3}M^3N_V^2(3\pi^2 - 16)^3\alpha_s^3}{81\beta_V\pi^{9/2}(1264 + 111\pi^2)^2} \left(1 - \frac{3\sqrt{6\pi a_1}(1 + 8a_1)}{(1 + 24a_1)} e^{6a_1} \text{erfc}(\sqrt{6a_1}) \right)^2, \quad (3.1)$$

$$\Gamma_2 = \frac{10\sqrt{3}M^3N_V^2(29\pi^2 - 280)^3\alpha_s^3}{81\beta_V\pi^{9/2}(265\pi^2 - 1424)^2} \left(1 - \frac{3\sqrt{6\pi a_2}(1 + 8a_2)}{(1 + 24a_2)} e^{6a_2} \text{erfc}(\sqrt{6a_2}) \right)^2, \quad (3.2)$$

$$\Gamma_3 = \frac{320\sqrt{3}M^3N_V^2\alpha_s^3}{81\beta_V^7\pi^{9/2}(88 + 27\pi^2)^5} \left(1 - \frac{6\sqrt{3\pi}(16 + a_3)}{\sqrt{a_3}(48 + a_3)} e^{12/a_3} \text{erfc}\left(\frac{2\sqrt{3}}{\sqrt{a_3}}\right) \right)^2, \quad (3.3)$$

$$\Gamma_4 = \frac{20480\sqrt{3}M^3N_V^2(\pi^2 - 9)^3\alpha_s^3}{81\beta_V\pi^{9/2}(48 + 121\pi^2)^2} \left(1 - \frac{12\sqrt{3\pi a_4}(1 + 64a_4)}{(1 + 192a_4)} e^{48a_4} \text{erfc}(4\sqrt{3a_4}) \right)^2, \quad (3.4)$$

where $\text{erfc}(x)$ is the complementary error function, and the auxiliary parameters are defined as $a_1 = \frac{(3\pi^2-16)M^2}{(1264+111\pi^2)\beta_V^2}$, $a_2 = \frac{(29\pi^2-280)M^2}{(265\pi^2-1424)\beta_V^2}$, $a_3 = \frac{(88+27\pi^2)\beta_V^2}{M^2}$, and $a_4 = \frac{(\pi^2-9)M^2}{(48+121\pi^2)\beta_V^2}$.

The explicit results for Γ_1 to Γ_4 are found to automatically satisfy the model-independent symmetry relations, thereby providing a direct and nontrivial verification of our calculational framework. Moreover, our numerical analysis will show that the $\hat{q}^2$ -order relativistic corrections yield unphysical results for $\psi(2S)$, while the phenomenologically improved expressions give physically consistent predictions. A detailed discussion follows in the next section.

Beyond this internal consistency, these polarized decay widths offer a richer set of observables compared to the unpolarized decay width alone. They encode specific correlations among the gluon helicities; these correlations are transferred through hadronization and can ultimately be probed indirectly via final-state hadrons. A prominent example is the decay $V \rightarrow B\bar{B}$ (where B denotes a baryon). Specifically, a key finding from Eqs. (A8)–(A11) [or (3.1)–(3.4)] is the strong dominance of decays from transversely polarized V ($\lambda_V = \pm 1$) over those from longitudinally polarized V ($\lambda_V = 0$). This pronounced transverse dominance implies the enhancement of transverse spin polarizations and correlations in the produced baryon pairs. Such effects can be probed experimentally via multidimensional angular-distribution fits and moment analyses of processes such as $V \rightarrow p\bar{p}$ or $\Lambda\bar{\Lambda}$ [32,33], thereby connecting our predictions for the polarized decay widths to measurable hadronic observables.

B. Unpolarized decay widths

The unpolarized decay width $\Gamma(V \rightarrow ggg)$ including $\hat{q}^2$ -order relativistic corrections reads

$$\Gamma(V \rightarrow ggg) = \frac{40(\pi^2 - 9)N_V^2\beta_V^3\alpha_s^3}{27\pi^{9/2}M} \left(1 - \kappa \frac{\beta_V^2}{M^2}\right) \quad (3.5)$$

with $\kappa \equiv \frac{7(48+121\pi^2)}{48(\pi^2-9)}$. This expression shows that the relativistic correction, quantified by $\kappa\beta_V^2/M^2$, is significantly larger for $\psi(2S)$ (where $\kappa\beta_{\psi(2S)}^2/M^2 \approx 2.70$) than for $\Upsilon(2S)$ [where $\kappa\beta_{\Upsilon(2S)}^2/M^2 \approx 0.88$], owing to the lighter charm-quark mass. More importantly, for $\psi(2S)$, the decay width calculated to $\hat{q}^2$ order yields an unphysical negative value. This directly indicates the poor convergence of the relativistic expansion and underscores the necessity of including higher-order effects.

To address this issue and obtain a physically consistent prediction, we incorporate a selected subset of higher-order relativistic corrections through our phenomenological treatment. Summing the nonvanishing polarized widths over the helicity configurations of the four equivalence classes listed

in the Appendix—four configurations for each of $\Gamma_{1,2,3}$ and two for Γ_4 —and averaging over the three polarizations of the initial vector meson, we reconstruct the unpolarized decay width from the polarized components as

$$\Gamma(V \rightarrow ggg) = \frac{1}{3}(4\Gamma_1 + 4\Gamma_2 + 4\Gamma_3 + 2\Gamma_4), \quad (3.6)$$

where Γ_1 – Γ_4 are given by Eqs. (3.1)–(3.4) and are calculated by incorporating $\hat{q}^2$ -order relativistic corrections and systematically including a selected subset of higher-order effects, thereby restoring physical positivity. Their combination according to Eq. (3.6) provides a more reliable theoretical description of the decay width.

Since relativistic corrections arising from the internal motion of the bound quarks and QCD radiative corrections originating from additional gluon emissions during the hard process are governed by distinct dynamical scales, it is standard to treat them as factorizable at the present level of accuracy. Incorporating both types of corrections, we obtain the unpolarized decay width as the product of the relativistic correction factor and the QCD radiative correction factor,

$$\Gamma(V \rightarrow ggg) = \frac{40(\pi^2 - 9)N_V^2\beta_V^3\alpha_s^3}{27\pi^{9/2}M} \left(1 - \kappa \frac{\beta_V^2}{M^2}\right) \left(1 - C_V \frac{\alpha_s}{\pi}\right). \quad (3.7)$$

Here, the relativistic correction factor $(1 - \kappa \frac{\beta_V^2}{M^2})$ includes only $\hat{q}^2$ -order corrections, while the QCD radiative correction factor $(1 - C_V \frac{\alpha_s}{\pi})$ contains the $\mathcal{O}(\alpha_s)$ contribution. The process-dependent coefficient C_V takes the values $C_{\psi(2S)} = 3.7$ and $C_{\Upsilon(2S)} = 4.9$, as determined in Ref. [34].

However, as noted earlier, the $\hat{q}^2$ -order treatment leads to unphysical results for $\psi(2S)$. To go beyond it, we include a phenomenological subset of higher-order relativistic effects. The improved expression is obtained by replacing the product of the leading-order width and the relativistic correction factor in Eq. (3.7) with the combination of phenomenologically improved polarized widths,

$$\Gamma(V \rightarrow ggg) = \frac{1}{3}(4\Gamma_1 + 4\Gamma_2 + 4\Gamma_3 + 2\Gamma_4) \left(1 - C_V \frac{\alpha_s}{\pi}\right), \quad (3.8)$$

where Γ_1 – Γ_4 are given by Eqs. (3.1)–(3.4), and C_V is understood as $C_{\psi(2S)}$ or $C_{\Upsilon(2S)}$ for the respective quarkonium.

We also compute the leptonic decay width $\Gamma(V \rightarrow e^+e^-)$. Including $\hat{q}^2$ -order relativistic and one-loop QCD corrections, it is given by

$$\Gamma(V \rightarrow e^+e^-) = \frac{12e_Q^2\alpha^2N_V^2\beta_V^3}{\pi^{7/2}M} \left(1 - \frac{28\beta_V^2}{3M^2}\right) \left(1 - \frac{16\alpha_s}{3\pi}\right), \quad (3.9)$$

where e_Q denotes the electric charge of the heavy quark. Applying an analogous phenomenological treatment to that employed for the three-gluon decay yields the improved form,

$$\begin{aligned} \Gamma(V \rightarrow e^+e^-) &= \frac{12e_Q^2\alpha^2 N_V^2 \beta_V^3}{\pi^{7/2}M} \left(\frac{27\pi \left(a_0 \left(1 + \frac{\beta_V^2}{M^2} \right) - \frac{2\sqrt{3}\beta_V}{9\sqrt{\pi}M} \left(3 + \frac{\beta_V^2}{M^2} \right)^2 \right)}{16 \left(\frac{\beta_V}{M} \right)^{10}} \right) \\ &\times \left(1 - \frac{16\alpha_s}{3\pi} \right), \end{aligned} \quad (3.10)$$

with $a_0 = e^{3M^2/(4\beta_V^2)} \text{erfc}(\frac{\sqrt{3}M}{2\beta_V})$.

To substantially reduce the theoretical uncertainties, we construct the ratio of the hadronic to leptonic decay widths,

$$R_V = \frac{\Gamma(V \rightarrow ggg)}{\Gamma(V \rightarrow e^+e^-)}. \quad (3.11)$$

In this ratio, the common nonperturbative factor $N_V^2 \beta_V^3$, which is associated with the quarkonium wave function at the origin, cancels exactly. Consequently, R_V provides a significantly cleaner observable, offering a more direct probe of the short-distance dynamics and thus yielding a more robust theoretical prediction.

For our numerical analysis, we take the meson masses and total widths from the PDG [35]. The QCD running coupling constants are taken as $\alpha_s(M_{\psi(2S)}/2) = 0.31$ and $\alpha_s(M_{\Upsilon(2S)}/2) = 0.21$, which are calculated through the two-loop renormalization group equation. We adopt the harmonic oscillator parameters $\beta_{\psi(2S)} = 420$ MeV for $\psi(2S)$ and $\beta_{\Upsilon(2S)} = 650$ MeV for $\Upsilon(2S)$. The value $\beta_{\psi(2S)} = 420$ MeV is chosen to be consistent with the nodal position of the $\psi(2S)$ wave function extracted from lattice QCD [8], while $\beta_{\Upsilon(2S)} = 650$ MeV lies within the range obtained in studies of highly excited bottomonium states [36]. The constants $N_{\psi(2S)} = 0.232 \text{ MeV}^{-1/2}$ for

$\psi(2S)$ and $N_{\Upsilon(2S)} = 0.167 \text{ MeV}^{-1/2}$ for $\Upsilon(2S)$ are determined by fitting to the experimental branching ratios of the corresponding leptonic decays $V \rightarrow e^+e^-$. With these inputs, we present our predictions for the branching ratios $\mathcal{B}(V \rightarrow ggg)$, $\mathcal{B}(V \rightarrow e^+e^-)$, and their ratio R_V in Table I. Theoretical uncertainties are omitted; they arise mainly from the scale choice for α_s and uncalculated higher-order effects, and are expected to be below 30%. The second column (“ $\hat{q}^2$ -order”) includes $\hat{q}^2$ -order relativistic corrections, while the third (“Higher-order”) incorporates phenomenologically improved higher-order corrections; both include the known $\mathcal{O}(\alpha_s)$ QCD radiative corrections.

The numerical results in Table I reveal several key features. First, as mentioned earlier, the predictions at $\hat{q}^2$ -order relativistic correction (second column) yield unphysical negative values for both the branching ratio $\mathcal{B}(\psi(2S) \rightarrow ggg)$ and the ratio $R_{\psi(2S)}$. This dramatic failure arises because the three-gluon decay amplitude involves a convolution with multiple quark propagators, making it highly sensitive to the momentum distribution across the entire wave function. The characteristic nodal structure of the $2S$ radial wave function introduces significant destructive interference in this convolution, which is severely misrepresented when only the leading $\hat{q}^2$ -order correction is included. In contrast, after applying our phenomenologically improved treatment, the results shift dramatically and come into good agreement with experimental data. This strongly indicates that higher-order relativistic corrections beyond $\hat{q}^2$ play a crucial role in the gluonic decay of the $2S$ state. A similar conclusion was previously reached in the exclusive hadronic decay $\psi(2S) \rightarrow p\bar{p}$ [13].

Second, in the leptonic decay channel $V \rightarrow e^+e^-$, the predictions at $\hat{q}^2$ -order (second column) are already very close to those with higher-order corrections (third column) and agree well with the experimental value. This suggests that the dominant relativistic corrections in the leptonic decay are largely captured already at the $\hat{q}^2$ level. The relatively mild impact of higher-order terms highlights the fast convergence of the relativistic expansion for this

TABLE I. Theoretical predictions at $\hat{q}^2$ -order relativistic correction and with higher-order relativistic corrections, together with experimental data, for the branching ratios $\mathcal{B}(V \rightarrow ggg)$, $\mathcal{B}(V \rightarrow e^+e^-)$, and the ratio R_V in $\psi(2S)$ and $\Upsilon(2S)$ decays.

	$\hat{q}^2$ -order	Higher-order	Experiment [35,37,38]
$\mathcal{B}(\psi(2S) \rightarrow ggg)$	-88.8%	9.4%	$(10.6 \pm 1.6)\%$
$\mathcal{B}(\psi(2S) \rightarrow e^+e^-)$	7.94×10^{-3}	8.04×10^{-3}	$(7.94 \pm 0.22) \times 10^{-3}$
$R_{\psi(2S)}$	-111.8	11.8	13.4 ± 2.4
$\mathcal{B}(\Upsilon(2S) \rightarrow ggg)$	13.8%	54.5%	$(58.8 \pm 1.2)\%$
$\mathcal{B}(\Upsilon(2S) \rightarrow e^+e^-)$	1.91%	1.91%	$(1.91 \pm 0.16)\%$
$R_{\Upsilon(2S)}$	7.2	28.5	30.8 ± 3.2

process and demonstrates its practical reliability for vector-meson leptonic widths.

Third, a striking contrast emerges when comparing the convergence behavior between the gluonic and leptonic decay channels. While the relativistic expansion converges rapidly for $\Gamma(V \rightarrow e^+e^-)$, it converges extremely slowly for $\Gamma(V \rightarrow ggg)$. This profound difference originates from the distinct dynamics of the two processes. The leptonic decay is mediated by a single virtual photon, and its width is predominantly sensitive to the wave function at the origin, where the nodal structure has a milder effect. In contrast, the amplitude for $V \rightarrow ggg$ involves a convolution with multiple quark propagators, making it acutely sensitive to the detailed momentum distribution across the entire wave function. The nodal structure introduces destructive interference in this convolution, which is severely misrepresented at $\hat{q}^2$ order. Thus, it is the combination of the nodal structure and the specific multigluon dynamics that leads to the dramatic failure of the low-order expansion in the gluonic channel.

Once both the dominant higher-order relativistic effects and the next-to-leading-order QCD radiative corrections are included, our predictions for $\mathcal{B}(V \rightarrow ggg)$, $\mathcal{B}(V \rightarrow e^+e^-)$, and R_V are in good agreement with the experimental data for both $\psi(2S)$ and $\Upsilon(2S)$.


C. Harmonic oscillator parameter

As a cross-check, we further extract the harmonic oscillator parameter β_V within our B-S framework. This parameter determines the characteristic momentum-space width of the quarkonium wave function. Using the ratio

$$R_V = \frac{\Gamma(V \rightarrow ggg)}{\Gamma(V \rightarrow e^+e^-)}, \quad (3.12)$$

together with the experimental values [35,37,38],

$$R_{\psi(2S)}^{\text{exp}} = 13.4 \pm 2.4, \quad R_{\Upsilon(2S)}^{\text{exp}} = 30.8 \pm 3.2, \quad (3.13)$$



we obtain the following allowed ranges:

$$\begin{aligned} \beta_{\psi(2S)} &\in (360, 430) \text{ MeV}, \\ \beta_{\Upsilon(2S)} &\in (540, 670) \text{ MeV}. \end{aligned} \quad (3.14)$$

The functional dependence of the ratio R_V on the harmonic oscillator parameter β_V is displayed in Fig. 1. Our extracted β_V values lie at the lower end of the ranges commonly reported in the literature for these states, with typical values being approximately (400–700) MeV for $\psi(2S)$ and (600–1300) MeV for $\Upsilon(2S)$ [39–46]. These broader ranges encompass values derived from various approaches, such as light-front quark models [39–42] and potential models [43–46]. Our analysis, which employs the distinct observable R_V , provides an independent determination that refines these phenomenological estimates, consistently favoring the lower portion of the allowed intervals.

This preference for smaller β_V values carries important physical implications. Because the B-S framework provides a fully relativistic treatment of bound-state dynamics, it yields a wave function that is more localized in momentum space than those from phenomenological models. This increased localization, which corresponds directly to a smaller β_V [45], consequently ensures a more reliable description of the high-momentum components that are often poorly captured by effective models. Hence, the larger β_V values of many phenomenological models can be interpreted as an effective increase in the width parameter that compensates for their missing relativistic and high-momentum dynamics. Therefore, by calibrating this key parameter through processes that are acutely sensitive to such dynamics, our work establishes a concrete benchmark for nonperturbative models and enables a deeper understanding of heavy quarkonium structure.

IV. SUMMARY

In this work, we have investigated the three-gluon decays of the radially excited vector quarkonia $\psi(2S)$ and $\Upsilon(2S)$ within the B-S formalism. By employing analytic harmonic




FIG. 1. Dependence of the ratio R_V on the harmonic oscillator parameter β_V . The green curve shows our theoretical prediction. The yellow band represents the experimental value with its 1σ uncertainty. The intersection determines the extracted β_V range.

oscillator wave functions that explicitly incorporate the nodal structure of the $2S$ states, we obtained fully analytic expressions for the decay amplitudes. To go beyond the $\hat{q}^2$ -order relativistic corrections, we introduced a simple yet effective phenomenological treatment that incorporates the dominant higher-order contributions while preserving the correct low-momentum limit. Model-independent relations among polarized decay widths, derived from helicity-flip and phase-space symmetries, allow us to classify the nonvanishing contributions into four distinct groups, each with compact analytic formulas.

Our numerical results reveal a striking contrast between decay channels. At $\hat{q}^2$ order the gluonic width for $\psi(2S)$ and the ratio $R_{\psi(2S)}$ become unphysical and negative, exposing the breakdown of the low-order expansion caused by destructive interference from the $2S$ nodal structure in the multigluon convolution. The improved treatment restores positive, physically consistent predictions that agree well with experimental branching ratios for both $\psi(2S)$ and $\Upsilon(2S)$. In contrast, the leptonic decay $V \rightarrow e^+e^-$ already converges rapidly at $\hat{q}^2$ order, underscoring that the three-gluon channel is far more sensitive to the detailed momentum distribution across the entire wave function. As a cross-check, the harmonic oscillator parameter β_V extracted from the ratio R_V further confirms the reliability of our framework.

It is worth noting that, while our phenomenological treatment is not a first-principles derivation, its remarkably concise and physically transparent form offers a practical yet intuitive way to incorporate the dominant higher-order relativistic corrections beyond $\hat{q}^2$, together with the nodal structure characteristic of the $2S$ radial wave function. These findings establish gluonic decays of radially excited heavy quarkonia as a precision probe of relativistic bound-state dynamics, demonstrating that the internal nodal structure dramatically amplifies the sensitivity to higher-order effects in multigluon channels. This work thereby provides valuable insight into the unique role of excited-state internal structure in heavy quarkonium physics.

ACKNOWLEDGMENTS

The work of C.-J. F. is supported by the Hubei Provincial Natural Science Foundation Youth Project under Grant No. 2024AFB151. The work of J.-K. H. is supported by the National Natural Science Foundation of China under Grant No. 12305086, the Open Fund of the Key Laboratory of Quark and Lepton Physics (MOE) under Grant No. QLPL2024P01, and the Project of Science and Technology Research Program of the Hubei Provincial Department of Education under Grant No. Q20222504.

DATA AVAILABILITY

The data that support the findings of this article are not publicly available. The data are available from the authors upon reasonable request.

APPENDIX: SYMMETRY ANALYSIS AND ANALYTIC $\hat{q}^2$ -ORDER EXPRESSIONS OF THE POLARIZED DECAY WIDTHS

For completeness we collect here the symmetry analysis of the polarized decay widths $\Gamma_{[\lambda_V, \lambda_1, \lambda_2, \lambda_3]}^{ggg}$ and the analytic $\hat{q}^2$ -order expressions for the four equivalence classes Γ_1 – Γ_4 . The derivations follow exactly the methodology established in our previous work on the ground-state decays $J/\psi \rightarrow ggg$ and $\Upsilon(1S) \rightarrow ggg$ [14], now with the Dirac structures extended to $\mathcal{O}(\hat{q}^2)$ and the $1S$ scalar radial function replaced by the $2S$ one in Eq. (2.10).

Owing to helicity-flip symmetry and phase-space symmetry, the following relations are satisfied:

$$\begin{aligned}\Gamma_{[\lambda_V, \lambda_1, \lambda_2, \lambda_3]}^{ggg} &= \Gamma_{[-\lambda_V, -\lambda_1, -\lambda_2, -\lambda_3]}^{ggg}, \\ \Gamma_{[\lambda_V, \lambda_1, \lambda_2, \lambda_3]}^{ggg} &= \Gamma_{[\lambda_V, \lambda_1, \lambda_3, \lambda_2]}^{ggg}.\end{aligned}\quad (A1)$$

Furthermore, after summing over the polarizations λ_V of the initial quarkonium to eliminate the preferred direction defined by the initial polarization, the full permutation symmetry among the three identical gluons is restored, leading to the additional equalities,

$$\begin{aligned}\sum_{\lambda_V=-1,0,1} \Gamma_{[\lambda_V, 1, 1, -1]}^{ggg} &= \sum_{\lambda_V=-1,0,1} \Gamma_{[\lambda_V, 1, -1, 1]}^{ggg} = \sum_{\lambda_V=-1,0,1} \Gamma_{[\lambda_V, -1, 1, 1]}^{ggg} \\ &= \sum_{\lambda_V=-1,0,1} \Gamma_{[\lambda_V, -1, 1, -1]}^{ggg} = \sum_{\lambda_V=-1,0,1} \Gamma_{[\lambda_V, 1, -1, -1]}^{ggg} \\ &= \sum_{\lambda_V=-1,0,1} \Gamma_{[\lambda_V, -1, -1, 1]}^{ggg}, \\ \sum_{\lambda_V=-1,0,1} \Gamma_{[\lambda_V, 1, 1, 1]}^{ggg} &= \sum_{\lambda_V=-1,0,1} \Gamma_{[\lambda_V, -1, -1, -1]}^{ggg}.\end{aligned}\quad (A2)$$

These relations originate directly from the fundamental symmetries of QCD and the bosonic nature of the gluons and provide a model-independent self-consistency check for the present calculation.

The following helicity configurations yield identically zero width:

$$\begin{aligned}\Gamma_{[1, 1, 1, 1]}^{ggg} &= \Gamma_{[-1, -1, -1, -1]}^{ggg} = 0, \\ \Gamma_{[1, -1, -1, -1]}^{ggg} &= \Gamma_{[-1, 1, 1, 1]}^{ggg} = 0, \\ \Gamma_{[0, 1, 1, 1]}^{ggg} &= \Gamma_{[0, -1, -1, -1]}^{ggg} = 0, \\ \Gamma_{[1, -1, 1, 1]}^{ggg} &= \Gamma_{[-1, 1, -1, -1]}^{ggg} = 0, \\ \Gamma_{[0, 1, -1, -1]}^{ggg} &= \Gamma_{[0, -1, 1, 1]}^{ggg} = 0.\end{aligned}\quad (A3)$$

The remaining nonvanishing polarized decay widths fall into four equivalence classes labeled Γ_1 , Γ_2 , Γ_3 , and Γ_4 ,

$$\Gamma_{[1, 1, 1, -1]}^{ggg} = \Gamma_{[-1, -1, -1, 1]}^{ggg} = \Gamma_{[-1, 1, 1, -1]}^{ggg} = \Gamma_{[1, -1, -1, 1]}^{ggg} = \Gamma_1, \quad (A4)$$

$$\Gamma_{[1,-1,1,-1]}^{ggg} = \Gamma_{[-1,1,-1,1]}^{ggg} = \Gamma_{[-1,1,1,-1]}^{ggg} = \Gamma_{[1,-1,-1,1]}^{ggg} = \Gamma_2, \quad (\text{A5})$$

$$\Gamma_{[0,1,1,-1]}^{ggg} = \Gamma_{[0,-1,-1,1]}^{ggg} = \Gamma_{[0,-1,1,-1]}^{ggg} = \Gamma_{[0,1,-1,1]}^{ggg} = \Gamma_3, \quad (\text{A6})$$

$$\Gamma_{[1,1,-1,-1]}^{ggg} = \Gamma_{[-1,-1,1,1]}^{ggg} = \Gamma_4, \quad (\text{A7})$$

where $\Gamma_{1,2,3}$ each collect four equivalent helicity configurations, while Γ_4 collects only two. Performing the analytic integration over the internal momentum of the heavy quarkonium and the three-body phase space, the four classes admit the closed-form $\hat{q}^2$ -order expressions,

$$\Gamma_1 = \frac{5\alpha_s^3 N_V^2 \beta_V^3 (3\pi^2 - 16)}{216M\pi^{9/2}} \left(1 - \frac{7(1264 + 111\pi^2)}{6(3\pi^2 - 16)} \frac{\beta_V^2}{M^2} \right), \quad (\text{A8})$$

$$\Gamma_2 = \frac{5\alpha_s^3 N_V^2 \beta_V^3 (29\pi^2 - 280)}{216M\pi^{9/2}} \left(1 - \frac{7(265\pi^2 - 1424)}{6(29\pi^2 - 280)} \frac{\beta_V^2}{M^2} \right), \quad (\text{A9})$$

$$\Gamma_3 = \frac{5\alpha_s^3 N_V^2 \beta_V^3}{27M\pi^{9/2}} \left(1 - \frac{7(88 + 27\pi^2)}{12} \frac{\beta_V^2}{M^2} \right), \quad (\text{A10})$$

$$\Gamma_4 = \frac{20\alpha_s^3 N_V^2 \beta_V^3 (\pi^2 - 9)}{27M\pi^{9/2}} \left(1 - \frac{7(48 + 121\pi^2)}{48(\pi^2 - 9)} \frac{\beta_V^2}{M^2} \right). \quad (\text{A11})$$

For $\psi(2S)$ these $\hat{q}^2$ -order expressions exhibit the slow convergence of the relativistic expansion discussed in the main text; the phenomenologically improved expressions used in the main analysis are given in Eqs. (3.1)–(3.4).

-
- [1] G. T. Bodwin, E. Braaten, and G. P. Lepage, Rigorous QCD analysis of inclusive annihilation and production of heavy quarkonium, *Phys. Rev. D* **51**, 1125 (1995); *Phys. Rev. D* **55**, 5853(E) (1997).
- [2] N. Brambilla *et al.* (Quarkonium Working Group Collaboration), Heavy quarkonium physics, *arXiv:hep-ph/0412158*.
- [3] C. Adloff *et al.* (H1 Collaboration), Diffractive photoproduction of $\psi(2S)$ mesons at HERA, *Phys. Lett. B* **541**, 251 (2002).
- [4] I. Abt *et al.* (ZEUS Collaboration), Measurement of the cross-section ratio $\sigma_{\psi(2S)}/\sigma_{J/\psi(1S)}$ in exclusive photoproduction at HERA, *J. High Energy Phys.* **12** (2022) 164.
- [5] H. S. Shao, H. Han, Y. Q. Ma, C. Meng, Y. J. Zhang, and K. T. Chao, Yields and polarizations of prompt J/ψ and $\psi(2S)$ production in hadronic collisions, *J. High Energy Phys.* **05** (2015) 103.
- [6] H.-f. Fu, X.-j. Chen, and G.-L. Wang, Annihilation rates of heavy 1^{--} S-wave quarkonia in Salpeter method, *Phys. Lett. B* **692**, 312 (2010).
- [7] Z.-K. Geng, T. Wang, Y. Jiang, G. Li, X.-Z. Tan, and G.-L. Wang, Relativistic effects in the semileptonic B_c decays to charmonium with the Bethe-Salpeter method, *Phys. Rev. D* **99**, 013006 (2019).
- [8] K. Nochi, T. Kawanai, and S. Sasaki, Bethe-Salpeter wave functions of $\eta_c(2S)$ and $\psi(2S)$ states from full lattice QCD, *Phys. Rev. D* **94**, 114514 (2016).
- [9] V. Bertone, J.-P. Lansberg, and K. Lynch, Impact of relativistic corrections to high-pT prompt- $\psi(2S)$ production at hadron colliders, *arXiv:2510.06456*.
- [10] M. Copeland, L. Dai, Y. Fu, and J. Roy, $\psi(2S)$ production in jets using NRQCD, *J. High Energy Phys.* **06** (2026) 063.
- [11] Z.-G. He, X.-B. Jin, and B. A. Kniehl, Relativistic corrections to prompt double charmonium hadroproduction near threshold, *Phys. Rev. D* **109**, 094013 (2024).
- [12] N. Kivel, Relativistic corrections to $\psi(nS) \rightarrow \rho\pi$ exclusive decays and their role in the understanding of the $\rho\pi$ -puzzle, *Phys. Rev. D* **107**, 094015 (2023).
- [13] N. Kivel, Relativistic corrections to $J/\psi \rightarrow p\bar{p}$ decay, *Phys. Rev. D* **107**, 054026 (2023).
- [14] H.-M. Jiang, C.-J. Fan, J.-K. He, and C. Kong, Heavy quarkonium decay $V \rightarrow ggg$ with both relativistic and QCD radiative corrections, *Phys. Rev. D* **112**, 114014 (2025).
- [15] K.-T. Chao, H.-W. Huang, and Y.-Q. Liu, Gluonic and leptonic decays of heavy quarkonia and the determination of $\alpha_s(m_c)$ and $\alpha_s(m_b)$, *Phys. Rev. D* **53**, 221 (1996).
- [16] E. E. Salpeter and H. A. Bethe, A relativistic equation for bound state problems, *Phys. Rev.* **84**, 1232 (1951).
- [17] E. E. Salpeter, Mass corrections to the fine structure of hydrogen—like atoms, *Phys. Rev.* **87**, 328 (1952).
- [18] A. N. Mitra and S. Bhatnagar, Hadron—quark vertex function. Interconnection between 3-D and 4D wave function, *Int. J. Mod. Phys. A* **07**, 121 (1992).
- [19] H. Negash and S. Bhatnagar, Spectroscopy of ground and excited states of pseudoscalar and vector charmonium and bottomonium, *Int. J. Mod. Phys. E* **25**, 1650059 (2016).
- [20] S. Bhatnagar and L. Alemu, Approach to calculation of mass spectra and two-photon decays of $c\bar{c}$ mesons in the framework of Bethe-Salpeter equation, *Phys. Rev. D* **97**, 034021 (2018).
- [21] E. Mengesha and S. Bhatnagar, Cross section for double charmonium production in electron-positron annihilation at energy $\sqrt{s} = 10.6$ GeV, *Int. J. Mod. Phys. E* **20**, 2521 (2011).

- [22] H. Negash and S. Bhatnagar, Mass spectrum and leptonic decay constants of ground and radially excited states of η_c and η_b in a Bethe-Salpeter equation framework, *Int. J. Mod. Phys. E* **24**, 1550030 (2015).
- [23] E. Gebrehana, S. Bhatnagar, and H. Negash, Analytic approach to calculations of mass spectra and decay constants of heavy-light quarkonia in the framework of Bethe-Salpeter equation, *Phys. Rev. D* **100**, 054034 (2019).
- [24] J.-K. He and C.-J. Fan, Revisiting the P -wave charmonium radiative decays $h_c \rightarrow \gamma\eta^{(\prime)}$ with relativistic corrections, *Phys. Rev. D* **103**, 114006 (2021).
- [25] S. Bhatnagar, S.-Y. Li, and J. Mahecha, Power counting of various Dirac covariants in hadronic Bethe-Salpeter wave functions for decay constant calculations of pseudoscalar mesons, *Int. J. Mod. Phys. E* **20**, 1437 (2011).
- [26] S. Bhatnagar, J. Mahecha, and Y. Mengesha, Relevance of various Dirac covariants in hadronic Bethe-Salpeter wave functions in electromagnetic decays of ground state vector mesons, *Phys. Rev. D* **90**, 014034 (2014).
- [27] C. H. Llewellyn-Smith, A relativistic formulation for the quark model for mesons, *Ann. Phys. (N.Y.)* **53**, 521 (1969).
- [28] G.-L. Wang, Decay constants of heavy vector mesons in relativistic Bethe-Salpeter method, *Phys. Lett. B* **633**, 492 (2006).
- [29] R. Alkofer, P. Watson, and H. Weigel, Mesons in a poincare covariant Bethe-Salpeter approach, *Phys. Rev. D* **65**, 094026 (2002).
- [30] S. Bhatnagar and S.-Y. Li, Generalized structure of hadron-quark vertex function in Bethe-Salpeter framework: Applications to leptonic decays of V -mesons, *J. Phys. G* **32**, 949 (2006).
- [31] G. T. Bodwin and A. Petrelli, Order- v^4 corrections to S -wave quarkonium decay, *Phys. Rev. D* **66**, 094011 (2002); *Phys. Rev. D* **87**, 039902(E) (2013).
- [32] M. Ablikim *et al.* (BESIII Collaboration), Polarization and entanglement in baryon-antibaryon pair production in electron-positron annihilation, *Nat. Phys.* **15**, 631 (2019).
- [33] M. Ablikim *et al.* (BESIII Collaboration), First observation of Λ hyperon transverse polarization in $\psi(3686) \rightarrow \Lambda\bar{\Lambda}$, [arXiv:2509.15276](#).
- [34] W. Kwong, P. B. Mackenzie, R. Rosenfeld, and J. L. Rosner, Quarkonium annihilation rates, *Phys. Rev. D* **37**, 3210 (1988).
- [35] S. Navas *et al.* (Particle Data Group), Review of particle physics, *Phys. Rev. D* **110**, 030001 (2024).
- [36] X.-Z. Weng, L.-Y. Xiao, W.-Z. Deng, X.-L. Chen, and S.-L. Zhu, Three body open flavor decays of higher charmonium and bottomonium, *Phys. Rev. D* **99**, 094001 (2019).
- [37] D. Besson *et al.* (CLEO Collaboration), Measurement of the direct photon momentum spectrum in $v(1S)$, $v(2S)$, and $v(3S)$ decays, *Phys. Rev. D* **74**, 012003 (2006).
- [38] J. Libby *et al.* (CLEO Collaboration), Inclusive radiative $\psi(2S)$ decays, *Phys. Rev. D* **80**, 072002 (2009).
- [39] T. Peng and B.-Q. Ma, Heavy quarkonium $2S$ states in light-front quark model, *Eur. Phys. J. A* **48**, 66 (2012).
- [40] H.-M. Choi, C.-R. Ji, Z. Li, and H.-Y. Ryu, Variational analysis of mass spectra and decay constants for ground state pseudoscalar and vector mesons in the light-front quark model, *Phys. Rev. C* **92**, 055203 (2015).
- [41] A. J. Arifi, H.-M. Choi, C.-R. Ji, and Y. Oh, Mixing effects on $1S$ and $2S$ state heavy mesons in the light-front quark model, *Phys. Rev. D* **106**, 014009 (2022).
- [42] R. Acharyya, S. Puan, H. Dahiya, and N. Kumar, Spectroscopy of excited quarkonium states in the light-front quark model, *Chin. Phys. C* **49**, 023104 (2025).
- [43] S. Godfrey and K. Moats, Bottomonium mesons and strategies for their observation, *Phys. Rev. D* **92**, 054034 (2015).
- [44] I. Asghar and N. Akbar, Spectrum and decay properties of bottomonium mesons, *Eur. Phys. J. A* **60**, 58 (2024).
- [45] X.-L. Gao, J.-X. Cui, Y.-H. Zhou, and Z.-Y. Zhou, Updated analysis of charmonium states in a relativized quark potential model, [arXiv:2504.14575](#).
- [46] Z. Ahmad, I. Asghar, B. Masud, and M. A. Sultan, Charmonium spectrum and its decay properties, *Eur. Phys. J. A* **61**, 200 (2025).