Heavy mesons from the QCD instanton vacuum beyond the static limit

Ki-Hoon Hong,[1,*] Yongwoo Choi,[2,3,†] Nurmukhammad Rakhimov,[1,‡] and Hyun-Chul Kim[4,1,5,§]

[1]*Institute of Quantum Science, Inha University, Incheon 22212, Republic of Korea*
[2]*School of AIT, Hoseo University, Asan 31499, Republic of Korea*
[3]*Quantum AI Innovative Technology Institute (Q-AIT), Hoseo University, Asan 31499, Republic of Korea*
[4]*Department of Physics, Inha University, Incheon 22212, Republic of Korea*
[5]*School of Physics, Korea Institute for Advanced Study (KIAS), Seoul 02455, Republic of Korea*

(Received 30 June 2026; accepted 30 September 2026; published 27 October 2026)

We study pseudoscalar heavy mesons in the QCD instanton vacuum beyond the static limit. Finite-mass effects in the heavy-light loop are encoded in a separable effective vertex built from a profile function $\phi(\vec{p})$, kept distinct from the static Wilson-line form factor $F_Q^{(\infty)}(\vec{q})$ of the $m_Q \to \infty$ limit. The pseudoscalar two-point function fixes the residual mass Λ and the residue-normalized meson-quark coupling, from which we evaluate the decay constant and the spin-independent kinetic matrix element including the retained $1/m_Q$ derivative and kinetic contributions. The Isgur-Wise function is evaluated separately in the static heavy-quark limit, and its zero-recoil slope is extracted by using the same residue-normalized heavy-light vertex. The subleading calculation is restricted to the kinetic (derivative) part of the heavy-quark effective theory operators. For a representative vertex, calibrated at fixed $m_b^{\rm eff} = 5.04$ GeV to the leading-order decay constant $f_B^{\rm LO} = 180.0$ MeV and the spin-averaged mass $\bar{m}_B = 5.313$ GeV, we obtain $\delta f_B^{(1/m_Q,\rm partial)} = +121.9$ MeV and $f_B^{\rm partial} = 301.9$ MeV, $\Lambda = 190.7$ MeV, $\lambda_1^{(\partial)} = -0.773$ GeV^2, and $\rho_{\rm IW}^2 = 1.153$. The kinetic contribution yields a mass shift of order $\Lambda/2$ and a sizable partial $1/m_Q$ correction, indicating that the spin-independent nonperturbative $1/m_Q$ sector is a sensitive probe of the finite-mass heavy-light vertex.

DOI: 10.1103/83wz-blj7

I. INTRODUCTION

Heavy mesons containing a single heavy quark provide a standard system for studying how QCD separates the short-distance physics at the scale m_Q from the nonperturbative dynamics at hadronic scales. In the heavy-quark mass limit $m_Q \to \infty$, the spin of the heavy quark decouples from the gluon field, so that the heavy-quark spin and the total angular momentum of the light degrees of freedom are separately conserved. The dynamics also become independent of the heavy-quark flavor, since the heavy quark enters only as a static color source. These properties constitute the heavy-quark spin-flavor symmetry [1–5]. In this limit, the internal dynamics of a heavy hadron is governed entirely by the light degrees of freedom. Because m_Q is much larger than the light-quark masses and the typical hadronic scale, $1/m_Q$ serves as a small expansion parameter. A systematic expansion of the heavy-quark QCD Lagrangian in $1/m_Q$ yields heavy-quark effective theory (HQET) [6–9].

In HQET, the perturbative contributions to matrix elements of heavy hadrons are contained in the Wilson coefficients, whereas the long-distance contributions must be supplied by nonperturbative methods [6–9]. The QCD instanton vacuum provides one such framework for computing these nonperturbative contributions. It realizes the spontaneous breakdown of chiral symmetry (SBχS) in QCD [10–12] (see also the reviews in Refs. [13,14]). The mechanism of the SBχS can be explained as follows: a quark propagates through the instanton medium, which consists of instantons (I) and anti-instantons ($\bar{I}$). As the quark moves between random Is and $\bar{I}$s, its helicity undergoes a change due to the different handedness of the I and $\bar{I}$ zero modes. This process yields the SBχS, and the quark acquires a dynamical quark mass, $M(k)$, which is momentum dependent. $M(k)$ comes from the Fourier transform of the fermionic zero mode [12,14]. The value at zero virtuality, $M(0) \sim 350$ MeV, is determined by the saddle-point equation. The effective low-energy QCD partition function derived from the instanton vacuum has been successfully employed to describe low-lying light hadrons [15–24].

*Contact author: kihoon.hong@inha.ac.kr
†Contact author: ywchoi@hoseo.edu
‡Contact author: n.r.rakhimov@gmail.com
§Contact author: hchkim@inha.ac.kr

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

This theoretical framework was subsequently extended to systems that include heavy quarks [25–31]. A heavy quark interacts with the background gauge field through a Wilson line, and its coupling to the instanton–anti-instanton ensemble fixes the instanton contributions to the heavy-quark mass and the heavy-quark potential. Combining the heavy quark with the light quarks yields an effective heavy-light quark interaction. This interaction is nonlocal, so that the light and heavy quarks each carry a form factor: the light-quark form factor $F(p)$ generates the momentum-dependent dynamical mass [12], while the Wilson line gives the static heavy-line form factor $F_Q^{(\infty)}(\vec{q})$ in the $m_Q \to \infty$ limit [31].

In our previous work [31], the momentum dependence of the heavy-line form factor was neglected by setting $F_Q^{(\infty)}(\vec{q}) \to F_Q^{(\infty)}(\vec{0}) = 1$. The normalized static Wilson-line form factor $F_Q^{(\infty)}(\vec{q})$ describes the momentum transfer along the heavy line in the $m_Q \to \infty$ limit. The present bound-state loop calculation, however, requires the nonlocal heavy-light vertex as a function of the relative three-momenta at the two heavy-light vertices. We therefore represent the finite-mass heavy-light kernel by the separable effective form

$$\mathcal{F}(\vec{p},\vec{l};m_Q) \simeq \frac{\phi(\vec{p})\phi(\vec{l})}{N_\phi^2}, \tag{1}$$

where $\phi(\vec{p})$ denotes the finite-mass heavy-light vertex profile and N_ϕ stands for a momentum-independent normalization factor. The two heavy-line functions play different roles: $F_Q^{(\infty)}(\vec{q})$ depends on the momentum transfer along the static heavy line, whereas $\phi(\vec{p})$ depends on the relative momentum at a heavy-light vertex. The pole condition and its residue fix the interaction strength used below. We then calculate the residual mass Λ, the pseudoscalar decay constant, the spin-independent kinetic matrix element, and the Isgur-Wise slope for a single light flavor. The explicit $1/m_Q$ calculation is limited to the ordinary-derivative part of the subleading current and kinetic insertion. Gauge-field terms in the covariant derivatives and the chromomagnetic operator require the corresponding effective gluonic operators and are not included here.

We sketch the structure of the present work: Sec. II provides the nonlocal heavy-light interaction, its separable form, and the pole-residue normalization. Sections III and IV contain the decay-constant and kinetic calculations, respectively. The Isgur-Wise function is discussed in Sec. V, and the numerical analysis is given in Sec. VI.

II. HEAVY-LIGHT INTERACTION FROM THE INSTANTON VACUUM

We start from the effective heavy-light quark action for the light flavor number $N_f = 1$ derived from the instanton vacuum [31], as follows:

$$S_{\rm eff} = S_0 + S_{\rm int}, \tag{2}$$

$$S_0 = -\int d^4x [q^\dagger(x)(i\partial\!\!\!/ + iM(i\partial))q(x) + h^\dagger(x)(iv\cdot\partial)h(x)], \tag{3}$$

$$S_{\rm int} = -M_q \int d^4z \int \frac{d^4p_q}{(2\pi)^4}\frac{d^4l_q}{(2\pi)^4} e^{-i(p_q-l_q)\cdot z} F(p_q)F(l_q) \int d^4x d^4y \frac{1}{N_c}\mathrm{tr}_c\langle x|\theta^{-1}(w-\theta)\theta^{-1}|y\rangle \times \left[-(q^\dagger(p_q)q(l_q))(h^\dagger(x)h(y)) + \frac{1}{8}(h^\dagger(x)\Gamma_i q(l_q))(q^\dagger(p_q)\Gamma_i h(y))\right], \tag{4}$$

where $\Gamma_i = \{\mathbf{1}, \gamma_5, \gamma_\mu, i\gamma_\mu\gamma_5, \sigma_{\mu\nu}/\sqrt{2}\}$, and S_0 and $S_{\rm int}$ represent the free and interaction actions, respectively. The fields q and h denote the corresponding light and heavy quark fields; a heavy-flavor label can be attached to h when needed. $M(i\partial)$ stands for the dynamical quark mass for the light quark, $M(k) = M_q F^2(k)$ with $M_q = 345$ MeV [12,14], where $F(k)$ is given by

$$F(k) = -z\frac{d}{dz}[I_0(z)K_0(z) - I_1(z)K_1(z)]|_{z=\bar{\rho}k/2}, \tag{5}$$

where $I_n(z)$ and $K_n(z)$ are the modified Bessel functions of the first and second kinds of order n, respectively, and $\bar{\rho}$ is the average instanton size. In the heavy-quark mass limit $m_Q \to \infty$, the action exhibits heavy-quark spin-flavor symmetry [4–6]. The interaction term in Eq. (4) is given as a nonlocal four-point interaction between heavy-light quarks, where the quark form factor $F(k)$ is attached to each light quark, whereas heavy quarks are dressed by a Wilson-line form factor [25,31]. Here, $\theta^{-1}(= d/dt)$ is the free static-heavy-quark propagator, and $w = (\theta^{-1} - iA_{I\mu}\dot{x}_\mu(t))^{-1}$ is the corresponding propagator in the field of a single (anti-) instanton [25]. The notation $(q^\dagger q) = q^\dagger_{ai}q^{ai}$ indicates contraction over spin and color indices, and the four-fermion terms arise from Fierz reordering in color and spin spaces. The overall signs in Eqs. (3) and (4) follow from the Wick rotation of the Minkowski action to the Euclidean one used throughout this work. The propagators and vertices below are defined consistently with this convention. Details of the

static Wilson-line kernel entering the heavy-light interaction are provided in the Appendix. In the HQET limit, retaining only the leading contribution in the residual energies $\omega_{1,2} \sim \Lambda_{\rm QCD}$ yields a normalized momentum-space form factor $F_Q^{(\infty)}(\vec{q})$, which characterizes the momentum dependence of the static reference kernel. Using this result, the heavy-light interaction can be written as

$$S_{qQ} = +g^2 \int \frac{d^4p_q}{(2\pi)^4}\frac{d^4l_q}{(2\pi)^4}\frac{d^4k_1}{(2\pi)^4}\frac{d^4k_2}{(2\pi)^4}(2\pi)^4 \times \delta^{(4)}(p_q - l_q + k_1 - k_2)F(p_q)F(l_q)F_Q^{(\infty)}(\vec{k}_1 - \vec{k}_2) \times \sum_i [q^\dagger(-p_q)\Gamma_i h(k_1)][h^\dagger(k_2)\Gamma_i q(-l_q)], \quad (6)$$

where the coupling constant is defined as $g^2 \equiv M_q \Delta M_Q/(16n)$, and the momentum-dependent dynamical quark mass is given by $M(p_q) = \frac{(2\pi\bar{\rho})^2\lambda}{N_c}F^2(p_q) \equiv M_q F^2(p_q)$. Here, we redefine $T(\vec{q}) \equiv -\frac{\Delta M_Q}{2n}F_Q^{(\infty)}(\vec{q}=\vec{k}_1-\vec{k}_2)$, using the instanton contribution $\Delta M_Q = \frac{16\pi n\bar{\rho}^3}{N_c}\int_0^\infty d\tilde{r}\tilde{r}^2 F_Q^{(\infty)}(\tilde{r}) = 68$ MeV to the heavy quark mass and the instanton density $n = N/V = (200\ \text{MeV})^4$, as follows:

$$F_Q^{(\infty)}(\vec{q}) = \frac{\int_0^\infty d\tilde{r}\,\tilde{r}^2 j_0(\bar{\rho}|\vec{q}|\tilde{r})F_Q^{(\infty)}(\tilde{r})}{\int_0^\infty d\tilde{r}\,\tilde{r}^2 F_Q^{(\infty)}(\tilde{r})} \quad (7)$$

$$F_Q^{(\infty)}(\tilde{r}) = \cos^2\left(\frac{\pi\tilde{r}}{2\sqrt{\tilde{r}^2+1}}\right). \quad (8)$$

In Ref. [31], this momentum dependence was replaced by the static momentum approximation, $F_Q^{(\infty)}(|\vec{q}| \to 0) = 1$. The normalization in Eq. (7) is chosen such that $F_Q^{(\infty)}(\vec{q}=0) = 1$, so the Wilson-line form factor reduces to the static result in the forward limit. The position-space function $F_Q^{(\infty)}(\tilde{r})$ is fixed by the Wilson-line structure in the instanton background and depends only on $\tilde{r} = r/\bar{\rho}$.

The overall strength ΔM_Q is factored out, so $F_Q^{(\infty)}(\vec{q})$ describes only the momentum dependence generated by the Wilson line in the large-m_Q limit. Since this form factor is a function of the momentum transfer $\vec{q}$ along the heavy line, we do not identify it directly with the heavy-light vertex needed in the bound-state loop calculation. Instead, we replace the direct use of $F_Q^{(\infty)}(\vec{q})$ by the phenomenological separable ansatz

$$\mathcal{F}(\vec{p},\vec{l};m_Q) \simeq \frac{\phi(\vec{p})\phi(\vec{l})}{N_\phi^2}, \quad (9)$$

where $\vec{p}$ and $\vec{l}$ are the relative three-momenta at the two heavy-light vertices. Here $\phi(\vec{p})$ is the heavy-light vertex profile and N_ϕ is its dimensionless overall normalization factor. This ansatz supplies the vertex momentum dependence used in the calculation of the residual mass, decay constant, kinetic correction, and Isgur-Wise function.

In the convention used here, the normalized $F_Q^{(\infty)}(\vec{q})$, the effective finite-mass kernel $\mathcal{F}(\vec{p},\vec{l};m_Q)$, and $\phi(\vec{p})$ are dimensionless. Imposing the ansatz in Eq. (9), one can rewrite the interaction term in Eq. (6) in terms of heavy–light currents as follows:

$$S_{qQ}^{\rm sep} = +G_0^2 \sum_i \int \frac{d^4p_H}{(2\pi)^4} J_i(p_H)J_i^\dagger(p_H), \quad (10)$$

$$J_i(p_H) \equiv \int \frac{d^4p}{(2\pi)^4} q^\dagger(-p)\Gamma_i F(p)\phi(\vec{p})h(p_H - p), \quad (11)$$

$$J_i^\dagger(p_H) \equiv \int \frac{d^4l}{(2\pi)^4} h^\dagger(p_H - l)\Gamma_i F(l)\phi(\vec{l})q(-l), \quad (12)$$

where $S_{qQ}^{\rm sep}$ follows from Eq. (6) after applying the separable ansatz in Eq. (9), integrating over k_2, and redefining $p_H = p_q + k_1$, $p_q = p$, and $l_q = l$. The heavy-light current entering the bosonized theory is therefore defined by a single four-momentum integral over the relative momentum. The finite-mass heavy-line effect is encoded in the rest-frame vertex profile $\phi(\vec{p})$. Before meson-field residue normalization, the separable interaction has the coefficient $G_0^2 = g^2/N_\phi^2$, ($[G_0] = [g]$).

The separable interaction can be bosonized as

$$S_{qQ\Phi_v}^{\rm sep} = \sum_i \int \frac{d^4p_H}{(2\pi)^4}[\Phi_{iv}^\dagger(p_H)\Phi_{iv}(p_H) - G_0\{\Phi_{iv}^\dagger(p_H)J_i(p_H) - J_i^\dagger(p_H)\Phi_{iv}(p_H)\}]. \quad (13)$$

Here, the index i specifies the meson channel and the subscript v indicates the HQET velocity sector. The relative-momentum dependence resides in the current $J_i(p_H)$ through the factors $F(p)$ and $\phi(\vec{p})$, so the quadratic Hubbard-Stratonovich term is diagonal in the residual momentum of the heavy meson p_H. The dimensional rescaling associated with the physical heavy-meson residue is performed only after the pole residue is extracted below.

Using the light- and heavy-quark Lagrangian in Eq. (3) and integrating out the quark fields, we obtain the quadratic effective action for the physical HQET meson field in terms of the heavy-meson residual momentum p_H, as follows:

$$S_{\rm eff}^{(2)} = \sum_i \int \frac{d^4p_H}{(2\pi)^4}\Phi_{iv}^\dagger(p_H)[1 - G_0^2\Sigma_i(p_H)]\Phi_{iv}(p_H), \quad (14)$$

where the heavy-light loop function Σ_i is given by

$$\Sigma_i(p_H) = \int \frac{d^4p}{(2\pi)^4} |\phi(\vec{p})|^2 F^2(p) \times \mathrm{tr}_D[S_q(-p)\Gamma_i S_h(p_H - p)\Gamma_i]. \quad (15)$$

Here, $p = (\vec{p}, p_4)$ represents the internal light-quark loop momentum. In our convention, the light- and heavy-quark propagators are given by

$$S_q(p) = \frac{\not{p} + iM(p)}{p^2 + M^2(p)}, \qquad S_h(k) = \frac{1 + \not{v}}{2v \cdot k}. \quad (16)$$

Due to the structure of the HQET propagator, the dependence of the loop function on the residual momentum reduces to the scalar form $v \cdot p_H$, which directly determines the inverse meson propagator,

$$S_{H,i}^{-1}(v \cdot p_H) = 1 - G_0^2 \Sigma_i(v \cdot p_H). \quad (17)$$

A physical heavy meson is obtained when this inverse propagator vanishes. In Euclidean space, we set $p_H = i\Lambda v$, so that the pole position defines the residual-mass parameter Λ in the present effective theory.

For the pseudoscalar channel, $\Gamma_P = \gamma_5$, the pole condition is given by $G_0^2 \Sigma_P(i\Lambda) = 1$. By exploiting the reflection symmetry of the p_4 integration domain, the odd terms in p_4 vanish, and the remaining even part of the integrand can be written as

$$\mathcal{I}_P(p;\Lambda) = \frac{F^2(p)}{\Lambda^2 + p_4^2} \frac{2M(p)\Lambda - 2p_4^2}{|\vec{p}|^2 + p_4^2 + M^2(p)}, \quad (18)$$

where $F(p)$ and $M(p)$ denote $F(\sqrt{|\vec{p}|^2 + p_4^2})$ and $M(\sqrt{|\vec{p}|^2 + p_4^2})$, respectively. We then define

$$\mu_P(i\Lambda, p_4) \equiv \int \frac{d^3p}{(2\pi)^3} |\phi(\vec{p})|^2 \mathcal{I}_P(p;\Lambda), \quad (19)$$

so that

$$\Sigma_P(i\Lambda) = \int_0^\infty \frac{dp_4}{\pi} \mu_P(i\Lambda, p_4). \quad (20)$$

The pole residue fixes the dimensional rescaling from the bosonized field Φ_{P_v} to the canonically normalized HQET meson field. Near $v \cdot p_H = i\Lambda$, the inverse propagator behaves as

$$S_H^{-1} = 1 - G_0^2 \Sigma_P(v \cdot p_H) \simeq -G_0^2 \Sigma_P'(i\Lambda)(v \cdot p_H - i\Lambda), \quad (21)$$

where the prime denotes the derivative with respect to $v \cdot p_H$ at the pole. Since this derivative is taken with respect to a dimensionful variable, the combination $iG_0^2\Sigma_P'(i\Lambda)$ has mass dimension M^{-1}. We define the pole-residue factor

$$\mathcal{R}_H \equiv iG_0^2 \Sigma_P'(i\Lambda), \qquad [\mathcal{R}_H] = M^{-1}. \quad (22)$$

The Hubbard-Stratonovich kernel is dimensionless. A physical HQET inverse propagator, by contrast, is linear in the residual energy, so the pole residue $\mathcal{R}_H$ supplies the dimensional rescaling as follows:

$$H_v(p_H) = \sqrt{\frac{\mathcal{R}_H}{2}} \Phi_{P_v}(p_H),$$
$$\Phi_{P_v}(p_H) = \sqrt{\frac{2}{\mathcal{R}_H}} H_v(p_H). \quad (23)$$

The residue-normalized heavy-light coupling is then defined as

$$G \equiv \sqrt{\frac{2}{\mathcal{R}_H}} G_0. \quad (24)$$

Only the combination $G_0 = g/N_\phi$ survives in the separable interaction; N_ϕ itself need not be chosen. Once the profile and pole position are specified, the pole condition fixes G_0. For the field normalization, we set $\mathcal{R}_H = 2\ \mathrm{GeV}^{-1}$. This convention does not add another interaction strength or observable. It also gives $[G] = M^{-1/2}$ from $[G_0] = [\mathcal{R}_H] = M^{-1}$. With this definition, one has

$$iG^2 \Sigma_P'(i\Lambda) = 2. \quad (25)$$

Explicitly,

$$\Sigma_P'(i\Lambda) = -i \int_0^\infty \frac{dp_4}{\pi} \frac{\partial \mu_P(i\Lambda, p_4)}{\partial \Lambda} = -i \int_0^\infty \frac{dp_4}{\pi} \int \frac{d^3p}{(2\pi)^3} \frac{|\phi(\vec{p})|^2 F^2(p)}{(\Lambda^2 + p_4^2)^2} \frac{4p_4^2\Lambda + 2M(p)(p_4^2 - \Lambda^2)}{|\vec{p}|^2 + p_4^2 + M^2(p)}. \quad (26)$$

Using the pole condition, we give the prerescaling quantities

$$G_0^2 = \frac{1}{\Sigma_P(i\Lambda)}, \qquad \mathcal{R}_H = i\frac{\Sigma_P'(i\Lambda)}{\Sigma_P(i\Lambda)}, \quad (27)$$

whereas the residue-normalized coupling entering external meson matrix elements is

$$G^2 = \frac{2}{i\Sigma_P'(i\Lambda)}. \tag{28}$$

The heavy-light vertex for the matrix elements is given by

$$\mathcal{V}_{H,i}(p) = \frac{G}{\sqrt{N_c}} F(p)\phi(\vec{p})\Gamma_i,$$
$$\mathcal{V}_{H^\dagger,i}(p) = -\frac{G}{\sqrt{N_c}} F(p)\phi(\vec{p})\Gamma_i. \tag{29}$$

Each internal quark line carries the four-momentum integration with momentum conservation $(2\pi)^4\delta^{(4)}(p_H - p_q - k)$ at the vertex. The pole condition is written in terms of G_0, while decay constants, kinetic matrix elements, and heavy-to-heavy form factors are written in terms of the residue-normalized coupling G.

III. HEAVY-LIGHT CURRENT AND WEAK DECAY CONSTANTS

For a fixed profile, the pseudoscalar two-point function determines Λ and the residue-normalized heavy-light vertex. These quantities are then used in the matrix element of the axial-vector current that defines the pseudoscalar decay constant. We restrict the explicit calculation to the pseudoscalar channel. Note that vector states and hyperfine observables require the chromomagnetic sector, which is not included here.

At order $1/m_Q$, the axial-vector matrix element contains the derivative correction to the heavy-light current and the time-ordered insertion of the kinetic operator from the HQET Lagrangian. With the conventions used below, we write

$$\langle 0|J_\mu^A|P(v)\rangle = \langle 0|q^\dagger\gamma_\mu\gamma_5 h|P(v)\rangle - \frac{1}{2m_Q}\langle 0|q^\dagger\gamma_\mu\gamma_5 \not{D}_\perp h|P(v)\rangle$$
$$- \frac{1}{2m_Q}\int d^4y \langle 0|T\{q^\dagger\gamma_\mu\gamma_5 h(0), h^\dagger iD_\perp^2 h(y)\}|P(v)\rangle + O(1/m_Q^2). \tag{30}$$

In Eq. (30), we keep the ordinary-derivative projection $D_\perp \to \partial_\perp$. The gauge-field pieces are left to the effective gluonic operator sector. This projection affects the long-distance matrix elements but not the need for QCD-HQET current matching: the heavy-light axial-vector current is not conserved, and its short-distance scale dependence is carried by the Wilson coefficients. The matching relation is

$$J_\mu^A = \sum_i C_i(\mu_0)\left(J_\mu^{A(i)} - \frac{1}{2m_Q}(O_\mu^{A(i)} + T_\mu^{A(i)})\right) + \mathcal{O}\left(\frac{1}{m_Q^2}\right), \tag{31}$$

where $C_i(\mu_0)$ denote the Wilson coefficients of the HQET heavy-light axial-vector current at the low-energy scale μ_0 [6].

The flavor number entering the perturbative running of these Wilson coefficients is denoted by $n_f^{\rm run}$, to distinguish it from the single light flavor used in the instanton-induced effective interaction. Since $\mu_0 \ll m_Q$, we use the renormalization-group improved expressions [32]

$$C_1(\mu_0) = x^{2/\beta_0}\left[1 + \frac{\alpha_s(m_Q) - \alpha_s(\mu_0)}{4\pi} S_{\rm hl} - \frac{4}{3}\frac{\alpha_s(m_Q)}{\pi}\right], \tag{32}$$

$$C_2(\mu_0) = -\frac{2}{3}x^{2/\beta_0}\frac{\alpha_s(m_Q)}{\pi}, \qquad x \equiv \frac{\alpha_s(\mu_0)}{\alpha_s(m_Q)}, \tag{33}$$

where

$$S_{\rm hl} = \frac{\gamma_1^{\rm hl}}{2\beta_0} - \frac{\beta_1\gamma_0^{\rm hl}}{2\beta_0^2}, \qquad \gamma_0^{\rm hl} = -3C_F = -4, \tag{34}$$

and

$$\gamma_1^{\rm hl} = C_F\left[C_F\left(\frac{5}{2} - 16\zeta(2)\right) + C_A\left(-\frac{49}{6} + 4\zeta(2)\right) + \frac{10}{3}T_F n_f^{\rm run}\right]$$
$$= -\frac{254}{9} - \frac{56\pi^2}{27} + \frac{20}{9}n_f^{\rm run}, \tag{35}$$

with $C_F = \frac{4}{3}$, $C_A = 3$, and $T_F = \frac{1}{2}$ for $\mathrm{SU}(3)_c$. The running coupling at two-loop order is given by

$$\alpha_s(\mu) = \frac{4\pi}{\beta_0 \ln(\mu^2/\Lambda_{\rm QCD}^2)}\left[1 - \frac{\beta_1}{\beta_0^2}\frac{\ln\ln(\mu^2/\Lambda_{\rm QCD}^2)}{\ln(\mu^2/\Lambda_{\rm QCD}^2)}\right], \tag{36}$$

where

$$\beta_0 = 11 - \frac{2}{3}n_f^{\rm run}, \qquad \beta_1 = 102 - \frac{38}{3}n_f^{\rm run}. \tag{37}$$

Each current is defined as

$$J_\mu^{A(1)} \equiv q^\dagger\gamma_\mu\gamma_5 h, \qquad J_\mu^{A(2)} \equiv q^\dagger v_\mu\gamma_5 h, \tag{38}$$

$$O_\mu^{A(1)} \equiv q^\dagger \gamma_\mu \gamma_5 \not{D}_\perp h, \qquad O_\mu^{A(2)} \equiv q^\dagger v_\mu \gamma_5 \not{D}_\perp h, \tag{39}$$

$$T_\mu^{A(1)} \equiv \int d^4y\, T\{J_\mu^{A(1)}(0), h^\dagger(y) i D_\perp^2 h(y)\},$$
$$T_\mu^{A(2)} \equiv \int d^4y\, T\{J_\mu^{A(2)}(0), h^\dagger(y) i D_\perp^2 h(y)\}. \tag{40}$$

The standard dimension-four basis also contains $O^{A(3,\ldots,6)}$ [6]. They first enter at $O(\alpha_s/m_Q)$, beyond the matching accuracy used here, and are therefore not included. Below, F_1 denotes the retained derivative-current term and F_2 the kinetic insertion in f_B; $\lambda_1^{(\partial)}$ refers to the ordinary-derivative kinetic matrix element.

The combinations in Eq. (31) can be reduced to single matrix elements:

$$\sum_i C_i(\mu_0)\langle 0|J_\mu^{A(i)}|P(v)\rangle = [C_1(\mu_0) - C_2(\mu_0)]\langle 0|J_\mu^{A(1)}|P(v)\rangle, \tag{41}$$

$$\sum_i C_i(\mu_0)\langle 0|O_\mu^{A(i)}|P(v)\rangle = [C_1(\mu_0) + C_2(\mu_0)]\langle 0|O_\mu^{A(1)}|P(v)\rangle, \tag{42}$$

$$\sum_i C_i(\mu_0)\langle 0|T_\mu^{A(i)}|P(v)\rangle = [C_1(\mu_0) - C_2(\mu_0)]\langle 0|T_\mu^{A(1)}|P(v)\rangle. \tag{43}$$

Applying the Feynman rules, the matrix elements are represented by

$$-F_0 v_\mu \equiv \langle 0|J_\mu^{A(1)}|P(v)\rangle = -2G\sqrt{N_c} v_\mu \int_0^\infty \frac{dp_4}{\pi} \int_0^\infty dp |\vec{p}|^2 F(p)\phi(\vec{p}) \frac{p_4^2 - \Lambda M(p)}{2\pi^2(p_4^2+\Lambda^2)(|\vec{p}|^2 + p_4^2 + M^2(p))}, \tag{44}$$

$$-F_1 v_\mu \equiv \langle 0|O_\mu^{A(1)}|P(v)\rangle = -2G\sqrt{N_c} v_\mu \int_0^\infty \frac{dp_4}{\pi} \int_0^\infty dp |\vec{p}|^2 F(p)\phi(\vec{p}) \frac{-|\vec{p}|^2 \Lambda}{2\pi^2(p_4^2+\Lambda^2)(|\vec{p}|^2 + p_4^2 + M^2(p))}, \tag{45}$$

$$-F_2 v_\mu \equiv \langle 0|T_\mu^{A(1)}|P(v)\rangle = -2G\sqrt{N_c} v_\mu \int_0^\infty \frac{dp_4}{\pi} \int_0^\infty dp |\vec{p}|^2 F(p)\phi(\vec{p}) \frac{-|\vec{p}|^2(2p_4^2\Lambda + M(p)(p_4^2 - \Lambda^2))}{2\pi^2(p_4^2+\Lambda^2)^2(|\vec{p}|^2 + p_4^2 + M^2(p))}. \tag{46}$$

So, if we define $\mathcal{C}_{12}^{\pm} \equiv C_1 \pm C_2$, we get the decay constant $f_P = F_P/\sqrt{M_P}$, given by

$$f_P = \frac{1}{\sqrt{m_P}} \left[\mathcal{C}_{12}^{-}(\mu_0) F_0 - \frac{1}{2m_Q} (\mathcal{C}_{12}^{+}(\mu_0) F_1 + \mathcal{C}_{12}^{-}(\mu_0) F_2) \right]. \tag{47}$$

IV. KINETIC TERM FROM THE $1/M_Q$ CORRECTION

At order $1/m_Q$, the spin-independent kinetic operator measures the residual motion of the heavy quark inside the meson. In the present nonlocal model, it is weighted by the same vertex profile that enters the pole condition, so its ordinary-derivative matrix element $\lambda_1^{(\partial)}$ directly probes the momentum dependence of the finite-mass kernel. The HQET Lagrangian at this order is [6,8,33,34]

$$\mathcal{L}_{1/m_Q}^{\mathrm{HQET}} = +\frac{i}{2m_Q} h^\dagger \left(D_\perp^2 - \frac{1}{2}\sigma_{\mu\nu} G_{\mu\nu} \right) h, \tag{48}$$

where $D_{\perp\mu} \equiv D_\mu - v_\mu (v \cdot D)$ and $G_{\mu\nu}$ is the gluon field strength. The first term defines the full HQET kinetic parameter λ_1, while the second represents the chromomagnetic operator. We evaluate only the ordinary-derivative part of the kinetic term, $D_\perp^2 \to \partial_\perp^2$, and denote the corresponding matrix element by $\lambda_1^{(\partial)}$. The omitted gauge-field and chromomagnetic pieces require an effective instanton representation of gluonic operators, as in the light-quark case [35–37]. With this restriction, the pole shift from the $|\vec{k}|^2$ insertion is

$$\delta m_H^{(\partial)} = \frac{\delta S_H^{-1}}{\partial_{i\Lambda} S_H^{-1}} = \frac{\frac{1}{2m_Q} G_0^2 \Sigma_{\mathrm{kin}}(i\Lambda)}{-i G_0^2 \Sigma_P'(i\Lambda)} \equiv -\frac{1}{2m_Q} \lambda_1^{(\partial)}. \tag{49}$$

Here, δS_H^{-1} is the first-order change of the inverse meson propagator. Expanding the pole condition $S_H^{-1}(i\Lambda) = 0$ yields the ratio in Eq. (49), and the same pole slope that fixes the residue therefore also enters the kinetic mass shift.

In Eq. (49), $\lambda_1^{(\partial)}$ (derivative part) is defined as

$$\begin{aligned}\lambda_1^{(\partial)} &= \frac{1}{2}\langle H_v|h^\dagger i\partial_\perp^2 h|H_v\rangle = \frac{G^2}{2}\Sigma_{\rm kin}(i\Lambda)\\ &= i\frac{G^2}{2}\int\frac{d^4k}{(2\pi)^4}F^2(k)\phi^2(\vec{k})\mathrm{tr}_D[S_q(-k)\Gamma_P S_h(i\Lambda v-k)|\vec{k}|^2 S_h(i\Lambda v-k)\Gamma_P]\\ &= -\frac{G^2}{2}\int_0^\infty\frac{d|\vec{k}|}{2\pi^2}\int_0^\infty\frac{dk_4}{\pi}|\vec{k}|^2F^2(k)\phi^2(\vec{k})\frac{|\vec{k}|^2(-2M(k)\Lambda^2+4\Lambda k_4^2+2M(k)k_4^2)}{(|\vec{k}|^2+k_4^2+M^2(k))(\Lambda^2+k_4^2)^2}.\end{aligned} \tag{50}$$

The first line of Eq. (50) defines $\lambda_1^{(\partial)}$; the remaining lines give its loop representation. The kinetic insertion supplies the explicit $|\vec{k}|^2$ weight, making this quantity particularly sensitive to the high-momentum tail of $\phi(\vec{k})$. It should not be identified with the full λ_1 until the gauge-field part of $D_\perp^2$ is included.

V. ISGUR-WISE FUNCTION

Heavy-quark symmetry implies that all heavy-to-heavy semileptonic form factors are governed, at leading order in $1/m_Q$, by a single universal function of the recoil parameter $w = v\cdot v'$ [1,2,38,39]. A nontrivial check of any heavy-light bound-state model is therefore whether it reproduces the normalization $\xi(1) = 1$ at zero recoil and yields a reasonable slope $\rho_{\rm IW}^2 = -d\xi(w)/dw|_{w=1}$. In the present work we do not introduce separate finite-mass heavy-light vertex form factors for different heavy flavors. Instead, the Isgur-Wise function is evaluated directly in the static heavy-quark limit with the same heavy-light vertex form factor $\phi(\vec{p})$ that appears in the heavy-light two-point function.

The heavy-to-heavy transition is represented by the static current $-ih_{v'}^\dagger\Gamma_\mu h_v$, where v and v' denote the velocities of the initial and final static heavy fields. The corresponding Feynman diagram is shown in Fig. 1.

For the pseudoscalar case, the static-limit vertex representation gives

$$\begin{aligned}&-i\langle H(v')|h_{v'}^\dagger\Gamma_\mu h_v|H(v)\rangle\\ &= -iG^2\int\frac{d^4p_q}{(2\pi)^4}F^2(p_q)\phi^2(\vec{p}_q)\mathrm{tr}_D[\Gamma_P S_q(-p_q)\Gamma_P S_h(i\Lambda v'\\ &\quad -p_q)\Gamma_\mu S_h(i\Lambda v-p_q)].\end{aligned} \tag{51}$$

Here, the two heavy propagators are the HQET propagators of Eq. (16) with velocities v' and v, respectively. The same coupling G, residual mass Λ, and heavy-light vertex form factor $\phi(\vec{p})$ are used at both heavy-light vertices, because the quantity extracted here is the universal static-limit function rather than a finite-mass flavor-changing form factor.

To calculate the correlation function, we must define the velocities v and v'. In Euclidean convention, we use $v_M^0 = v_{E4}$ and $v_M^i = iv_{Ei}$, then the velocity satisfies $v_M^2 = v_E^2 = 1$ and $\frac{1+\not{v}_M}{2} = \frac{1+\not{v}_E}{2}$. If we set the Euclidean velocity $v = (\vec{0}, 1)$ as an initial state, then the final state in the center of mass frame has v' as

$$v' = (0, 0, \sqrt{1-w^2}, w), \tag{52}$$

where $v'^2 = 1$ and $w = v\cdot v'$. With the Euclidean normalization $v^2 = v'^2 = 1$, one has $w = \cos\theta$ and therefore $-1 \le w \le 1$ in purely Euclidean kinematics. Physical recoil corresponds to the Minkowski region $w \ge 1$. In this nonlocal model, however, a direct continuation of the Euclidean kinematics to $w > 1$ can lead to unstable or unphysical behavior because the heavy-line contribution and the nonlocal form factors become complex. For this reason, we evaluate the loop integrals only for $-1 \le w \le 1$ in Euclidean kinematics and extract the recoil dependence from the behavior around zero recoil.

Hence, we can rewrite the correlation function (51) as

$$\begin{aligned}&-i\langle H(v')|h_{v'}^\dagger\Gamma_\mu h_v|H(v)\rangle\\ &= -iG^2\mathrm{tr}_D\left[\gamma_5\frac{1+\not{v}'}{2}\Gamma_\mu\frac{1+\not{v}}{2}\gamma_5\mathcal{M}\right],\end{aligned} \tag{53}$$

where $\mathcal{M}$ is defined as




FIG. 1. Feynman diagram for the heavy-to-heavy form factor. The blob I is the nonlocal heavy-light vertex induced by the instanton vacuum.

$$\mathcal{M} \equiv \int \frac{d^4 p_q}{(2\pi)^4} \frac{-\not{p}_q + iM(p_q)}{p_q^2 + M^2(p_q)} \frac{F^2(p_q)\phi^2(\vec{p}_q)}{(i\Lambda - v\cdot p_q)(i\Lambda - v'\cdot p_q)}. \tag{54}$$

Having computed the trace over spin space, we obtain

$$-i\langle H(v')|h^\dagger_{v'}\Gamma_\mu h_v|H(v)\rangle = -iG^2 \int \frac{d^4 p}{(2\pi)^4} \frac{-(v'\cdot v - 1)p_\mu + [v_\mu(v'\cdot p) + v'_\mu(v\cdot p)] + iM(p)(v_\mu + v'_\mu)}{p^2 + M^2(p)} \times \frac{F^2(p)\phi^2(\vec{p})}{(i\Lambda - v\cdot p)(i\Lambda - v'\cdot p)}. \tag{55}$$

Guided by the standard HQET decomposition of the pseudoscalar heavy-to-heavy matrix element, we project the loop result onto the coefficient of $(v+v')_\mu$. Although the separable vertex $\phi(\vec{p})$ is implemented in the meson rest frame and does not make full covariance manifest, the leading heavy-quark-symmetry structure can be written as $F(v,v') = f_+(w)(v+v') + f_-(w)(v-v')$ [6,8]. Therefore, Eq. (55) is rewritten in this standard HQET basis. To obtain this form, the momentum p can be represented in the basis of v and v', as follows:

$$p_\mu = a(v+v')_\mu + b(v-v')_\mu + p^\perp_\mu, \tag{56}$$

where the perpendicular momentum $p^\perp$ satisfies $v\cdot p^\perp = 0$ and $v'\cdot p^\perp = 0$. Multiplying by $(v+v')$ and $(v-v')$, respectively, gives

$$a = \frac{(v+v')\cdot p}{2(1+v\cdot v')}, \qquad b = \frac{(v-v')\cdot p}{2(1-v\cdot v')}. \tag{57}$$

Then Eq. (55) is reduced to the following simple relation:

$$\begin{aligned} &-i\langle H(v')|h^\dagger_{v'}\gamma_\mu h_v|H(v)\rangle \\ &= -iG^2 \int \frac{d^4 p}{(2\pi)^4} \frac{F^2(p)\phi^2(\vec{p})}{(p^2 + M^2(p))(i\Lambda - v\cdot p)(i\Lambda - v'\cdot p)} \\ &\quad \times \left[iM(p) + \frac{(v+v')\cdot p}{1+v\cdot v'}\right](v+v')_\mu \\ &\equiv \xi(w)(v+v')_\mu, \end{aligned} \tag{58}$$

where $\xi(w)$ denotes the universal Isgur-Wise function in the present separable approximation. The result is generalized as

$$\langle H'(v')|h^\dagger_{v'}\Gamma_\mu h_v|H(v)\rangle = i\xi(w)\mathrm{Tr}\left[\Gamma' \frac{1+\not{v}'}{2}\Gamma_\mu \frac{1+\not{v}}{2}\Gamma\right]. \tag{59}$$

The decomposition in Eq. (56) defines the Isgur-Wise function as the coefficient of $(v+v')_\mu$ in the heavy-quark limit. At zero recoil, $v = v'$, and the current becomes the heavy-flavor symmetry generator. In this limit, Eq. (58) reduces, with the same Euclidean sign convention as in Eq. (21), to

$$\xi(1) = \frac{iG^2}{2}\Sigma'_P(i\Lambda). \tag{60}$$

The residue normalization $iG^2\Sigma'_P(i\Lambda) = 2$ then gives $\xi(1) = 1$. This is a nontrivial check for our framework because the same residue-normalized vertex coupling is used in both the two-point and three-point functions.

For nonzero recoil, the w dependence of $\xi(w)$ is determined by the heavy-light vertex form factor together with the nonlocal structure of the current insertion. In Euclidean kinematics, one may write $w_E = \cos\theta = 1 - \theta^2/2 + \cdots$, whereas the physical Minkowski recoil variable is $w_M = \cosh\eta = 1 + \eta^2/2 + \cdots$. Thus, the Euclidean and Minkowski regions approach the common zero-recoil point $w = 1$ from opposite sides. If the loop amplitude is analytic in a neighborhood of $w = 1$, the zero-recoil derivative is fixed unambiguously. We use the standard Isgur-Wise slope [40–42]

$$\rho^2_{\mathrm{IW}} \equiv -\left.\frac{d\xi(w)}{dw}\right|_{w=1}. \tag{61}$$

Equivalently, near zero recoil, the same analytic function is expanded as

$$\xi(w) = 1 - \rho^2_{\mathrm{IW}}(w-1) + O((w-1)^2). \tag{62}$$

In the Euclidean calculation, this expansion is evaluated for $w_E < 1$, so that

$$\begin{aligned} \xi_E(w_E) &= 1 - \rho^2_{\mathrm{IW}}(w_E - 1) + O((w_E-1)^2) \\ &= 1 + \rho^2_{\mathrm{IW}}(1 - w_E) + O((1-w_E)^2). \end{aligned} \tag{63}$$

Thus, the Euclidean derivative gives

$$\rho^2_{\mathrm{IW}} = -\left.\frac{d\xi_E(w_E)}{dw_E}\right|_{w_E = 1^-}. \tag{64}$$

This identification concerns the derivative with respect to w, not the coefficient of an expansion in the Euclidean

angle θ or the Minkowski rapidity η. The derivative is evaluated from Euclidean loop integrals in the interval $-1 \le w_E \le 1$.

VI. NUMERICAL RESULTS

We present a representative numerical solution of this framework. The instanton parameters are kept fixed at $M_q = 345$ MeV, $\bar{\rho}^{-1} = 600$ MeV, and $n = (200\ \text{MeV})^4$, with the standard dynamical-mass input discussed above [12,14]. For the heavy-quark sector, $m_b^{\text{eff}} = 5.04$ GeV is used in the leading mass relation, while $m_b = 4.695$ GeV is used in the short-distance matching and in the explicit $1/m_b$ terms. The finite-mass heavy-line kernel is represented by the same heavy-light vertex form factor that enters the pole condition, the decay constant, the kinetic correction, and the Isgur-Wise matrix element. For the numerical example, we use

$$\phi(\vec{k}) = \frac{1}{(1+|\vec{k}|^2/b^2)^a}. \tag{65}$$

Equation (65) is our central power-law (PL) choice. The separable ansatz does not determine the detailed momentum dependence of the finite-mass vertex, however, so using only one shape would leave that choice untested. We therefore refit two alternatives. The first is the rational (R) form

$$\phi_{\text{R}}(\vec{k}) = \frac{1}{1+(|\vec{k}|/b)^a}, \tag{66}$$

and, for a qualitatively different falloff, we use the Bessel-inspired (B) form

$$\phi_{\text{B}}(\vec{k}) = [\mathcal{B}(z_B)]^a, \qquad z_B \equiv \frac{|\vec{k}|}{2b}, \tag{67}$$

$$\mathcal{B}(z) \equiv 2z[I_0(z)K_1(z) - I_1(z)K_0(z)] - 2I_1(z)K_1(z),$$
$$\mathcal{B}(0) = 1. \tag{68}$$

Only the shape of $\mathcal{B}(z)$ is borrowed from the modified-Bessel structure of the instanton zero mode. Its parameters a and b are fitted anew. No extra light-quark factor $F(k)$ is introduced, so ϕ_{B} is simply another finite-mass heavy-light vertex profile. All three choices obey $\phi(0) = 1$.

Each profile is fitted independently. For PL, b fixes the momentum scale and a the power of the falloff. The pole condition used in the fit is

$$G_0^2 \Sigma_P(i\Lambda) = 1, \tag{69}$$

and the residue-normalized coupling

$$G^2 = \frac{2}{i\Sigma_P'(i\Lambda)}. \tag{70}$$

Equivalently, $\mathcal{R}_H = i\Sigma_P'(i\Lambda)/\Sigma_P(i\Lambda)$, and we impose $\mathcal{R}_H = 2\ \text{GeV}^{-1}$. Together with $f_B^{\text{LO}} = 180.0$ MeV and $\bar{m}_B = 5.313$ GeV, these conditions determine the profile parameters. Fits are restricted to $0.25 \le a \le 20$ and $0.015 \le b \le 3$ GeV.

TABLE I. Fitted parameters for the three finite-mass heavy-light vertex profiles. Each profile is calibrated to the same residue, f_B^{LO}, and spin-averaged mass conditions.

Profile	a	b [GeV]
PL	0.6560	0.1048
R	1.3631	0.0926
B	0.4103	0.0517

Table I gives the fitted a and b values; Fig. 2 shows the corresponding profiles.

Having refitted, we find that the three shapes yield very similar observables. Relative to PL, the largest shifts are 1.3% in Λ, 2.9% in $-\lambda_1^{(\partial)}$, 1.3% in ρ_{IW}^2, and 4.0% in the retained partial decay correction. We use these shifts as the functional-form spread of the vertex ansatz.

Table II summarizes the sensitivity to the instanton inputs, with the PL functional form retained. Our central choice is close to the conventional Diakonov-Petrov set [12,14]. Note that the set of Ref. [43] uses $M_q = 398$ MeV and $\bar{\rho} = 0.313$ fm. In each case a and b are refitted. Values from the two parametrizations are not mixed.

The partial decay correction hardly moves between the two sets: its change is 0.9%. By comparison, Λ changes by 2.7%, $-\lambda_1^{(\partial)}$ by 6.2%, and the recoil slope by 12.5%. Thus, ρ_{IW}^2 carries the largest input dependence in this test. We do not scan the instanton density independently. At fixed M_q




FIG. 2. Fitted PL, R, and B heavy-light vertex profiles. The three forms satisfy the same residue, f_B^{LO}, and spin-averaged mass conditions.

TABLE II. Sensitivity to two representative instanton-vacuum parameter sets used in the literature. The PL profile is independently refitted for each row to $\mathcal{R}_H = 2\ \mathrm{GeV}^{-1}$, $f_B^{\mathrm{LO}} = 180.0$ MeV, and $\bar{m}_B = 5.313$ GeV.

Set	M_q [MeV]	$\bar{\rho}$ [fm]	a	b [GeV]	Λ [MeV]	$-\lambda_1^{(\partial)}$ [GeV^2]	ρ^2_{IW}	$\delta f_B^{(1/m_Q,\mathrm{partial})}$ [MeV]
Present [12,14]	345	0.329	0.6560	0.1048	190.7	0.7730	1.1529	121.9
Ref. [43]	398	0.313	0.7446	0.1835	185.6	0.8211	1.2966	120.8

TABLE III. Representative numerical results. The quoted reference values are used only for orientation. The values taken from the literature may correspond to different mass schemes and conventions. In particular, the entries in the Λ column should be interpreted as the usual HQET parameter Λ in those references. The range quoted for $-\lambda_1$ is scheme-dependent and is taken from representative HQE/HQET analyses [44–46], while the Isgur-Wise slope bound is the QCD sum-rule lower bound [47]. The Belle II value is the experimentally fitted slope parameter at zero recoil [48].

Profile	f_B^{LO} [MeV]	$\delta f_B^{(1/m_Q,\mathrm{partial})}$ [MeV]	f_B^{partial} [MeV]	f_B^{ref} [MeV]	Λ [MeV]	m_b^{eff} [GeV]	$-\lambda_1^{(\partial)}$ [GeV^2]	ρ^2_{IW}
PL	180.0	+121.9	301.9	⋯	190.7	5.04	0.773	1.153
R	180.0	+119.8	299.8	⋯	189.6	5.04	0.783	1.145
B	180.0	+126.8	306.8	⋯	193.1	5.04	0.751	1.168
PDG [49]				190.0	⋯	⋯	⋯	⋯
Ref. [44]				⋯	314–588	4.70–4.90	0.26–0.97	⋯
Ref. [45]				⋯	240–320	5.02	0.11–0.17	⋯
Ref. [46]				⋯	⋯	4.518–4.564	0.336–0.492	⋯
Ref. [47]				⋯	⋯	⋯	⋯	> 0.75
Belle II [48]				⋯	⋯	⋯	⋯	1.09 ± 0.06

and $\bar{\rho}$, the explicit n dependence of $g^2 = M_q \Delta M_Q/(16n)$ cancels through $\Delta M_Q \propto n$, while G_0 is fixed by the pole condition.

The chromomagnetic sector is not included in this calculation, so the mass constraint is applied to the spin average

$$\bar{m}_B = \frac{m_B + 3m_{B^*}}{4}. \tag{71}$$

This removes the leading hyperfine splitting. With the heavy-quark inputs specified above, the fitted mass relation becomes

$$\bar{m}_B = m_b^{\mathrm{eff}} + \Lambda - \frac{\lambda_1^{(\partial)}}{2m_b}. \tag{72}$$

The same fit also imposes the residue condition and $f_B^{\mathrm{LO}} = 180.0$ MeV. The resulting quantities are collected in Table III.

Two entries in Table III should be distinguished from the phenomenological reference columns. The quantity f_B^{partial} contains only the retained derivative-current and kinetic insertions, so the Particle Data Group (PDG) value is shown separately as f_B^{ref}. Likewise, the quoted literature values of $-\lambda_1$ are for the full covariant HQET kinetic matrix element, not for our $-\lambda_1^{(\partial)}$.

For PL, the pole condition gives $\Lambda = 190.7$ MeV. Together with $\lambda_1^{(\partial)} = -0.773\ \mathrm{GeV}^2$, the retained kinetic piece contributes

$$-\frac{\lambda_1^{(\partial)}}{2m_b} \simeq 82.3\ \mathrm{MeV}, \tag{73}$$

which is approximately 43% of Λ. The retained partial decay-constant correction, $\delta f_B^{(1/m_Q,\mathrm{partial})} = +121.9$ MeV, amounts to approximately 68% of the leading-order value $f_B^{\mathrm{LO}} = 180.0$ MeV. These results show that the retained derivative sector is numerically important. Neither quantity, however, represents the complete correction at order $1/m_b$. Their magnitudes alone therefore do not establish a breakdown of the heavy-quark expansion; a quantitative assessment requires the omitted contributions at this order and an estimate of higher-order effects.

After independent refits to the same conditions, the PL, R, and B profiles yield partial decay-constant corrections between 119.8 and 126.8 MeV, with a maximum deviation of 4.0% from the PL result. Thus, the sizable partial correction persists across the three tested functional forms, with limited profile dependence under the adopted fitting conditions.

For the recoil dependence, the loop calculation fixes the zero-recoil slope. We therefore quote ρ^2_{IW} as the primary result. When an illustrative recoil-shape representation is needed, we use the slope-preserving parametrization

$$\xi_{\mathrm{fit}}(w) = \left(\frac{2}{1+w}\right)^{2\rho^2_{\mathrm{IW}}}. \tag{74}$$

With $\rho^2_{\rm IW} = 1.153$, this parametrization is normalized at zero recoil by construction. It should be understood only as a reconstruction from the fitted zero-recoil slope, not as an independent calculation of the full recoil dependence. The slope value satisfies the QCD sum-rule lower bound [47] and is close to the Belle II slope quoted in Table III.

VII. SUMMARY AND OUTLOOK

We have formulated a nonlocal instanton-based description of heavy-light mesons in which the static Wilson-line form factor $F_Q^{(\infty)}(\vec{q})$ is clearly separated from the effective finite-mass heavy-line vertex used in the calculation. The separable effective kernel represents the finite-mass vertex $\mathcal{F}(\vec{p},\vec{l};m_Q) \simeq \phi(\vec{p})\phi(\vec{l})/N_\phi^2$, where $\phi(\vec{p})$ is the dimensionless heavy-light vertex profile and N_ϕ is its dimensionless overall normalization factor. With this representation, the heavy-light current is written as a single four-momentum integral over the relative momentum, and the Hubbard-Stratonovich field is introduced as an auxiliary field depending only on the total momentum. At the same time, the nonlocal relative-momentum structure is kept in the current.

Only the combination $G_0 = g/N_\phi$ enters the separable interaction, so N_ϕ need not be fixed separately. The pseudoscalar pole condition $G_0^2\Sigma_P(i\Lambda) = 1$ fixes G_0, while the residue factor $\mathcal{R}_H = iG_0^2\Sigma_P'(i\Lambda)$ supplies the dimensional rescaling from the auxiliary bosonized field to the canonical HQET meson field. We impose $\mathcal{R}_H = 2\ \mathrm{GeV}^{-1}$ as this normalization condition and define $G = \sqrt{2/\mathcal{R}_H}G_0$. The same residue-normalized vertex $GF(p)\phi(\vec{p})\Gamma_i$ enters the decay constant, the kinetic $1/m_Q$ correction, and the heavy-to-heavy matrix element. The framework, therefore, treats the two-point function and the matrix elements within one common nonlocal structure while keeping the preresidue coupling G_0 distinct from the physical coupling G.

For the representative vertex form factor in Eq. (65), we obtained $f_B^{\rm LO} = 180.0\,\mathrm{MeV}$, $\delta f_B^{(1/m_Q,\mathrm{partial})} = +121.9$ MeV, $f_B^{\rm partial} = 301.9$ MeV, $\Lambda = 190.7$ MeV, $m_b^{\rm eff} = 5.04$ GeV (fixed) for $\bar{m}_B = 5.313$ GeV, $\lambda_1^{(\partial)} = -0.773\ \mathrm{GeV}^2$, and $\rho^2_{\rm IW} = 1.153$. The leading decay constant and the spin-averaged mass are used in the fit, while the remaining quantities follow from the fitted profile. The R and B profiles give only a few-percent spread around the central PL result: the largest changes are 1.3% in Λ, 2.9% in $-\lambda_1^{(\partial)}$, 1.3% in $\rho^2_{\rm IW}$, and 4.0% in the retained partial decay-constant correction. Changing to the alternative instanton-vacuum parameter set has a larger effect on the Isgur-Wise slope, which shifts by 12.5%, while the partial correction to f_B changes by less than 1%.

The kinetic contribution to the residual mass is about 43% of the leading residual contribution Λ, and the corresponding partial $1/m_Q$ correction to the decay constant is numerically sizable. This partial result should not be interpreted as a complete $O(1/m_b)$ prediction; rather, it shows that the retained spin-independent subleading sector is a sensitive probe of the finite-mass heavy-light vertex. The kinetic insertion is treated explicitly here because it is directly tied to the momentum dependence of the finite-mass heavy-light vertex. The remaining gauge-field and spin-dependent contributions can be incorporated through the corresponding effective gluonic operators and matching conditions within the same instanton-based framework. This extension is under investigation.

ACKNOWLEDGMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korean government under Grants No. RS-2025-02634319 (K.-H. H. and Y. C.) and No. RS-2025-00513982 (H.-C. K. and N. R.).

DATA AVAILABILITY

There are no publicly available research data or software supporting this manuscript. Requests for further information or data should be sent to the authors.

APPENDIX: WILSON-LINE KERNEL

Here, we specify the details of the Wilson-line calculation entering Eq. (4). In the heavy-quark static limit, the free heavy-quark propagator can be written in terms of the operator θ as

$$S_0(x,y) = \theta(x_4 - y_4)\delta^{(3)}(\vec{x} - \vec{y}), \tag{A1}$$

$$S_0^{-1}(x,y) = \delta^{(4)}(x-y)\partial_{y_4}, \tag{A2}$$

where $\theta(x_4 - y_4)$ denotes the Heaviside function. The propagator in the presence of a single instanton field A_I is

$$S_A(x,y) = \theta(x_4 - y_4)\delta^{(3)}(\vec{x} - \vec{y})W(\vec{x} - \vec{z}; x_4, y_4), \tag{A3}$$

$$W(\vec{x} - \vec{z}; x_4, y_4) = \mathcal{P}\exp\left[i\int_{y_4}^{x_4} d\tau\, A_{I4}(\vec{x} - \vec{z}, \tau - z_4)\right], \tag{A4}$$

where the instanton gauge field in singular gauge is given by

$$A_{I\mu}(x) = \frac{\bar{\eta}^a_{\mu\nu}\tau^a x_\nu \rho^2}{x^2(x^2+\rho^2)}, \tag{A5}$$

and only the temporal component A_{I4} enters the static Wilson line. Here $\bar{\eta}^a_{\mu\nu}$ denotes the 't Hooft symbol.

In momentum space, the heavy-quark contribution reads

$$\int d^4x d^4y h^\dagger(y)\mathrm{tr}_c[\langle y|\theta^{-1}(w-\theta)\theta^{-1}|x\rangle]h(x)$$
$$= \int \frac{d^4p_2}{(2\pi)^4}\frac{d^4p_1}{(2\pi)^4}h^\dagger(p_2)\int d^3x e^{-i(\vec{p}_2-\vec{p}_1)\cdot\vec{x}}\mathrm{tr}_c\left[\int dx_4\, dy_4 e^{-i\omega_{p_2}y_4+i\omega_{p_1}x_4}\langle y_4|\theta^{-1}(w-\theta)\theta^{-1}|x_4\rangle\right]h(p_1). \tag{A6}$$

To evaluate the Wilson-line operator, we use

$$\langle t_1|\theta|t_2\rangle = \theta(t_1-t_2), \qquad \langle t_1|\theta^{-1}|t_2\rangle = -\partial_{t_1}\delta(t_1-t_2). \tag{A7}$$

Inserting the completeness relation $\int d\tau|\tau\rangle\langle\tau| = 1$, one obtains

$$\int dx_4\, dy_4\, e^{-i\omega_{p_2}y_4+i\omega_{p_1}x_4}\langle y_4|\theta^{-1}\left(\int d\tau_2|\tau_2\rangle\langle\tau_2|\right)(w-\theta)\left(\int d\tau_1|\tau_1\rangle\langle\tau_1|\right)\theta^{-1}|x_4\rangle$$
$$= -\omega_{p_1}\omega_{p_2}e^{-i(p_2-p_1)\cdot z}\int_{-\infty}^{\infty}dT_1\int_{-\infty}^{\infty}dT_2\, e^{-i\omega_{p_2}T_2+i\omega_{p_1}T_1}\theta(T_2-T_1)\left\{\mathcal{P}\exp\left[i\int_{T_1}^{T_2}dr_4\frac{\bar{\eta}^a_{4\nu}\tau^a r_\nu\rho^2}{r^2(r^2+\rho^2)}\right]-1\right\}, \tag{A8}$$

where $r_\mu = (\vec{x}-\vec{z},\tau-z_4)$ and $T_{1,2} = \tau_{1,2}-z_4$.

After taking the color trace, Eq. (A8) becomes

$$-2\omega_{p_1}\omega_{p_2}e^{-i(p_2-p_1)\cdot z}\int_{-\infty}^{\infty}dT_1\int_{-\infty}^{\infty}dT_2\, e^{-i\omega_{p_2}T_2+i\omega_{p_1}T_1}\theta(T_2-T_1)\sin^2\left[\frac{1}{2}(\Omega(T_2)-\Omega(T_1))\right], \tag{A9}$$

with

$$\Omega(T) \equiv \tan^{-1}\left(\frac{T}{r}\right)-\frac{r}{\sqrt{r^2+\rho^2}}\tan^{-1}\left(\frac{T}{\sqrt{r^2+\rho^2}}\right). \tag{A10}$$

With the HQET momentum decomposition, $P_Q^\mu = m_Q v^\mu + k^\mu$ ($v^2=1$), where k^μ denotes the residual momentum and the residual energies satisfy $\omega_{1,2}\sim v\cdot k \ll m_Q$. Since the heavy quark remains close to its mass shell,

$$P_Q^2 = m_Q^2 + 2m_Q v\cdot k + k^2 \simeq m_Q^2, \tag{A11}$$

the Wilson-line kernel may be evaluated at $\omega_{1,2}=0$ to leading order in the HQET expansion.

In this limit, the characteristic time scale is $T\sim 1/\omega$, which is parametrically large. Since the instanton form factor suppresses hard contributions, the dominant support of the time integrals comes from the large-$|T|$ region. Therefore,

$$\lim_{T\to\pm\infty}\Omega(T) = \pm\frac{\pi}{2}\left(1-\frac{r}{\sqrt{r^2+\rho^2}}\right) \equiv \Omega_\infty(r)\mathrm{sgn}(T), \tag{A12}$$

so that

$$\Omega(T_2)-\Omega(T_1) \simeq \Omega_\infty(r)[\mathrm{sgn}(T_2)-\mathrm{sgn}(T_1)]. \tag{A13}$$

The difference vanishes when T_1 and T_2 have the same sign, so the only nonvanishing contribution comes from the region $T_1<0<T_2$. Consequently,

$$-2\omega_{p_1}\omega_{p_2}\int_{-\infty}^{0}dT_1\int_{0}^{\infty}dT_2\, e^{-i\omega_{p_2}T_2+i\omega_{p_1}T_1}\int d^3\, r\, e^{-i(\vec{p}_1-\vec{p}_2)\cdot\vec{r}}\sin^2[\Omega_\infty(r)]$$
$$= 8\pi\int_0^\infty dr\, r^2\, j_0(|\vec{p}_1-\vec{p}_2|r)\sin^2\left(\frac{\pi}{2}-\frac{\pi r}{2\sqrt{r^2+\rho^2}}\right) = 8\pi\int_0^\infty dr\, r^2\, j_0(|\vec{p}_1-\vec{p}_2|r)\cos^2\left(\frac{\pi r}{2\sqrt{r^2+\rho^2}}\right). \tag{A14}$$

The time integrations therefore reduce to the boundary contribution $W(\infty,-\infty,\vec{r})-1$, yielding the static momentum-space kernel

$$T(\vec{q})=\frac{1}{N_c}\int d^3\,r\,e^{i\vec{q}\cdot\vec{r}}\mathrm{tr}_c\langle\infty|(w-\theta)|-\infty\rangle=-\frac{8\pi}{N_c}\int_0^\infty dr\,r^2\,j_0(|\vec{q}|r)\cos^2\left(\frac{\pi r}{2\sqrt{r^2+\rho^2}}\right). \tag{A15}$$

Restoring the overall color normalization and the retarded/advanced sign convention gives the static Wilson-line kernel used in the main text.

[1] N. Isgur and M. B. Wise, Weak decays of heavy mesons in the static quark approximation, Phys. Lett. B **232**, 113 (1989).
[2] N. Isgur and M. B. Wise, Weak transition form factors between heavy mesons, Phys. Lett. B **237**, 527 (1990).
[3] H. Georgi, An effective field theory for heavy quarks at Low-energies, Phys. Lett. B **240**, 447 (1990).
[4] H. Georgi, Heavy quark effective field theory, in *Perspectives in the Standard Model: Proceedings of the Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 91)* (World Scientific, Singapore, 1992), pp. 589–628.
[5] M. B. Wise, Combining chiral and heavy quark symmetry, in *CCAST Symposium on Particle Physics at the Fermi Scale* (1993), pp. 71—114, arXiv:hep-ph/9306277.
[6] M. Neubert, Heavy quark symmetry, Phys. Rep. **245**, 259 (1994).
[7] M. A. Shifman, Lectures on heavy quarks in quantum chromodynamics, in Theoretical Advanced Study Institute in Elementary Particle Physics (TASI 95): QCD and Beyond (1995), pp. 409–514, arXiv:hep-ph/9510377.
[8] M. Neubert, Heavy quark effective theory, Subnucl. Ser. **34**, 98 (1997), arXiv:hep-ph/9610266.
[9] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, Phenomenology of heavy meson chiral Lagrangians, Phys. Rep. **281**, 145 (1997).
[10] E. V. Shuryak, The role of instantons in quantum chromodynamics. I. Physical vacuum, Nucl. Phys. **B203**, 93 (1982).
[11] D. Diakonov and V. Y. Petrov, Instanton based vacuum from Feynman variational principle, Nucl. Phys. **B245**, 259 (1984).
[12] D. Diakonov and V. Y. Petrov, A theory of light quarks in the instanton vacuum, Nucl. Phys. **B272**, 457 (1986).
[13] T. Schäfer and E. V. Shuryak, Instantons in QCD, Rev. Mod. Phys. **70**, 323 (1998).
[14] D. Diakonov, Instantons at work, Prog. Part. Nucl. Phys. **51**, 173 (2003).
[15] D. Diakonov, V. Y. Petrov, and P. V. Pobylitsa, A chiral theory of nucleons, Nucl. Phys. **B306**, 809 (1988).
[16] M. M. Musakhanov and H.-C. Kim, A test of the instanton vacuum with low-energy theorems of the axial anomaly, Phys. Lett. B **572**, 181 (2003).
[17] S.-i. Nam and H.-C. Kim, Electromagnetic form factors of the pion and kaon from the instanton vacuum, Phys. Rev. D **77**, 094014 (2008).
[18] S.-i. Nam and H.-C. Kim, Kaon semileptonic decay (K_{l3}) form factors from the instanton vacuum, Phys. Rev. D **75**, 094011 (2007).
[19] S.-i. Nam and H.-C. Kim, Generalized form factors and spin structures of the kaon, Phys. Lett. B **707**, 546 (2012).
[20] H.-D. Son, S.-i. Nam, and H.-C. Kim, Weak $K\to\pi$ generalized form factors and transverse transition quark-spin density from the instanton vacuum, Phys. Lett. B **747**, 460 (2015).
[21] S.-I. Shim and H.-C. Kim, Pion radiative weak decay from the instanton vacuum, Phys. Lett. B **772**, 687 (2017).
[22] S.-I. Shim, A. Hosaka, and H.-C. Kim, Vector and Axial-vector form factors in radiative kaon decay and flavor SU(3) symmetry breaking, Phys. Lett. B **795**, 438 (2019).
[23] C. V. Christov, A. Blotz, H.-C. Kim, P. Pobylitsa, T. Watabe, T. Meissner, E. Ruiz Arriola, and K. Goeke, Baryons as nontopological chiral solitons, Prog. Part. Nucl. Phys. **37**, 91 (1996).
[24] D. Diakonov, Chiral quark–soliton model, in: Advanced Summer School on Nonperturbative Quantum Field Physics (1997), pp. 1—55, arXiv:hep-ph/9802298.
[25] D. Diakonov, V. Y. Petrov, and P. V. Pobylitsa, The Wilson loop and heavy quark potential in the instanton vacuum, Phys. Lett. B **226**, 372 (1989).
[26] S. Chernyshev, M. A. Nowak, and I. Zahed, Heavy hadrons and QCD instantons, Phys. Rev. D **53**, 5176 (1996).
[27] U. T. Yakhshiev, H.-C. Kim, M. M. Musakhanov, E. Hiyama, and B. Turimov, Instanton effects on the heavy-quark static potential, Chin. Phys. C **41**, 083102 (2017).
[28] U. Yakhshiev, H.-C. Kim, and E. Hiyama, Instanton effects on charmonium states, Phys. Rev. D **98**, 114036 (2018).
[29] M. Musakhanov, N. Rakhimov, and U. T. Yakhshiev, Heavy quark correlators in an instanton liquid model with perturbative corrections, Phys. Rev. D **102**, 076022 (2020).
[30] K.-H. Hong, H.-C. Kim, and U. Yakhshiev, Instanton effects on electromagnetic transitions of charmonia, Prog. Theor. Exp. Phys. **2022**, 103D02 (2022).
[31] K.-H. Hong, H.-C. Kim, M. M. Musakhanov, and N. Rakhimov, Heavy-light quark systems from the QCD instanton vacuum: $N_f=1$ light flavor case, Phys. Rev. D **110**, 114044 (2024).
[32] M. Neubert, Short distance expansion of heavy–light currents at order 1/mQ, Phys. Rev. D **49**, 1542 (1994).

[33] A. F. Falk, B. Grinstein, and M. E. Luke, Leading mass corrections to the heavy quark effective theory, Nucl. Phys. **B357**, 185 (1991).
[34] A. F. Falk, M. E. Luke, and M. J. Savage, Nonperturbative contributions to the inclusive rare decays $B \to X_s\gamma$ and $B \to X_s l^+ l^-$, Phys. Rev. D **49**, 3367 (1994).
[35] D. Diakonov, M. V. Polyakov, and C. Weiss, Hadronic matrix elements of gluon operators in the instanton vacuum, Nucl. Phys. **B461**, 539 (1996).
[36] M. V. Polyakov and C. Weiss, Mixed quark–gluon condensate from instantons, Phys. Lett. B **387**, 841 (1996).
[37] J. Balla, M. V. Polyakov, and C. Weiss, Nucleon matrix elements of higher twist operators from the instanton vacuum, Nucl. Phys. **B510**, 327 (1998).
[38] Z. Ligeti, Y. Nir, and M. Neubert, The Subleading Isgur-Wise form-factor $\xi_3(v\,v\prime)$ and its implications for the decays $\bar{B} \to D^{(*)}\ell\bar{\nu}$, Phys. Rev. D **49**, 1302 (1994).
[39] I. Caprini and M. Neubert, Improved bounds for the slope and curvature of $B \to D(*)\ell\nu$ form-factors, Phys. Lett. B **380**, 376 (1996).
[40] N. Isgur and M. B. Wise, Excited charm mesons in semileptonic $\bar{B}$ decay and their contributions to a bjorken sum rule, Phys. Rev. D **43**, 819 (1991).
[41] I. Caprini, Slope of the Isgur-Wise function from a QSSR constraint on the YBB couplings, Phys. Lett. B **339**, 187 (1994).
[42] M. Neubert, Two loop corrections to the Isgur-Wise function in QCD sum rules, Phys. Rev. D **47**, 4063 (1993).
[43] W.-Y. Liu, E. Shuryak, and I. Zahed, Glue in hadrons at medium resolution and the QCD instanton vacuum, Phys. Rev. D **110**, 054005 (2024).
[44] K. K. Jeong and C. S. Kim, Determination of HQET parameter λ_1 from inclusive semileptonic B meson decay spectrum, Phys. Rev. D **59**, 114019 (1999).
[45] V. Chernyak, What are the right values of Lambda-bar and the heavy quark kinetic energy?, Phys. Lett. B **387**, 173 (1996).
[46] P. Gambino and C. Schwanda, Inclusive semileptonic fits, heavy quark masses, and V_{cb}, Phys. Rev. D **89**, 014022 (2014).
[47] N. Uraltsev, New exact heavy quark sum rules, Phys. Lett. B **501**, 86 (2001).
[48] I. Adachi *et al.*, Determination of $|V_{cb}|$ using B → D$\ell\nu l$ decays at Belle II, Phys. Rev. D **112**, 112009 (2025).
[49] S. Navas *et al.*, Review of particle physics, Phys. Rev. D **110**, 030001 (2024).