

Composite quarks and leptons with embedded QCD

Benoît Assi^{1,2} and Bogdan A. Dobrescu¹

¹Particle Theory Department, *Fermilab*, Batavia, Illinois 60510, USA

²Department of Physics, *University of Cincinnati*, Cincinnati, Ohio 45221, USA



(Received 30 January 2025; accepted 18 September 2025; published 3 October 2025)

We construct a model of quark and lepton compositeness based on an $SU(15)$ gauge interaction that confines chiral preons, which are also charged under the weakly coupled $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge group. The breaking of the latter, down to the Standard Model group, is achieved by scalar $SU(15)$ bound states at a scale in the 30–100 TeV range. The embedding of the QCD gauge group in $SU(4)_{\text{PS}}$ slows down the running of α_s in the UV. We estimate the effects of the strongly coupled $SU(15)$ dynamics on the running of the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge couplings, which likely remain perturbative beyond the compositeness scale of about 10^3 – 10^4 TeV, and even above a unification scale. A composite vectorlike lepton doublet acquires a mass in the TeV range probed at future colliders, and an extended Higgs sector arises from six-preon bound states.

DOI: [10.1103/bns1-b9fy](https://doi.org/10.1103/bns1-b9fy)

I. INTRODUCTION

The quest for uncovering deeper building blocks of matter was at the heart of physics research throughout the 20th century, and yielded outstanding scientific breakthroughs. Since the beginning of this century, that quest has been wavering, probably due to the inherent difficulties of making progress towards understanding strongly coupled quantum field theories with chiral fermions [1]. Chiral gauge theories appear to be necessary for describing the quarks and leptons as bound states, given that all Standard Model (SM) fermions have chiral charges under the electroweak gauge group.

A recently proposed theory of quark and lepton compositeness [2] attempts to describe the observed properties of the quarks and leptons based on an $SU(15)_p$ gauge interaction that confines an anomaly-free set of chiral fermions, referred to as preons (for historical reasons, e.g., [3–5]). The latter include 15 fermions charged (the same way as a single generation of SM fermions) under the SM gauge group, $SU(3)_c \times SU(2)_W \times U(1)_Y$, which does not play a role in confining the preons. That theory has several successful features: it implies the existence of three generations of composite quarks and leptons with the SM gauge charges, it provides a dynamical origin for the SM Higgs doublet as a six-preon bound state, and it predicts the existence of certain composite vectorlike quarks and

leptons that could be searched for at the LHC and future colliders.

At the same time, quark and lepton compositeness based on $SU(15)_p$ is facing some challenges [6]. First, dimension-8 proton decay operators are likely induced at the compositeness scale, such that the latter must satisfy $\Lambda_{\text{pre}} \gtrsim 10^3$ TeV, with uncertainties of more than an order of magnitude due to unknown nonperturbative effects. Second, the presence of several composite vectorlike quarks implies that the QCD coupling loses asymptotic freedom above a scale of about 10 TeV, and barely remains perturbative up to Λ_{pre} . This suggests that $SU(3)_c$ should be embedded in a larger group, so that the gauge coupling has a smaller positive β function (i.e., slower running), or even a negative one (i.e., asymptotic freedom is recovered at higher scales).

Moreover, due to the composite vectorlike quarks and leptons, the $U(1)_Y$ coupling loses perturbativity at a scale above Λ_{pre} but far below the GUT scale. Although we cannot reliably predict what occurs in the region near the scale Λ_{pre} where the $SU(15)_p$ interaction is strongly coupled, a self-consistent description of the hypercharge gauge interaction requires embedding $U(1)_Y$ in a non-Abelian group. Only in that case the hypercharge β function becomes smaller or even negative, due to the negative contribution from the gauge boson self-coupling.

For the above reasons, in this paper we study a preonic $SU(15)_p$ theory in which the preons are also charged under the minimal non-Abelian extension of the SM gauge group: $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$. While the latter was proposed in [7] to provide a unified origin for quarks and leptons, we study it here as a weakly coupled non-Abelian

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

gauge group acting on preons as well as on preonic bound states. Embedding the QCD gauge group in $SU(4)_{\text{PS}}$ has the advantage that its gauge coupling runs slower at scales above the masses of the composite vectorlike quarks. Moreover, we show that the binding due to the additional gauge bosons may push the masses of several composite vectorlike fermions towards the Λ_{pre} scale, further softening the running of the gauge couplings.

We also show that the composite scalars formed as di-prebaryon states (akin to the deuteron in QCD) may provide the origin of $SU(4)_{\text{PS}} \times SU(2)_R$ breaking down to $SU(3)_c \times U(1)_Y$ at a scale $\Lambda_{\text{PS}} < \Lambda_{\text{pre}}$. While robust proofs of the mechanisms outlined here are not possible in the absence of a better understanding of strongly coupled chiral gauge theories, which so far eludes lattice field theory (for recent attempts, see [8]), the $SU(15)_p \times SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge theory proposed here appears to have the necessary ingredients for a realistic theory of quark and lepton compositeness.

The paper is organized as follows. In Sec. II we introduce the preonic model above the compositeness scale. In Sec. III we study the effective theory below the compositeness scale. In particular, we examine the spontaneous breaking of $SU(4)_{\text{PS}} \times SU(2)_R$ due to composite dynamics, followed by an analysis of the theory at scales below Λ_{PS} . In Sec. IV, we compute within this framework the running of the three gauge couplings up to the compositeness scale, and then investigate how the gauge couplings may evolve above Λ_{pre} . We summarize our findings and provide an outlook in Sec. V.

II. PREONIC THEORY

The theory proposed in this paper has the same preonic gauge group as the one analyzed in [2,6], but crucially it has a larger weakly coupled gauge group, which we will argue later on (Sec. III) that it is spontaneously broken down to the SM gauge group by the dynamics of the preonic gauge interaction.

A. Field content

We consider a preonic theory with the $SU(15)_p \times SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge symmetry. The $SU(15)_p$ group becomes strongly coupled at a scale Λ_{pre} , a few orders of magnitude above the electroweak scale. $SU(4)_{\text{PS}}$ is the weakly coupled group that embeds QCD, $SU(2)_L$ is the SM gauge group associated with the weak interaction, and $SU(2)_R$ is the weakly coupled gauge group that together with the $U(1)_{B-L}$ subgroup of $SU(4)_{\text{PS}}$ eventually breaks down to the SM hypercharge gauge group.

The preons are chiral fermions which transform nontrivially under $SU(15)_p$: several of them ($\Psi_W, \Psi_{W'}, \psi_1, \psi_2, \psi_3$) belong to the fundamental representation, and one fermion (Ω) belongs to the conjugate symmetric representation. Only the Ψ_W and $\Psi_{W'}$ fields are charged under

TABLE I. Field content of the preonic model. All fields (six chiral fermions and two scalars) belong to nontrivial representations of the confining $SU(15)_p$ group. Only the Ψ_W and $\Psi_{W'}$ preons carry charges of the weakly coupled $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge group.

Field	Spin	$SU(15)_p$	$SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$
Ψ_W	1/2	$\square$	(4, 2, 1)
$\Psi_{W'}$	1/2	$\square$	($\bar{4}$, 1, 2)
ψ_1, ψ_2, ψ_3	1/2	$\square$	(1, 1, 1)
Ω	1/2	$\overline{\square\square}$	(1, 1, 1)
$\mathcal{A}, \mathcal{A}'$	0	$\square$	(1, 1, 1)
		$\square$	

$SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$, as shown in Table I. This set of six chiral fermion representations is free of gauge anomalies. In particular, the $SU(15)_p$ gauge anomaly of Ω is canceled by 19 fermions transforming in the fundamental representation [9]; this is achieved by the ψ_1, ψ_2, ψ_3 fermions together with the 16 fundamentals of $SU(15)_p$ packaged in Ψ_W and $\Psi_{W'}$.

The only other elementary fields present at scales of the order of Λ_{pre} are two scalars ($\mathcal{A}, \mathcal{A}'$) in the conjugate antisymmetric representation of $SU(15)_p$, whose Yukawa couplings are ultimately responsible for the flavor dependence of the SM fermion masses (see Sec. III D). The Yukawa couplings of the antisymmetric scalars to preons can be written as

$$\mathcal{L}_{\text{Yuk}} = \lambda_{ij} \mathcal{A}_{ab} \psi_i^a \psi_j^b + \lambda'_{ij} \mathcal{A}'_{ab} \psi_i^a \psi_j^b + \text{H.c.}, \quad (2.1)$$

where $i, j = 1, 2, 3$ are preonic flavor indices, and $a, b = 1, \dots, 15$ are $SU(15)_p$ -color indices. The Yukawa couplings λ_{ij} form a complex symmetric matrix. Through an appropriate redefinition of the fermion fields—implementing a Takagi factorization—this matrix can be diagonalized with all entries (λ_i) real. In that field basis, λ'_{ij} generically form a complex symmetric matrix with both off-diagonal and diagonal nonzero entries.

B. Preonic dynamics

Given that the second Dynkin indices for the symmetric and the antisymmetric $SU(15)$ representations are $T_2(\Omega) = 17/2$ and $T_2(\mathcal{A}) = 13/2$, respectively [9], we find the following one-loop coefficient of the $SU(15)_p$ β function:

$$b_{15} = 55 - \frac{2}{3} \left(\frac{19}{2} + T_2(\Omega) \right) - \frac{2}{3} T_2(\mathcal{A}) = \frac{116}{3}. \quad (2.2)$$

Thus, $b_{15} > 0$ (i.e., negative β function), implying asymptotic freedom for the $SU(15)_p$ gauge interaction. Consequently, it is natural that $SU(15)_p$ is weakly coupled near the Planck scale (M_p), and that its gauge coupling

grows logarithmically towards lower scales until it becomes nonperturbative at Λ_{pre} .

The dynamical behavior of the $SU(15)_p$ theory at the scale Λ_{pre} cannot be decided (for now) through a direct computation. The reason is that this is a strongly coupled chiral gauge interaction, and the only tool available for first-principle based computations is lattice field theory, which currently cannot be applied to chiral theories. Nevertheless, there are consistency checks that indicate some possible dynamics. Specifically, there are three mutually excluded proposals in the literature for the dynamical outcomes of an $SU(N)_p$ gauge theory with chiral fermions in one conjugate symmetric two-tensor and $N + 4$ fundamental representations.

The first one, which we assume to be the correct one in Secs. III and IV, is that there is confinement at the scale Λ_{pre} , and that the chiral symmetry of the elementary fermions is unbroken. Even though the latter behavior is different from QCD, which spontaneously breaks the chiral symmetry of the quarks, it is a self-consistent dynamics [4,10–14] supported by both the 't Hooft anomaly matching condition and large N arguments [10,12].

The second proposal is that there is no confinement, and instead there are chiral condensates that break completely the $SU(N)_p$ gauge symmetry [14]. Intriguingly, the spectrum of massless fermions is similar to the case of confinement discussed above (almost consistent with the conjecture of complementarity [5,15]). In [14] it is argued that confinement and no chiral symmetry breaking is a more likely scenario than the formation of chiral condensates, based on a conjecture regarding the thermodynamic free energy applied to chiral gauge theories [13]. In [16], the opposite is argued based on discrete symmetry matching; however, that argument was disputed in [11] (for follow-ups, see [12,17]).

The third proposal is that there is confinement, but chiral condensates partially break the $SU(N)_p$ gauge symmetry [18]. This is based on supersymmetric dualities perturbed by small anomaly mediated supersymmetry breaking terms. As emphasized in [12,18], for larger supersymmetry breaking it is possible that a phase transition occurs, and the result obtained in [18] might not apply to the nonsupersymmetric theory. In fact, some evidence that this is the case is provided by the large N arguments: the phase identified in [18] is valid for *any* $N \geq 13$, while the derivation of the unbroken chiral symmetry is robust at $N \rightarrow \infty$ [10,12]. In our case, $N = 15$ seems large enough to satisfy the large N arguments, but for now other phases cannot be conclusively ruled out.

It is worth emphasizing that the theory studied here has additional ingredients compared to $SU(N)_p$ theory with a symmetric tensor and $N + 4$ fundamental representations: there are two antisymmetric scalars (see Table I) with masses presumably of the order of or slightly above Λ_{pre} ,

and there are three other gauge groups that are weakly coupled at Λ_{pre} . We will argue in Sec. III that their effects may be sufficient to induce small vacuum expectation values (VEVs) for composite scalars, and masses for the composite quarks and leptons.

Turning now to the running of the weakly coupled gauge couplings in the presence of preons (the running at scales below Λ_{pre} is analyzed in Sec. IV), the one-loop coefficient of the β function is $b_4 = -16/3$ for $SU(4)_{\text{PS}}$, and $b_2 = -38/3$ for both $SU(2)_L$ and $SU(2)_R$. As $b_4, b_2 < 0$, the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge interactions are not asymptotically free. Furthermore, the large values of $|b_4|$ and $|b_2|$ indicate that these interactions could become strongly coupled at a UV scale below M_P . Note though that $|b_4|$ is smaller than the analogous quantity for QCD, if $SU(3)_c$ were not embedded in $SU(4)_{\text{PS}}$. Therefore, the embedding pushes the UV scale to higher values. The same is true for the embedding of the hypercharge gauge group in $SU(4)_{\text{PS}} \times SU(2)_R$.

There are, however, mitigating effects due to $SU(15)_p$ dynamics near Λ_{pre} (see Sec. IV C) that decrease the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge couplings such that they may remain weakly coupled up to M_P . As it is difficult to estimate the size of these effects, we note that if the couplings become strongly coupled at a scale below M_P , then either there is some strongly coupled UV fixed point, or $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ must be embedded in a larger gauge group at a scale Λ_{422} that satisfies $\Lambda_{\text{pre}} < \Lambda_{422} < M_P$. That larger group can be a grand unified one, such as $SO(10)$, or just a product of groups with higher rank than $SU(4) \times SU(2)^2$.

III. BOUND STATES OF $SU(15)_p$

At the scale Λ_{pre} , the $SU(15)_p$ interactions become strong and form $SU(15)_p$ -singlet bound states. The chiral symmetry of the preons remains unbroken, so some fermionic bound states, referred to as chiral prebaryons, remain massless at this stage. The 't Hooft anomaly matching conditions [1] indicate that the only chiral prebaryons are three-preon states formed of one preon belonging to the $SU(15)$ conjugate symmetric tensor representation (Ω in our model) and two preons belonging to the fundamental representation ($\Psi_W, \Psi_{W'}$, or $\psi_i, i = 1, 2, 3$) [10,15]. Thus, the following three-preon states (which are two-component fermions) remain massless: $\Omega\Psi_W\psi_i$, $\Omega\Psi_{W'}\psi_i$, $\Omega\psi_i\psi_j$, $\Omega\Psi_W\Psi_{W'}$, $\Omega\Psi_W\Psi_W$, and $\Omega\Psi_{W'}\Psi_{W'}$, where $i, j = 1, 2, 3$ and $i \neq j$. Since the Ψ_W and $\Psi_{W'}$ preons are charged under the weakly coupled gauge group (see Table I), the chiral prebaryons belong to the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ representations displayed in Table II.

We label the chiral prebaryons as $\mathcal{P}$ with a lower index that specifies its $SU(4)_{\text{PS}}$ representation, and when this does not uniquely determine the prebaryon, the $SU(2)_L$ representation is also displayed after a comma, as shown

TABLE II. Chiral prebaryons of $SU(15)_p$, their preon content, and their representations under the weakly coupled gauge group. The preon flavor index is $i = 1, 2, 3$, so there are three $\mathcal{P}_4^i$ prebaryons and three $\mathcal{P}_{\bar{4}}^i$ prebaryons. The anticommutation property of the preons implies that the $\mathcal{P}_{1,1}^{ij}$ prebaryons are antisymmetric in the i, j indices, so there are also three of them.

Prebaryon	Preon content	$SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$
$\mathcal{P}_{15}, \mathcal{P}_{1,2}$	$\Omega\Psi_W\Psi_{W'}$	$(15, 2, 2), (1, 2, 2)$
$\mathcal{P}_{10}, \mathcal{P}_{6,3}$	$\Omega\Psi_W\Psi_W$	$(10, 1, 1), (6, 3, 1)$
$\mathcal{P}_{\bar{10}}, \mathcal{P}_{6,1}$	$\Omega\Psi_{W'}\Psi_{W'}$	$(\bar{10}, 1, 1), (6, 1, 3)$
$\mathcal{P}_4^i$	$\Omega\Psi_W\psi_i$	$3 \times (4, 2, 1)$
$\mathcal{P}_{\bar{4}}^i$	$\Omega\Psi_{W'}\psi_i$	$3 \times (\bar{4}, 1, 2)$
$\mathcal{P}_{1,1}^{ij}$	$\Omega\psi_i\psi_j$	$3 \times (1, 1, 1)$

in Table II. For example, both $\mathcal{P}_{6,3}$ and $\mathcal{P}_{6,1}$ belong to the six of $SU(4)_{\text{PS}}$, but the first prebaryon is a triplet under $SU(2)_L$ while the second one is an $SU(2)_L$ singlet. In the case of prebaryons that have one or two ψ_i preons as constituents, one or two flavor upper indices ($i, j = 1, 2, 3$) are also included.

At scales below the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ breaking, the $\mathcal{P}_4^i$ and $\mathcal{P}_{\bar{4}}^i$ prebaryons include three generations of SM quarks and leptons (see discussions in Secs. III C and III D). There are also six SM-singlet fermions, three in $\mathcal{P}_4^i$ and three in the $\mathcal{P}_{1,1}^{ij}$ prebaryons, which mix with the SM neutrinos. The remaining prebaryons, listed in the first three rows of Table II, are vectorlike under the SM gauge group, and acquire masses much larger than the electro-weak scale, as discussed below.

A. Scalar di-prebaryons

As the chiral symmetry of the preons is preserved, there are no light pionlike states bound by the $SU(15)_p$ interactions. We expect, however, that bound states of two or more chiral prebaryons form due to remnant interactions of $SU(15)_p$, somewhat similar to nuclear interactions in QCD. Scalars lighter than Λ_{pre} likely arise as bound states of two chiral prebaryons [2], and are referred to as di-prebaryons. Let us model the interaction between chiral prebaryons due to $SU(15)_p$ remnant effects as four-fermion contact terms in the Lagrangian:

$$-\frac{g_p^2}{\Lambda_{\text{pre}}^2}(\bar{\mathcal{P}}\sigma^\mu\mathcal{P})(\bar{\mathcal{P}}\sigma_\mu\mathcal{P}), \quad (3.1)$$

where the σ^μ matrices account for spin-1 premeson exchange between prebaryon Weyl spinors, g_p is a premeson coupling to the prebaryons, and we have not displayed the prebaryon indices. Additional operators involving four chiral prebaryons, such as tensor-tensor

couplings, are also expected to have important effects, but we will not further discuss them.

A Fierz transformation of (3.1) gives operators of the type $(\bar{\mathcal{P}}\bar{\mathcal{P}})(\mathcal{P}\mathcal{P})$, which are attractive and produce scalar $\mathcal{P}_\kappa\mathcal{P}_\lambda$ bound states (di-prebaryons), generically labeled $\phi_{\kappa-\lambda}$. Here κ and λ are prebaryon lower indices, according to the labeling in Table II. Since these interactions between chiral prebaryons are nonconfining, the low-energy effective theory includes Yukawa interactions of the di-prebaryons to their three-preon constituents:

$$-y_{\kappa-\lambda}\phi_{\kappa-\lambda}^\dagger\mathcal{P}_\kappa\mathcal{P}_\lambda, \quad (3.2)$$

where $y_{\kappa-\lambda}$ are nearly nonperturbative Yukawa couplings.

The squared-mass $M_{\kappa-\lambda}^2$ of the scalar bound state $\phi_{\kappa-\lambda}$ receives a positive contribution of the order of Λ_{pre}^2 due to the mass of the spin-1 premeson mediator, and a negative contribution of the order of $y_{\kappa-\lambda}^2\Lambda_{\text{pre}}^2/(2\pi^2)$ due to prebaryon loops involving the attractive Yukawa interaction. Thus, if the effective Yukawa coupling $y_{\kappa-\lambda}$ has a value near (but smaller than) a critical coupling $y_{\text{cr}} \approx \pi/\sqrt{2}$, then it is possible to have $0 < M_{\kappa-\lambda}^2 \ll \Lambda_{\text{pre}}^2$. A similar phenomenon has been studied [19] in the context of top quark condensation [20,21] and composite Higgs models [22], where a second-order chiral phase transition allows a large hierarchy between the scale of compositeness and the mass of the composite scalar.

Although the binding of these ϕ di-prebaryons is due mostly to $SU(15)_p$ remnant interactions, there are additional contributions from $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge interactions and scalar ($\mathcal{A}, \mathcal{A}'$) exchange. The additional contributions are larger when the constituent prebaryons belong to higher gauge representations or have large Yukawa couplings in Eq. (2.1). In those cases, the squared mass of ϕ may turn negative, implying that the di-prebaryons develop VEVs. If a di-prebaryon acquires a VEV, then its two prebaryon constituents form a Dirac fermion that acquire a vectorlike mass.

The binding potential $V(r)$ between two prebaryons $\mathcal{P}_\kappa$ and $\mathcal{P}_\lambda$ induced by one gauge boson exchange can be written as

$$V_{\kappa-\lambda}(r) \approx -\frac{1}{2r}(C_4^{\kappa-\lambda}\alpha_4 + C_L^{\kappa-\lambda}\alpha_L + C_R^{\kappa-\lambda}\alpha_R), \quad (3.3)$$

where $\alpha_4, \alpha_L, \alpha_R$ are the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ coupling constants at the Λ_{pre} scale. The binding coefficients C_4, C_L, C_R are given by the sum of the quadratic Casimirs of the gauge representations in which the two prebaryons transform, minus the quadratic Casimir of the $\phi_{\kappa\lambda}$ di-prebaryon (the $\mathcal{P}_\kappa - \mathcal{P}_\lambda$ bound state) [23]:

$$C_4^{\kappa-\lambda} = C_4(\mathcal{P}_\kappa) + C_4(\mathcal{P}_\lambda) - C_4(\phi_{\kappa-\lambda}), \quad (3.4)$$

and analogously for C_L and C_R . Thus, the most deeply bound states are gauge singlets, i.e., $\phi_{\kappa-\lambda}$ with

$C_4(\phi) = C_L(\phi) = C_R(\phi) = 0$, and are formed of prebaryons in representations with large quadratic Casimirs. In particular, the symmetric representations of $SU(4)_{\text{PS}}$, namely $\mathcal{P}_{10}$ and $\mathcal{P}_{\overline{10}}$, bind with $C_4^{10-\overline{10}} = 9$. As this binding is large, we assume that the ensuing di-prebaryon $\phi_{10-\overline{10}}$ acquires a large VEV, which induces a Dirac mass near Λ_{pre} for $\mathcal{P}_{10}$ and $\mathcal{P}_{\overline{10}}$.

The adjoint representation of $SU(4)_{\text{PS}}$, $\mathcal{P}_{15}$, binds to itself and produces a gauge singlet ϕ_{15-15} with a smaller $C_4^{15-15} = 8$. Additional binding in this channel is due to the $SU(2)_L \times SU(2)_R$ interactions, with $C_L^{15-15} = C_R^{15-15} = 3/2$. However, the SM coupling constants indicate $\alpha_L \lesssim \alpha_4/3$ and $\alpha_R \lesssim \alpha_L/3$ (there is some uncertainty in the running of the three gauge couplings, as discussed in Sec. IV), so it follows that the ϕ_{15-15} di-prebaryon (again a gauge singlet) has a smaller VEV than $\phi_{10-\overline{10}}$, and generates a Majorana mass for $\mathcal{P}_{15}$ that is smaller than the Dirac mass for $\mathcal{P}_{10}$ and $\mathcal{P}_{\overline{10}}$.

The $\mathcal{P}_{6,3}$ binds to itself with $C_4^{6,3-6,3} = 5$ and $C_L^{6,3-6,3} = 4$, and produces a gauge-singlet $\phi_{6,3-6,3}$ di-prebaryon. Likewise $\mathcal{P}_{6,1}$ binds to itself with $C_4^{6,1-6,1} = 5$ and $C_R^{6,1-6,1} = 4$, so the $\phi_{6,1-6,1}$ di-prebaryon has a slightly smaller VEV than $\phi_{6,3-6,3}$. The ensuing Majorana mass for $\mathcal{P}_{6,1}$ is slightly smaller than that for $\mathcal{P}_{6,3}$, which in turn is smaller than the one for $\mathcal{P}_{15}$.

Bound states of the gauge-singlet prebaryons, $\mathcal{P}_{1,1}^i$, may form depending on the strength of the Yukawa couplings of the $\mathcal{A}$ and $\mathcal{A}'$ scalars in Eq. (2.1). For example, if $\lambda_3, \lambda'_{32} > 1$, then $\mathcal{A}$ and $\mathcal{A}'$ exchanges produce a $\mathcal{P}_{1,1}^{32} - \mathcal{P}_{1,1}^{32}$ scalar bound state, which may acquire a VEV. Thus, the $\mathcal{P}_{1,1}^{32}$ prebaryon (a gauge singlet Weyl fermion) gets a large Majorana mass, which may produce a seesaw mechanism responsible for the tiny SM neutrino masses.

B. Di-prebaryons charged under $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$

Besides the di-prebaryons discussed so far, which are all gauge singlets, it is likely that there some di-prebaryons that transform nontrivially under $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$. As the $\mathcal{P}_{10}$, $\mathcal{P}_{\overline{10}}$, $\mathcal{P}_{15}$ (and to a lesser extent $\mathcal{P}_{6,3}$, $\mathcal{P}_{6,1}$) prebaryons are very heavy, nonsinglet bound states involving two of these are expected to have a large squared mass. Bound states involving only one heavy prebaryon and a $\mathcal{P}_4^i$ or $\mathcal{P}_{\overline{4}}^j$ are more likely to acquire a VEV when the binding coefficients are large enough. We will refer to these bound states as heavy-light di-prebaryons. In particular, this is the case of the $\mathcal{P}_{15} - \mathcal{P}_4^i$ bound states, labeled ϕ_{15-4_i} : the tensor product of the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ representations in this case is $(15, 2, 2) \times (4, 2, 1) \rightarrow (4, 1, 2)$, so the binding coefficients are given by $C_4^{15-4} = 4$, $C_L^{15-4} = 3/2$, and $C_R^{15-4} = 0$.

Given that the ϕ_{15-4_i} di-prebaryons transform in the $(4, 1, 2)$ representation, a VEV for one of them would lead

to a phenomenologically correct breaking pattern of the gauge symmetry, as discussed in Sec. III C. Furthermore, the symmetry breaking pattern remains correct when all three ϕ_{15-4_i} acquire VEVs because bilinear terms in the scalar potential of the type $\phi_{15-4_i}^\dagger \phi_{15-4_j}$ align the VEVs in the $SU(4)_{\text{PS}} \times SU(2)_R$ space. These bilinear terms are induced by the antisymmetric scalar $\mathcal{A}'$ running in a loop together with a ψ_k preon, and depend on the product of two different Yukawa couplings shown in (2.1): a $\psi_k \psi_j \mathcal{A}'$ coupling and the Hermitian conjugate of the $\psi_i \psi_k \mathcal{A}'$ coupling.

In the case of the $\mathcal{P}_{15} - \mathcal{P}_4^i$ bound states, labeled ϕ_{15-4_i} , the tensor product is $(15, 2, 2) \times (\bar{4}, 1, 2) \rightarrow (\bar{4}, 2, 1)$ so $C_4^{15-\bar{4}} = 4$, $C_L^{15-\bar{4}} = 0$, $C_R^{15-\bar{4}} = 3/2$, implying weaker binding compared to ϕ_{15-4_i} . Thus, ϕ_{15-4_i} acquire smaller VEVs than ϕ_{15-4_i} , and break the electroweak symmetry because they are doublets under $SU(2)_L$. The remaining heavy-light bound states are $\mathcal{P}_{10} - \mathcal{P}_{\overline{4}}^i$ (labeled $\phi_{10-\bar{4}_i}$) and $\mathcal{P}_{\overline{10}} - \mathcal{P}_4^i$ (labeled $\phi_{\overline{10}-4_i}$), which transform again as $(4, 1, 2)$ and $(\bar{4}, 2, 1)$, respectively. Since $\mathcal{P}_{10}$ is heavier than $\mathcal{P}_{15}$, we expect that the VEVs of $\phi_{10-\bar{4}_i}$ and $\phi_{\overline{10}-4_i}$ are smaller than those of ϕ_{15-4_i} , and could vanish. Even if all of these di-prebaryons acquire VEVs, the symmetry breaking pattern remains correct because these VEVs are aligned with $\langle \phi_{15-4_i} \rangle$ and $\langle \phi_{15-\bar{4}_i} \rangle$ through bilinear terms in the scalar potential generated by tree-level exchange diagrams as the one shown in Fig. 1.

$\mathcal{P}_4^i$ and $\mathcal{P}_{\overline{4}}^j$ may also bind. Although their binding coefficient is smaller, $C_4^{4-\bar{4}} = 15/4$, scalar exchange enhances their binding. We will assume $\lambda_3 \gg \lambda_2, \lambda_1$, such that $\mathcal{A}$ exchange in combination with premeson exchange interactions and $SU(4)_{\text{PS}}$ gauge boson exchange produces a $\mathcal{P}_4^3 - \mathcal{P}_{\overline{4}}^3$ bound state ($\phi_{4_3-\bar{4}_3}$) with nonzero VEV. That bound state transforms as $(1, 2, 2)$ under $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$, implying that under the SM gauge group it has exactly the same quantum numbers as two Higgs doublets. These provide the likely origin for SM quark and lepton masses, as discussed in Sec. III D.

C. Spontaneous breaking of $SU(4)_{\text{PS}} \times SU(2)_R$

As discussed in Sec. III B, there are several scalar bound states transforming in the $(4, 1, 2)$ representation of $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$: ϕ_{15-4_i} and $\phi_{10-\bar{4}_i}$ for $i = 1, 2, 3$. We expect that some of these acquire VEVs, which are smaller than Λ_{pre} and are aligned in the $SU(4)_{\text{PS}} \times SU(2)_R$ space. There is only one linear combination of these heavy-light di-prebaryons that has a nonzero VEV, and we label it here by Φ .

Since the scalar Φ transforms as $(4, 1, 2)$ under $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$, the general renormalizable potential for Φ is

$$V(\Phi) = -M_\Phi^2 \text{Tr} \Phi^\dagger \Phi + \frac{\lambda}{2} (\text{Tr} \Phi^\dagger \Phi)^2 + \frac{\kappa}{2} \text{Tr} (\Phi^\dagger \Phi)^2. \quad (3.5)$$




FIG. 1. Preon exchange diagram inducing the $\phi_{15-4_i}^\dagger \phi_{10-\bar{4}_i}$ bilinear terms in the scalar potential, which are responsible for vacuum alignment along the $SU(4)_{\text{PS}} \times SU(2)_{\text{R}}$ space. Gray bubbles represent the prebaryons (see Table II): $\mathcal{P}_{15}$ and $\mathcal{P}_4^i$ form the ϕ_{15-4_i} di-prebaryons, while $\mathcal{P}_{10}$ and $\mathcal{P}_4^i$ form the $\phi_{10-\bar{4}_i}$ di-prebaryons.

We assume $M_\Phi^2 > 0$ so that the potential has minima away from the origin, and for concreteness we choose $M_\Phi > 0$. The above potential is asymptotically stable for $\kappa > 0$ or $0 \leq -2\kappa < \lambda/2$. To analyze the minima of this potential, we first use an $SU(4)_{\text{PS}} \times SU(2)_{\text{R}}$ transformation to write without loss of generality the VEV of Φ as

$$\langle \Phi \rangle = e^{i\alpha/2} \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & s_2 \\ s_1 & 0 \end{pmatrix}, \quad (3.6)$$

where $s_1, s_2 \geq 0$ have mass dimension 1, and the phase satisfies $-\pi < \alpha \leq \pi$. There are two nontrivial stationarity conditions for the potential:

$$\begin{aligned} \frac{1}{2} \frac{\partial V}{\partial s_1} &= s_1[(\lambda + \kappa)s_1^2 + \lambda s_2^2 - M_\Phi^2] = 0, \\ \frac{1}{2} \frac{\partial V}{\partial s_2} &= s_2[(\lambda + \kappa)s_2^2 + \lambda s_1^2 - M_\Phi^2] = 0. \end{aligned} \quad (3.7)$$

The case where $s_2 = 0$ and $s_1 \neq 0$ gives

$$s_1 = \frac{M_\Phi}{\sqrt{\lambda + \kappa}}, \quad V(\Phi)|_{s_2=0} = -\frac{M_\Phi^4}{2(\lambda + \kappa)}. \quad (3.8)$$

The case where $s_1 = 0$ and $s_2 \neq 0$ is the same up to an $SU(4)_{\text{PS}} \times SU(2)_{\text{R}}$ transformation. Stationarity conditions (3.7) are also satisfied for $s_1, s_2 \neq 0$ provided

$$s_1 = s_2 = \frac{M_\Phi}{\sqrt{\lambda + 2\kappa}}, \quad V(\Phi)|_{s_1=s_2} = -\frac{M_\Phi^4}{\lambda + 2\kappa}. \quad (3.9)$$

The global minimum of $V(\Phi)$ is the one shown in (3.8) if and only if

$$0 < -\lambda < \kappa. \quad (3.10)$$

That vacuum breaks $SU(4)_{\text{PS}} \times SU(2)_{\text{R}}$ down to $SU(3)_c \times U(1)$ where the generator of $U(1)$ is a linear combination

of the diagonal $T_{15} = \text{diag}(1, 1, 1, -3)/\sqrt{24}$ generator of $SU(4)_{\text{PS}}$ and the diagonal generator T_3 of $SU(2)_{\text{R}}$. The gauge couplings of $SU(4)_{\text{PS}}$ and $SU(3)_c$ are the same, g_s . In order to find the gauge coupling of the unbroken $U(1)$, consider the kinetic term for the scalar Φ and focus on the terms that involve the generators mentioned above:

$$\begin{aligned} (D^\mu \Phi^\dagger)(D_\mu \Phi) &\supset |(g_s G^{15\mu} T_{15} + g_R R^{3\mu} T_3) \langle \Phi \rangle|^2 \\ &= \frac{(3g_s^2 + 4g_R^2)M_\Phi^2}{8(\lambda + \kappa)} Z'^\mu Z'_\mu. \end{aligned} \quad (3.11)$$

Here $G^{15\mu}$ and $R^{3\mu}$ are $SU(4)_{\text{PS}}$ and $SU(2)_{\text{R}}$ gauge bosons, respectively, while Z'^μ is their linear combination that becomes massive,

$$Z'^\mu = G^{15\mu} \cos \theta_z - R^{3\mu} \sin \theta_z. \quad (3.12)$$

The mixing angle $\theta_z \in (0, \pi/2)$ satisfies

$$\tan \theta_z = \frac{2g_R}{\sqrt{3}g_s}. \quad (3.13)$$

The linear combination of $G^{15\mu}$ and $R^{3\mu}$ that remains massless at this stage,

$$B^\mu = G^{15\mu} \sin \theta_z + R^{3\mu} \cos \theta_z, \quad (3.14)$$

is the SM hypercharge gauge boson. To see that, note that the interactions of this gauge boson are determined by the couplings of the $G^{15\mu}$ and $R^{3\mu}$ gauge bosons to the quarks and leptons contained in the $\mathcal{P}_4^i$ and $\mathcal{P}_{\bar{4}}^i$ prebaryons (see Tables II and III), which belong to the $(4, 1)$ and $(\bar{4}, 2)$ representations of $SU(4)_{\text{PS}} \times SU(2)_{\text{R}}$:

$$\begin{aligned} &\frac{g_s}{2\sqrt{6}} G^{15\mu} (\bar{q}_L \gamma^\mu q_L - 3\bar{\ell}_L \gamma^\mu \ell_L) \\ &= g_Y B^\mu (Y_{q_L} \bar{q}_L \gamma^\mu q_L + Y_{\ell_L} \bar{\ell}_L \gamma^\mu \ell_L) + g_z Z'^\mu J_{L\mu}^{\prime}, \end{aligned} \quad (3.15)$$

TABLE III. Prebaryons of $SU(15)_p$ lighter than Λ_{PS} and their representations under the SM gauge group. The SM quarks and leptons arise as $\Omega\Psi_W\psi_i$ and $\Omega\Psi_{W'}\psi_i$ prebaryons, with the generation index $i = 1, 2, 3$ identified as the preon flavor index. Fermions beyond the SM (below Λ_{PS}) include five SM singlets and one vectorlike lepton doublet. The last column includes the original prebaryons, before $SU(4)_{PS} \times SU(2)_R$ breaking (see Table II).

Prebaryons	Preon content	$SU(3)_c \times SU(2)_L \times U(1)_Y$	Origin
q_{iL}, ℓ_{iL}	$\Omega\Psi_W\psi_i$	$(3, 2, +1/6), (1, 2, -1/2)$	$\mathcal{P}_4^i$
$u_i^c, d_i^c, e_i^c, N_i^c$	$\Omega\Psi_{W'}\psi_i$	$(\bar{3}, 1, -2/3), (\bar{3}, 1, +1/3), (1, 1, +1), (1, 1, 0)$	$\mathcal{P}_4^i$
N_{31}, N_{21}	$\Omega\psi_3\psi_1, \Omega\psi_2\psi_1$	$(1, 1, 0)$	$\mathcal{P}_{1,1}^{31}, \mathcal{P}_{1,1}^{21}$
$\mathcal{L}^c, \mathcal{L}_L$	$\Omega\Psi_W\Psi_{W'}$	$(1, 2, +1/2) + (1, 2, -1/2)$	$\mathcal{P}_{1,2}$

$$\begin{aligned}
& \frac{g_s}{2\sqrt{6}} G^{15\mu} (\bar{q}_R \gamma^\mu q_R - 3\bar{\ell}_R \gamma^\mu \ell_R) \\
& + \frac{g_R}{\sqrt{2}} R^{3\mu} (\bar{u}_R \gamma^\mu u_R - \bar{d}_R \gamma^\mu d_R + \bar{N}_R \gamma^\mu N_R - \bar{e}_R \gamma^\mu e_R) \\
& = g_Y B^\mu (Y_{u_R} \bar{u}_R \gamma^\mu u_R + Y_{d_R} \bar{d}_R \gamma^\mu d_R + Y_{e_R} \bar{e}_R \gamma^\mu e_R) \\
& + g_z Z'^\mu J'_{R\mu}. \tag{3.16}
\end{aligned}$$

Here we are using the right-handed quark and lepton field, which are the conjugates of the $\mathcal{P}_4^i$ prebaryons (all prebaryons in Table II are left-handed fermion fields).

From (3.15) follows that the relation between the gauge couplings for B^μ and for gluons can be written as

$$g_Y = \sqrt{\frac{3}{2}} g_s \sin \theta_z. \tag{3.17}$$

Equations (3.15) and (3.16) are satisfied provided

$$Y_{q_L} = \frac{1}{6}, \quad Y_{\ell_L} = -\frac{1}{2}, \quad Y_{u_R} = \frac{2}{3}, \quad Y_{d_R} = -\frac{1}{3}, \quad Y_{e_R} = -1, \tag{3.18}$$

which confirms that B^μ is the SM $U(1)_Y$ gauge boson. The $J'_{L\mu}$ and $J'_{R\mu}$ represent the left-handed and right-handed quark and lepton currents coupled to Z'^μ , with appropriate charges, while g_z is the gauge coupling of the heavy gauge boson.

D. Effective theory below the $SU(4)_{PS} \times SU(2)_R$ breaking scale

At scales below Λ_{PS} , where the $SU(4)_{PS} \times SU(2)_L \times SU(2)_R$ gauge symmetry is spontaneously broken down to $SU(3)_c \times SU(2)_L \times U(1)_Y$, the effective theory includes the composite fermions that do not form di-prebaryons with large VEVs. These include the $\mathcal{P}_4^i$, $\mathcal{P}_4^i$, $\mathcal{P}_{1,2}$, and $\mathcal{P}_{1,1}^{i1}$ prebaryons, described in Table II. Their SM gauge charges are shown in Table III.

1. SM fermions and composite right-handed neutrinos

The prebaryons with preon content $\Omega\Psi_W\psi_i$ represent the left-handed doublets of the SM: the q_{iL} quarks and ℓ_{iL}

leptons. The index $i = 1, 2, 3$ labels the three generations, arising from the flavor index of the preons ψ_i . The prebaryons with preon content $\Omega\Psi_{W'}\psi_i$ represent the charge conjugates of the right-handed SM fermions: up-type quarks u_i^c , down-type quarks d_i^c , and charged leptons e_i^c . In addition, the three conjugates of right-handed neutrinos, N_i^c , also arise as $\Omega\Psi_{W'}\psi_i$ bound states; they are gauge singlets under the SM gauge group, but couple to the heavy gauge bosons associated with the breaking of $SU(4)_{PS} \times SU(2)_R$.

The $\mathcal{P}_{1,1}^{ij}$ prebaryons, whose preon content is $\Omega\psi_i\psi_j$ (with $i \neq j$), are neutral not only under the SM gauge interactions but also under all $SU(4)_{PS} \times SU(2)_L \times SU(2)_R$ gauge symmetries. Following the discussion at the end of Sec. III B, we assume that $\mathcal{P}_{1,1}^{32}$ has a Majorana mass above Λ_{PS} , due to $\mathcal{A}$ exchanges; this can be responsible for a high-mass seesaw mechanism involving the SM neutrinos. The other two $\mathcal{P}_{1,1}^{ij}$ prebaryons, which for brevity we label here N_{12} and N_{13} , have a Dirac mass below Λ_{PS} ; nevertheless, they likely have some mixings with the N_i^c fermions as well as with the SM neutrinos, and may mediate additional seesaw contributions to the neutrino masses.

Hence, the three N_i^c fermions play the role of sterile neutrinos. We will not explore here the phenomenological implications of this rich composite neutrino sector. It is also possible that one or two of the N_i^c fermions are stable, in which case they would constitute a dark matter component. Other dark matter candidates could in principle be provided by physics above Λ_{pre} .

2. Composite vectorlike leptons

The lepton doublets $\mathcal{L}_L$ and $\mathcal{L}^c$ arise from the prebaryon $\mathcal{P}_{1,2} = \Omega\Psi_W\Psi_{W'}$. These doublets have effective Yukawa couplings to the di-prebaryons $\phi_{10-\overline{10}}$ and ϕ_{15-15} , which are generated by the diagrams shown in Figs. 2 and 3. The VEVs of the di-prebaryons thus lead to a Dirac mass $m_{\mathcal{L}}$ for $\mathcal{L}_L$ and $\mathcal{L}^c$, which form a vectorlike lepton. As argued in [6], the two contributions to $m_{\mathcal{L}}$ are accidentally of the same order of magnitude. The first contribution, the preon-interchange diagram in Fig. 2, is suppressed by $1/N$ where $N = 15$ for the gauge group responsible for preon confinement. The second contribution, the $SU(4)_{PS}$ -exchange




FIG. 2. Preon exchange diagram inducing the effective Yukawa interaction of the $\phi_{10-\bar{10}}$ scalar (a gauge-singlet bound state of the $\mathcal{P}_{10}$ and $\mathcal{P}_{\bar{10}}$ prebaryons) to a pair of $\mathcal{P}_{1,2}$ prebaryons.

diagram in Fig. 3, is suppressed by α_4/π . As a result, $m_{\mathcal{L}}$ may be at the TeV scale, and thus the existence of the vectorlike lepton doublet may be probed at the LHC and future colliders.

If the vectorlike lepton doublet $\mathcal{L} = (\mathcal{L}^0, \mathcal{L}^-)$ mixes with the SM leptons, then the standard decay modes are $\mathcal{L}^\pm \rightarrow \nu W^\pm, \tau^\pm Z, \tau^\pm h^0$ (ν generically stands here for ν_τ or $\bar{\nu}_\tau$) and $\mathcal{L}^0 \rightarrow \tau^- W^+, \nu Z, \nu h^0$; note that the antiparticle of $\mathcal{L}^0$ then has the decay modes $\bar{\mathcal{L}}^0 \rightarrow \tau^+ W^-, \nu Z, \nu h^0$. In that case, the current LHC limit set by the CMS [24] and ATLAS [25] collaborations is $m_{\mathcal{L}} > 1.0$ TeV. The high-luminosity LHC will be sensitive to heavier vectorlike leptons. At a $\mu^+\mu^-$ collider with center-of mass energy $\sqrt{s} > 2m_{\mathcal{L}}$, Drell-Yan production of a pair of vectorlike leptons would allow precision measurements of their properties.

In this theory of quark and lepton compositeness, however, the vectorlike lepton doublet may also have more exotic decay modes, especially if the pseudo-Nambu-Goldstones arising from neutral di-prebaryons have masses below $m_{\mathcal{L}}$. If $A_{\mathcal{L}}$ is one of those pseudoscalars, then $\mathcal{L}^\pm \rightarrow \tau^\pm A_{\mathcal{L}}$ and $\mathcal{L}^0 \rightarrow \nu A_{\mathcal{L}}$ may be the dominant decay modes. The production of vectorlike leptons at the LHC with the




FIG. 3. Effective Yukawa interaction of the ϕ_{15-15} di-prebaryon (a gauge-singlet bound state of two $\mathcal{P}_{15}$ prebaryons) to a pair of $\mathcal{P}_{1,2}$ prebaryons, induced by an $SU(4)_{\text{PS}}$ gauge boson exchange.

largest cross section is via an off-shell W boson, so that the main processes are $pp \rightarrow W^{*+} \rightarrow \mathcal{L}^+ \mathcal{L}^0 \rightarrow (\tau^\pm A_{\mathcal{L}})(\nu_\tau A_{\mathcal{L}})$ and $pp \rightarrow W^{*-} \rightarrow \mathcal{L}^- \bar{\mathcal{L}}^0 \rightarrow (\tau^- A_{\mathcal{L}})(\bar{\nu}_\tau A_{\mathcal{L}})$. The process with a W^{*+} has the largest cross section (about 1 fb for $m_{\mathcal{L}} = 1$ TeV at $\sqrt{s} = 14$ TeV [26]), because the parton distribution is larger for the up quark than for the down quark.

The subsequent decays of $A_{\mathcal{L}}$ depend on its mass, and on possible couplings to the b or t quarks. For a $A_{\mathcal{L}}$ mass in the 3.5–10 GeV, the dominant decay is $A_{\mathcal{L}} \rightarrow \tau^+ \tau^-$, which leads to the $5\tau + \nu$ final state shown in Fig. 4. Searches in this final state have not yet been performed, and should be pursued by the ATLAS and CMS collaborations, as well as at future collider experiments. For $m_{\mathcal{L}} < 3.5$ GeV, the one-loop decay $A_{\mathcal{L}} \rightarrow \gamma\gamma$ competes with the $A_{\mathcal{L}} \rightarrow \mu^+ \mu^-$ decay, which is suppressed by the VEV of the neutral di-prebaryon; thus, the relevant final states are of the type $(\tau^\pm A_{\mathcal{L}})(\nu A_{\mathcal{L}}) \rightarrow \tau^\pm \nu + 4\gamma$ or $\tau^\pm \nu + 4\mu$, with the possibility of displaced vertices due to the long lifetime of $A_{\mathcal{L}}$.

It is interesting that the above phenomenological signatures are different from those predicted in the theory of quark and lepton compositeness [2,6] where the weakly coupled gauge group at the Λ_{pre} scale is the SM gauge group, instead of $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ as analyzed in this paper. The lightest composite vectorlike lepton in the theory of Refs. [2,6] is a weak singlet, which is produced at colliders via an off-shell Z and γ , and leads to an even number of τ 's in the final state [27]. Furthermore, in that case, there are some vectorlike quarks of charge $-1/3$ with mass potentially within the reach of the LHC.

3. Composite Higgs sector

As discussed in Sec. III B, the three $\phi_{15-\bar{4}_i}$ di-prebaryons transform as $(\bar{4}, 2, 1)$ and acquire VEVs smaller than the $SU(4)_{\text{PS}} \times SU(2)_R$ breaking VEVs of ϕ_{15-4_i} . Given that all $SU(4)_{\text{PS}}$ breaking VEVs are aligned, below Λ_{PS} each of the $\phi_{15-\bar{4}_i}$ consists of a scalar $\tilde{q}_i^c$ transforming as $(\bar{3}, 2, -1/6)$ under the SM gauge group plus a scalar H_i transforming as $(1, 2, +1/2)$. Consequently, there are three Higgs doublets,




FIG. 4. Main process for pair production of composite vectorlike leptons at the LHC, yielding a $5\tau + E_{\text{f}}$ final state, provided that the mass of the composite pseudoscalar $A_{\mathcal{L}}$ satisfies $2m_\tau < M_A < m_{\mathcal{L}}$. For $M_A < 2m_\tau$, $A_{\mathcal{L}}$ predominantly decays into $\gamma\gamma$ or $\mu^+ \mu^-$, so similar diagrams lead to $4\gamma + \tau + E_{\text{f}}$ or $4\mu + \tau + E_{\text{f}}$ final states.

H_i for $i = 1, 2, 3$, which have approximately equal VEVs (with small corrections due to $\mathcal{A}$ and $\mathcal{A}'$ loops).

Another source of electroweak symmetry breaking is the $\phi_{4_3-\bar{4}_3}$ di-prebaryon discussed at the end of Sec. III B, which includes two Higgs doublets, H_u and H_d . These generate the SM fermion masses through a mechanism similar to the one presented in [2]. The $\mathcal{P}_4^3$ and $\mathcal{P}_4^{\bar{3}}$ prebaryons, which are identified with third-generation SM fermions as displayed in Table III, form the two composite Higgs doublets. Consequently, it is expected that the Yukawa couplings of third-generation SM fermions are larger than one. If the $\phi_{4_3-\bar{4}_3}$ VEV is predominantly tilted in the direction of H_u , then the top quark gets a mass near the electroweak scale. Note that the Yukawa coupling larger than one is compensated by the H_u VEV that is below $v \approx 246$ GeV (because the H_i VEVs also contribute to electroweak symmetry breaking). The b quark and τ lepton, which are also part of the $\mathcal{P}_4^3$ and $\mathcal{P}_4^{\bar{3}}$ prebaryons, have Yukawa couplings to H_d that are comparable to the top coupling to H_u , so this is a type-II two-Higgs-doublet model in the large $\tan\beta$ regime, supplanted by the three fermiophobic Higgs doublets H_i .

The SM fermions of the second and first generations have effective Yukawa couplings to H_u and H_d which are mediated by $\mathcal{A}$ and $\mathcal{A}'$ scalar exchange (the relevant diagram for the particular case of the charm quark is shown in Fig. 5). As a result, their masses are proportional to a sum of products of the Yukawa couplings of $\mathcal{A}$ and $\mathcal{A}'$ shown in Eq. (2.1). For example, the up-type quark mass matrix $\mathcal{M}_u$ has the 22 element proportional to $\lambda_{33}\lambda_{22} + (\lambda'_{33})^\dagger \lambda'_{22}$. Its off-diagonal elements get contributions only from $\mathcal{A}'$ exchange in the basis where $\mathcal{A}$ has only diagonal couplings; e.g., the 23 element of $\mathcal{M}_u$ is proportional to $(\lambda'_{33})^\dagger \lambda'_{23}$.

Thus, the SM quarks and leptons acquire masses, with the third generation of fermions naturally heavier than the

other two, which are suppressed by a scalar exchange. The same would be true even if only one scalar in the antisymmetric representation of $SU(15)_p$ existed. However, in the absence of $\mathcal{A}'$ (as was the case in [2]), there would not be off-diagonal Cabibbo–Kobayashi–Maskawa matrix elements unless there is an additional source of preon flavor violation (such as some higher-dimensional operators). This justifies the introduction of $\mathcal{A}'$ in Sec. II.

Given that the SM quarks and leptons obtain masses from an effective two-Higgs-doublet model of type II, there are no tree-level flavor-changing neutral processes. There may be, however, flavor-violating higher-dimensional operators induced at the compositeness scale. We expect that their effects are adequately suppressed (given that proton stability implies $\Lambda_{\text{pre}} \gtrsim \times 10^3$ TeV), but dedicated studies should analyze whether flavor processes induced by effects at the compositeness scale may eventually be observed.

IV. RUNNING OF GAUGE COUPLINGS

We now explore the behavior of the gauge couplings at energy scales above the electroweak scale and up to the compositeness scale Λ_{pre} , using renormalization group equations (RGEs) [28–30]. The evolution of a coupling g_i takes the form

$$\frac{dg_i}{d\ln\mu} = \beta_i(\{g_j\}), \quad (4.1)$$

where μ is the renormalization scale, and the beta function β_i of coupling g_i depends on the set of couplings denoted by $\{g_j\}$, which includes g_i . The beta functions are calculated perturbatively from the relationship between the bare and renormalized couplings. For a generic field theory with gauge couplings g_i and Yukawa couplings Y_{jk}^i (from terms of the form $Y_{jk}^i \phi_i \bar{\psi}_j \psi_k$ in the Lagrangian), the beta functions can be written as $\beta_i = -b_i g_i^3 / (4\pi)^2$ with coefficients given by

$$b_i = b_i^{(1)} + \frac{1}{4\pi} \left(b_i^{(2)} \alpha_i + \sum_{j \neq i} b_{ij}^{(2)} \alpha_j + \sum_j \mathcal{Y}_j \right). \quad (4.2)$$

Here $\alpha_i = g_i^2 / (4\pi)$, and the one- and two-loop contributions are given by

$$\begin{aligned} b_i^{(1)} &= \frac{11}{3} C(G_i) - \frac{2}{3} T_i(F) - \frac{1}{6} T_i(S), \\ b_i^{(2)} &= \frac{34}{3} C(G_i)^2 - C(G_i) \left(\frac{10}{3} T_i(F) + \frac{1}{3} T_i(S) \right), \\ b_{ij}^{(2)} &= -2 \left(C_i(F) T_j(F) + C_i(S) T_j(S) \right), \\ \mathcal{Y}_i(F) &= \frac{1}{d(G_i)} \text{Tr}[C_i(F) Y^i (Y^i)^\dagger]. \end{aligned} \quad (4.3)$$




FIG. 5. Effective Yukawa coupling induced by preon interactions: the H_u Higgs doublet (a six-preon bound state) couples to the second-generation SM quark doublet (q_{2L}) and up-type quark singlet (u_2^c), producing the charm quark mass. The antisymmetric scalars $\mathcal{A}$ and $\mathcal{A}'$ couple to different preon flavors as in Eq. (2.1). The preon ψ_Q is the color-triplet component of Ψ_W , while the preon ψ_U is the color-antitriplet and upper- $SU(2)_R$ component of $\Psi_{W'}$.

The quadratic Casimir invariants appearing here, C , correspond to those of Weyl fermions (F), real scalars (S), and gauge bosons (G). The second Dynkin index, T , refers to Weyl fermions and real scalars, and the dimension of the Lie algebra of G is given by $d(G)$.

The RGEs will be for the SM gauge couplings up to the scale Λ_{PS} where there is a transition to the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ couplings. The SM coupling constants associated with the $SU(3)_c \times SU(2)_W \times U(1)_Y$ gauge interactions are α_s ,

$$\alpha_2 = \frac{\alpha}{\sin^2 \theta_W}, \quad \alpha_Y = \frac{\alpha}{\cos^2 \theta_W}, \quad (4.4)$$

where α is the electromagnetic coupling and θ_W is the weak mixing angle. The transition between the two sets of coupling constants at the Λ_{PS} scale satisfies the following tree-level matching conditions [31]:

$$\begin{aligned} \alpha_Y^{-1}(\Lambda_{\text{PS}}) &= \alpha_R^{-1}(\Lambda_{\text{PS}}) + \frac{2}{3} \alpha_4^{-1}(\Lambda_{\text{PS}}), \\ \alpha_2(\Lambda_{\text{PS}}) &= \alpha_L(\Lambda_{\text{PS}}), \\ \alpha_s(\Lambda_{\text{PS}}) &= \alpha_4(\Lambda_{\text{PS}}). \end{aligned} \quad (4.5)$$

One can also evolve the Yukawa couplings as they begin to play a role at the two-loop level in the running of the gauge couplings, as can be seen in Eq. (4.3).

In our analysis of gauge-Yukawa theories, we have utilized two-loop order beta functions from Refs. [32–35], evaluated using the packages SARAH 4 [36] and Rgbeta [37]. We set initial conditions for the SM gauge couplings at the top quark mass using MR:C++ [38], as shown in Ref. [39]. At scales above m_t , the running required recalculating beta functions at each mass threshold described in Sec. III.

A. Beta functions between mass thresholds

The lightest new particles charged under the SM gauge group are the vectorlike lepton doublet (of mass m_L), and the scalars associated with the second Higgs doublet (a linear combination of the states in H_u and H_d), and the three $\phi_{15-\bar{4}_i}$ scalars (which include three $\tilde{q}_i^c$ and three H_i), as discussed in Sec. III D. We assume for simplicity that all of these new particles are approximately degenerate in mass, so that the gauge coupling running up to the m_L scale is governed just by the SM. Above m_L , the coefficients of the beta functions for the three SM gauge couplings are

$$\begin{aligned} b_Y &= -\frac{25}{3} - \frac{1}{4\pi} \left(\frac{52}{3} \alpha_s + \frac{27}{2} \alpha_2 + \frac{245}{18} \alpha_Y \right), \\ b_2 &= \frac{1}{3} - \frac{1}{\pi} \left(9\alpha_s + \frac{253}{24} \alpha_2 + \frac{9}{8} \alpha_Y \right), \\ b_s &= 6 + \frac{1}{4\pi} \left(4\alpha_s - \frac{27}{2} \alpha_2 - \frac{13}{6} \alpha_Y \right). \end{aligned} \quad (4.6)$$

Here we have included the one-loop contributions, as well as the two-loop gauge contributions. At two loops there are also contributions from the Yukawa couplings, especially those of the third generation fermions to H_u and H_d , which for simplicity we do not include here.

The next threshold occurs around Λ_{PS} , where the SM gauge couplings transition to $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ couplings according to Eq. (4.5). The following left-handed fermion representations are present here: $3 \times (4, 2, 1)$, $3 \times (\bar{4}, 1, 2)$, $(1, 2, 2)$, corresponding to the prebaryons $\mathcal{P}_4^i$, $\mathcal{P}_{\bar{4}}^i$, $\mathcal{P}_{1,2}$, respectively (see Table II). The scalar representations discussed in Sec. III B are exactly the conjugates of the above fermion ones: $3 \times (\bar{4}, 2, 1)$, $3 \times (4, 1, 2)$, $(1, 2, 2)$, due to the di-prebaryons $\phi_{15-\bar{4}_i}$, ϕ_{15-4_i} , $\phi_{4_3-\bar{4}_3}$, respectively. Interestingly, although the underlying theory is not supersymmetric, the pair of bound states $\{\mathcal{P}_4^i, \phi_{15-\bar{4}_i}^\dagger\}$ form a chiral supermultiplet for each $i = 1, 2, 3$, and the same is true for $\{\mathcal{P}_{\bar{4}}^i, \phi_{15-4_i}^\dagger\}$ and $\{\mathcal{P}_{1,2}, \phi_{4_3-\bar{4}_3}\}$. The pure gauge β function coefficients at one and two loops, for scales between Λ_{PS} and $m_{\mathcal{P}_6}$ (the mass of the antisymmetric prebaryons) are given by

$$\begin{aligned} b_R &= \frac{1}{3} - \frac{1}{8\pi} \left(135\alpha_4 + \frac{253}{3} \alpha_L + 9\alpha_R \right), \\ b_L &= \frac{1}{3} - \frac{1}{8\pi} \left(135\alpha_4 + 9\alpha_L + \frac{253}{3} \alpha_R \right), \\ b_4 &= \frac{26}{3} + \frac{1}{8\pi} \left(\frac{107}{3} \alpha_4 - 27\alpha_L - 27\alpha_R \right). \end{aligned} \quad (4.7)$$

At scales above $m_{\mathcal{P}_6}$ we include intermediate-mass states transforming under the representations $(6, 1, 3)$ and $(6, 3, 1)$, corresponding to the $\mathcal{P}_{6,1}$ and $\mathcal{P}_{6,3}$ prebaryons, respectively. The pure gauge coupling beta function coefficients become

$$\begin{aligned} b_R &= -\frac{23}{3} - \frac{1}{8\pi} \left(255\alpha_4 + 9\alpha_L + \frac{1021}{3} \alpha_R \right), \\ b_L &= -\frac{23}{3} - \frac{1}{8\pi} \left(255\alpha_4 + \frac{1021}{3} \alpha_L + 9\alpha_R \right), \\ b_4 &= \frac{14}{3} - \frac{1}{8\pi} \left(\frac{553}{3} \alpha_4 + 51\alpha_L + 51\alpha_R \right). \end{aligned} \quad (4.8)$$

At a scale of order $m_{\mathcal{P}_6}$ two di-prebaryons also become relevant: $\phi_{6,1-6,1}$ and $\phi_{6,3-6,3}$. These are gauge-singlet scalars with large Yukawa couplings to the $\mathcal{P}_{6,1}$ and $\mathcal{P}_{6,3}$ prebaryons. The latter belong to a higher representation of $SU(4)_{\text{PS}}$ (the antisymmetric one), so their Yukawa couplings have larger two-loop effects on the gauge running compared to those of the top and bottom quarks. We incorporate the effects of $\phi_{6,1-6,1}$ and $\phi_{6,3-6,3}$ in the numerical study presented in Sec. IV B but their effects on the gauge running will turn out to be only barely visible.

Next, we encounter the threshold corresponding to the heaviest prebaryons, which transform under the representations $(15, 2, 2)$, $(10, 1, 1)$, and $(\bar{10}, 1, 1)$, as shown in Table II. These prebaryons significantly impact the beta functions due to their large $SU(4)_{\text{PS}}$ representations. The pure gauge beta functions at scales above $m_{\mathcal{P}_{10}}$ have the following coefficients:

$$\begin{aligned} b_R &= -\frac{46}{3} - \frac{1}{4\pi} \left(\frac{405}{2} \alpha_4 + 24\alpha_L + \frac{787}{3} \alpha_R \right), \\ b_L &= -\frac{46}{3} - \frac{1}{4\pi} \left(\frac{405}{2} \alpha_4 + \frac{787}{3} \alpha_L + 24\alpha_R \right), \\ b_4 &= -8 - \frac{1}{8\pi} (1013\alpha_4 + 81\alpha_L + 81\alpha_R). \end{aligned} \quad (4.9)$$

These prebaryons also form two gauge-singlet scalars, ϕ_{15-15} and $\phi_{10-\bar{10}}$. For the purpose of calculating the running of the gauge couplings, we assume that the masses of these prebaryons and their associated scalars are degenerate at the scale $m_{\mathcal{P}_{10}}$. Their Yukawa couplings again contribute to the running of the gauge couplings at the two-loop order, but the effects are larger due to the higher $SU(4)_{\text{PS}}$ representations. These are numerically included in Sec. IV B, with the starting value for the Yukawa couplings at the scale $m_{\mathcal{P}_{10}}$ being close to the nonperturbative regime, $y_{10-\bar{10}} \approx y_{15-15} \approx 2$.

At this stage, the presence of many composite fields has lead to the loss of asymptotic freedom for each of the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge interaction. Nevertheless, we will see next that these interactions remain perturbative at least up to the compositeness scale.

B. Gauge coupling evolution

We perform the running with $\overline{\text{MS}}$ renormalized beta functions. At the mass of the top quark, $m_t \approx 173$ GeV [40], the SM gauge coupling constants are given by [39]

$$\begin{aligned} \alpha_Y(m_t) &= 0.01023, & \alpha_2(m_t) &= 0.03337, \\ \alpha_s(m_t) &= 0.01074. \end{aligned} \quad (4.10)$$

The mass thresholds, discussed in Sec. IV A, satisfy the following hierarchy:

$$m_t < m_{\mathcal{L}} < \Lambda_{\text{PS}} < m_{\mathcal{P}_6} < m_{\mathcal{P}_{10}} < \Lambda_{\text{pre}}. \quad (4.11)$$

Using a reasonable set of values for these mass thresholds, we show the evolution of the inverse coupling constants in Fig. 6. There, we set the mass of the composite vectorlike lepton at $m_{\mathcal{L}} = 2$ TeV (the current lower limit is 1 TeV, see Sec. III D). For the scale of $SU(4)_{\text{PS}} \times SU(2)_R$ breaking we take $\Lambda_{\text{PS}} = 30$ TeV. Note that in perturbative, minimal Pati-Salam models, constraints from rare meson decays impose a lower limit on the symmetry breaking




FIG. 6. Two-loop running of the inverse coupling constants (α^{-1}) for $SU(3)_c \times SU(2)_L \times U(1)_Y$ from the m_t scale up to Λ_{PS} , and for $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ between Λ_{PS} and Λ_{pre} . Vertical lines mark various thresholds. Prebaryons $\mathcal{P}_{10}$, $\mathcal{P}_{\bar{10}}$, and $\mathcal{P}_{15}$, assumed to have same mass $m_{\mathcal{P}_{10}}$, reduce the three α^{-1} near Λ_{pre} . Prebaryons $\mathcal{P}_{6,3}$ and $\mathcal{P}_{6,1}$ (of mass $m_{\mathcal{P}_6}$), as well as the charged di-prebaryons and the composite vectorlike lepton (of mass $m_{\mathcal{L}}$), have a similar but less notable impact. The two-loop effects of the composite singlet scalars are shown in each lower line fork near Λ_{pre} . The gray-shaded vertical band at Λ_{pre} represents the region of strongly coupled preonic $SU(15)_p$ dynamics. The yellow-shaded horizontal band indicates the region where the three α 's would no longer be perturbative.

scale of 80 TeV [41,42]. However, in the presence of vectorlike fermions, that limit can be lowered to about 5 TeV [41,43].

The remaining composite vectorlike fermions, which are heavier as they belong to higher representations of $SU(4)_{\text{PS}}$ (see Table II), are assumed to have masses at $m_{\mathcal{P}_6} = 100$ TeV (for the $\mathcal{P}_{6,3}$ and $\mathcal{P}_{6,1}$ prebaryons) and $m_{\mathcal{P}_{10}} = 300$ TeV (for the $\mathcal{P}_{15}$, $\mathcal{P}_{10}$, and $\mathcal{P}_{\overline{10}}$ prebaryons). The benchmark choice for confinement scale $\Lambda_{\text{pre}} = 3 \times 10^3$ TeV is in line with the bounds from proton decay estimated in Ref. [6].

As can be seen in Fig. 6, above $m_{\mathcal{L}}$ the three gauge couplings run to larger values compared to the SM values at the same scale. The strong coupling starts decreasing faster above Λ_{PS} due to the additional non-Abelian gauge bosons. The hypercharge coupling is replaced by the $SU(2)_R$ coupling at Λ_{PS} , according to the matching condition (4.5). As expected based on Eqs. (4.7) and (4.8), the $SU(2)_L \times SU(2)_R$ couplings remain almost constant at scales in the $\Lambda_{\text{PS}} - m_{\mathcal{P}_6}$ range, while above the $m_{\mathcal{P}_6}$ scale they lose asymptotic freedom. Above the scale of the heaviest chiral prebaryons, $m_{\mathcal{P}_{10}}$, the $SU(4)_{\text{PS}}$ interaction also loses asymptotic freedom, but easily remains perturbative up to the compositeness scale Λ_{pre} . The two-loop effects of the gauge-singlet scalars are shown in Fig. 6 for each coupling as the lower branch near Λ_{pre} , with the starting value (at the scalar masses) for all Yukawa couplings taken to be $y \approx 2$. For comparison, the upper branches show the results without Yukawa couplings.

C. Compositeness scale and premeson effects

Understanding the behavior of the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge couplings near the compositeness scale Λ_{pre} is challenging due to the presence of numerous charged fields and the nonperturbative effects of the $SU(15)_p$ interactions. Nevertheless, at least a rough estimate of the shifts in the gauge couplings near Λ_{pre} is necessary for assessing the viability of the theory analyzed here. To see that, recall that we already encountered (below Λ_{pre}) a loss of asymptotic freedom in the gauge couplings of $SU(2)_L \times SU(2)_R$ above the scale $m_{\mathcal{P}_6}$, and of $SU(4)_{\text{PS}}$ above the scale $m_{\mathcal{P}_{10}}$, due to the presence of several prebaryons charged under these gauge groups. Moreover, above Λ_{pre} ,

the theory includes multiple preons that contribute to the running of the gauge couplings, preventing asymptotic freedom (see Sec. II B) and making it difficult for the couplings to remain perturbative towards higher energy scales.

To investigate this, we consider the impact on the coupling running of the premesons of spin-1 that are charged under $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$. These premesons have masses around the compositeness scale Λ_{pre} , and provide an uplift of the three inverse coupling constants. The specific premesons formed from the preon fields Ψ_W , Ψ'_W , and ψ_i are constructed as bilinears of the form $\bar{\Psi}_W \sigma^\mu \Psi_{W'}$, $\bar{\Psi}_W \sigma^\mu \Psi_W$, $\bar{\psi}_i \sigma^\mu \Psi_W$, etc. They are labeled by $\rho_{WW'}$, ρ_{WW} , ρ_{iW} , etc., and transform under the gauge representations listed in Table IV. Because of their large quadratic Casimir invariants or large multiplicities, these spin-1 fields substantially decrease the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge couplings.

The one-loop contributions from all spin-1 premesons on the coefficient of the beta function for an $SU(N)$ gauge group ($N = 4$ or 2 in our case) are

$$b(\rho) = \frac{11}{3} \left[N_{\text{Ad}} C_2(\text{Ad}) + N_S C_2(S) + N_A C_2(A) + N_F C_2(F) \right]. \quad (4.12)$$

Here N_{Ad} , N_S , N_A , and N_F are the numbers of spin-1 premesons transforming in the adjoint, symmetric, anti-symmetric, and fundamental representations, as outlined in Table IV. Note that $b(\rho) > 0$ because the contribution of spin-1 fields in non-Abelian gauge theories is towards asymptotic freedom. In the specific case of $SU(4)_{\text{PS}}$, the quadratic Casimir invariants for these representations are $C_2(\text{Ad}) = 4$, $C_2(S) = 9/2$, $C_2(A) = 5/2$, and $C_2(F) = 15/8$. Using the number of pre-mesons in each representation (specifically, $N_{\text{Ad}} = N_A = N_S = 8$, $N_F = 24$), the beta function coefficient for the $SU(4)_{\text{PS}}$ gauge coupling is found to be very large, $b_4(\rho) = 1463/3$. Similarly, $b_{L,R}(\rho) = 1078/3$, which is almost 2 orders of magnitude larger than typical one-loop beta function coefficients in the SM.

Besides spin-1 premesons, there are also many spin-0 premesons and vectorlike prebaryons with masses around

TABLE IV. Spin-1 premesons (of mass around Λ_{pre}) transforming under the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge group. The generation index is $i = 1, 2, 3$.

Spin-1 premesons	Preon content	$SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ representations
$\rho_{WW'}$, $\rho_{W'W}$	$\bar{\Psi}_W \sigma^\mu \Psi_{W'}$, $\bar{\Psi}_{W'} \sigma^\mu \Psi_W$	$(6, 2, 2) + (\overline{10}, 2, 2)$, conjugates
ρ_{WW}	$\bar{\Psi}_W \sigma^\mu \Psi_W$	$(15, 3, 1) + (15, 1, 1) + (1, 3, 1)$
$\rho_{W'W'}$	$\bar{\Psi}_{W'} \sigma^\mu \Psi_{W'}$	$(15, 1, 3) + (15, 1, 1) + (1, 1, 3)$
ρ_{iW} , ρ_{Wi}	$\bar{\psi}_i \sigma^\mu \Psi_W$, $\bar{\Psi}_W \sigma^\mu \psi_i$	$3 \times (4, 2, 1)$, conjugates
$\rho_{iW'}$, $\rho_{W'i}$	$\bar{\psi}_i \sigma^\mu \Psi_{W'}$, $\bar{\Psi}_{W'} \sigma^\mu \psi_i$	$3 \times (\bar{4}, 1, 2)$, conjugates

the Λ_{pre} scale. Their effect is harder to estimate, but it is in the opposite direction compared to the spin-1 states. Thus, the large positive contributions $b_4(\rho)$ and $b_{L,R}(\rho)$ are at least partially compensated by other prehadrons. Both the sign and size of contributions from higher-spin prehadrons are difficult to determine. In addition, the RGEs of α_4 , α_L , and α_R are affected by purely nonperturbative effects due to $SU(15)_p$ dynamics, which presently cannot be computed in chiral gauge theories.

To obtain a rough lower bound for α_4 , α_L , and α_R at a scale above preon confinement, we estimate the change in the gauge coupling due to the spin-1 premesons listed in Table IV. The question is over what range of scales do the spin-1 premesons act as relevant degrees of freedom? For a rough estimate, we use the information provided by QCD, although one should keep in mind that the behavior of spin-1 bound state in our chiral preon dynamics may be very different from QCD. The measured mass of the ρ mesons is $m_\rho \approx 770$ MeV, while the scale where quarks and gluons become perturbative particles is around 2 GeV (see, e.g., [40]). Thus, in QCD the ρ meson is a relevant particle up to a scale of about $2.6m_\rho$.

Note that the running of the electromagnetic coupling constant α is pushed by ρ mesons towards larger values because QED is an Abelian gauge theory. Therefore, we cannot use information about α to draw conclusions about α_4 , α_L , and α_R (for a lattice study of QCD effects on α , see [44]).

For a preonic confinement scale of $\Lambda_{\text{pre}} \approx 3 \times 10^3$ TeV, consistent with proton decay constraints from Ref. [6], the gauge coupling constants obtained in Sec. IV B are

$$\begin{aligned} \alpha_4(\Lambda_{\text{pre}}) &= 0.071, & \alpha_L(\Lambda_{\text{pre}}) &= 0.042, \\ \alpha_R(\Lambda_{\text{pre}}) &= 0.013. \end{aligned} \quad (4.13)$$

If the influence of the ρ premesons persists up to a scale $\Lambda'_{\text{pre}} \approx 2.6\Lambda_{\text{pre}} \approx 8 \times 10^3$ TeV (analogous to the end point of the ρ meson regime in QCD), then we find that the one-loop running gives the following coupling ratios:

$$\frac{\alpha_4^{-1}(\Lambda'_{\text{pre}})}{\alpha_4^{-1}(\Lambda_{\text{pre}})} = 6.1, \quad \frac{\alpha_L^{-1}(\Lambda'_{\text{pre}})}{\alpha_L^{-1}(\Lambda_{\text{pre}})} = 3.2, \quad \frac{\alpha_R^{-1}(\Lambda'_{\text{pre}})}{\alpha_R^{-1}(\Lambda_{\text{pre}})} = 1.7. \quad (4.14)$$

These large ratios illustrate the substantial impact that the premesons have on the running of gauge couplings. Thus, $SU(15)_p$ dynamics may substantially push up the three α^{-1} running curves shown in Fig. 6 once they enter the right-handed (gray) shaded region.

At scales beyond Λ_{pre} , where the preons rather than the prehadrons are the relevant degrees of freedom, the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ interactions lose again asymptotic freedom (see discussion in Sec. II B).

However, the decrease of the couplings due to $SU(15)_p$ dynamics around Λ_{pre} may be sufficient to keep α_4 , α_L , and α_R perturbative up to a unification scale Λ_{422} . There, $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ may be embedded in the $SO(10)$ gauge group, which is large enough to have asymptotic freedom in the presence of all of the preons.

Whether the three couplings unify at the Λ_{422} scale depends on various threshold effects and symmetry breaking patterns [45]. Compared to usual, perturbative GUT models, the unification scale can be higher (e.g., $\Lambda_{422} \approx 10^{17}$ GeV) in the framework discussed here due to the intricate running of the three gauge couplings. As a result, Planck-scale suppressed operators may be large enough to already split α_4 , α_L , and α_R at Λ_{422} .

V. CONCLUSIONS

We have developed a comprehensive framework for quark and lepton compositeness, based on an $SU(15)_p$ chiral gauge theory that confines preons. This theory not only yields three SM generations of composite quarks and leptons (as three-preon bound states), but also provides a dynamical origin for symmetry breaking and the Higgs sector (as six-preon bound states). The preons are also charged under the weakly coupled $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge group, which is broken down to the SM gauge group at a scale Λ_{PS} in the 30–100 TeV range.

By embedding the QCD gauge group within $SU(4)_{\text{PS}}$, the ultraviolet behavior of the strong coupling is moderated, at least up to the preon confinement scale Λ_{pre} , which is about 2 orders of magnitude above Λ_{PS} . Our analysis of renormalization group evolution, incorporating the effects of composite vectorlike fermions and scalar di-prebaryons, shows that each of the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ interactions lose asymptotic freedom below Λ_{pre} . Near the compositeness scale there are many spin-1 premesons whose effect is to decrease the gauge couplings. If that effect is not fully counterbalanced by other prehadrons and nonperturbative effects at Λ_{pre} that cannot be currently computed, the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge group may remain under perturbative control up to a unification scale, where it may be embedded in $SO(10)$.

The theory opens avenues for novel mechanisms of quark and lepton mass generation rooted in underlying preon dynamics. It also predicts composite vectorlike leptons not far above the TeV scale, as well as an extended Higgs sector, offering observable signatures at the LHC and future colliders. We have found that the vectorlike fermion spectrum is modified in the presence of the $SU(4)_{\text{PS}} \times SU(2)_L \times SU(2)_R$ gauge interactions, so the collider probes of this theory are different compared to the case where the weakly coupled gauge symmetry is just the SM one [2,6]. Moreover, the composite neutrino sector is complex, with several sterile neutrinos that may have interesting phenomenological implications.

There remain, however, many important questions, especially related to the behavior of the strongly coupled chiral $SU(15)_p$ gauge interactions. For example, the size of the VEVs of di-prebaryons is highly dependent on the coupling strength in the attractive channels. Consequently, a quantitative estimate of the mass spectrum for the composite vectorlike fermions is hampered by the lack of information regarding how close to the critical value is the binding due to premeson exchange.

Furthermore, the possibility that the preonic bound states associated with the Higgs sector have masses several orders of magnitude below Λ_{pre} needs to be put on a firmer ground. Nevertheless, the framework presented here offers a promising path for the preonic $SU(15)_p$ chiral gauge dynamics to be a realistic substructure origin of the SM.

ACKNOWLEDGMENTS

We would like to thank Jure Zupan and Ryan Plestid for helpful conversations. B. A. acknowledges support in part by the DOE Grant No. DE-SC1019775, and the NSF Grants No. OAC-2103889, No. OAC-2411215, and No. OAC-2417682. The work of B. A. was performed in part at the Aspen Center for Physics, with support by a grant from the Simons Foundation (1161654, Troyer). The work of B. D. was supported by Fermi Forward Discovery Group, LLC under Contract No. 89243024CSC000002 with the U.S. Department of Energy, Office of Science, Office of High Energy Physics.

DATA AVAILABILITY

No data were created or analyzed in this study.

-
- [1] G. 't Hooft, Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking, in *Recent Developments in Gauge Theories*, NATO Advanced Study Institutes Series Vol. 59 (Springer, Boston, MA, 1980), p. 135.
 - [2] B. A. Dobrescu, Quark and lepton compositeness: A renormalizable model, *Phys. Rev. Lett.* **128**, 241804 (2022).
 - [3] J. C. Pati, A. Salam, and J. A. Strathdee, A Preon model with hidden electric and magnetic type charges, *Nucl. Phys.* **B185**, 416 (1981).
 - [4] I. Bars and S. Yankielowicz, Composite quarks and leptons as solutions of anomaly constraints, *Phys. Lett.* **101B**, 159 (1981); I. Bars, Family structure with composite quarks and leptons, *Phys. Lett.* **106B**, 105 (1981).
 - [5] C. Q. Geng and R. E. Marshak, Two realistic preon models with $SU(N)$ metacolor satisfying complementarity, *Phys. Rev. D* **35**, 2278 (1987).
 - [6] B. Assi and B. A. Dobrescu, Proton decay from quark and lepton compositeness, *J. High Energy Phys.* **12** (2022) 116.
 - [7] J. C. Pati and A. Salam, Lepton number as the fourth color, *Phys. Rev. D* **10**, 275 (1974); **11**, 703(E) (1975).
 - [8] D. B. Kaplan, Chiral gauge theory at the boundary between topological phases, *Phys. Rev. Lett.* **132**, 141603 (2024); D. B. Kaplan and S. Sen, Weyl fermions on a finite lattice, *Phys. Rev. Lett.* **132**, 141604 (2024); S. Aoki, H. Fukaya, and N. Kan, A lattice formulation of Weyl fermions on a single curved surface, *Prog. Theor. Exp. Phys.* **2024**, 043B05 (2024); M. Golterman and Y. Shamir, Conserved currents in five-dimensional proposals for lattice chiral gauge theories, *Phys. Rev. D* **109**, 114519 (2024); D. B. Kaplan and S. Sen, Regulating chiral gauge theory at $\theta = 0$, [arXiv:2412.02024](https://arxiv.org/abs/2412.02024).
 - [9] E. Eichten, K. Kang, and I. G. Koh, Anomaly free complex representations in $SU(N)$, *J. Math. Phys. (N.Y.)* **23**, 2529 (1982).
 - [10] E. Eichten, R. D. Peccei, J. Preskill, and D. Zeppenfeld, Chiral gauge theories in the $1/N$ expansion, *Nucl. Phys.* **B268**, 161 (1986).
 - [11] P. B. Smith, A. Karasik, N. Lohitsiri, and D. Tong, On discrete anomalies in chiral gauge theories, *J. High Energy Phys.* **01** (2022) 112.
 - [12] A. Karasik, K. Önder, and D. Tong, Chiral gauge dynamics: Candidates for non-supersymmetric dualities, *J. High Energy Phys.* **11** (2022) 122.
 - [13] T. Appelquist, A. G. Cohen, M. Schmaltz, and R. Shrock, New constraints on chiral gauge theories, *Phys. Lett. B* **459**, 235 (1999).
 - [14] T. Appelquist, Z. y. Duan, and F. Sannino, Phases of chiral gauge theories, *Phys. Rev. D* **61**, 125009 (2000).
 - [15] S. Dimopoulos, S. Raby, and L. Susskind, Light composite fermions, *Nucl. Phys.* **B173**, 208 (1980).
 - [16] S. Bolognesi, K. Konishi, and A. Luzio, Strong anomaly and phases of chiral gauge theories, *J. High Energy Phys.* **08** (2021) 028.
 - [17] S. Bolognesi, K. Konishi, and A. Luzio, Anomalies and phases of strongly coupled chiral gauge theories: Recent developments, *Int. J. Mod. Phys. A* **37**, 2230014 (2022); The $\mathbb{Z}_2$ anomaly in some chiral gauge theories, *J. High Energy Phys.* **08** (2023) 125.
 - [18] C. Csáki, H. Murayama, and O. Telem, More exact results on chiral gauge theories: The case of the symmetric tensor, *Phys. Rev. D* **105**, 045007 (2022).
 - [19] H. Collins, A. K. Grant, and H. Georgi, Dynamically broken topcolor at large N , [arXiv:hep-ph/9907477](https://arxiv.org/abs/hep-ph/9907477); W. A. Bardeen, C. T. Hill, and D. U. Jungnickel, Chiral hierarchies, compositeness and the renormalization group, *Phys. Rev. D* **49**, 1437 (1994).
 - [20] W. A. Bardeen, C. T. Hill, and M. Lindner, Minimal dynamical symmetry breaking of the standard model, *Phys. Rev. D* **41**, 1647 (1990); C. T. Hill, Topcolor: Top quark

- condensation in a gauge extension of the standard model, *Phys. Lett. B* **266**, 419 (1991).
- [21] C. T. Hill, A new-old approach to composite scalars with chiral fermion constituents, *Nucl. Phys. B* **1011**, 116788 (2025); Natural top quark condensation (a Redux), *arXiv*: 2503.21518.
- [22] B. A. Dobrescu and C. T. Hill, Electroweak symmetry breaking via top condensation seesaw, *Phys. Rev. Lett.* **81**, 2634 (1998); R. S. Chivukula, B. A. Dobrescu, H. Georgi, and C. T. Hill, Top quark Seesaw theory of electroweak symmetry breaking, *Phys. Rev. D* **59**, 075003 (1999); B. A. Dobrescu, Minimal composite Higgs model with light bosons, *Phys. Rev. D* **63**, 015004 (2001).
- [23] S. Raby, S. Dimopoulos, and L. Susskind, Tumbling gauge theories, *Nucl. Phys. B* **169**, 373 (1980).
- [24] A. Tumasyan *et al.* (CMS Collaboration), Inclusive non-resonant multilepton probes of new phenomena at $\sqrt{s} = 13$ TeV, *Phys. Rev. D* **105**, 112007 (2022).
- [25] G. Aad *et al.* (ATLAS Collaboration), Search for third-generation vector-like leptons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **07** (2023) 118.
- [26] P. N. Bhattiprolu and S. P. Martin, Prospects for vectorlike leptons at future proton-proton colliders, *Phys. Rev. D* **100**, 015033 (2019).
- [27] E. Bernreuther and B. A. Dobrescu, Vectorlike leptons and long-lived bosons at the LHC, *J. High Energy Phys.* **07** (2023) 079; V. Chekhovsky *et al.* (CMS Collaboration), Search for vector-like leptons with long-lived particle decays in the CMS muon system in proton-proton collisions at $\sqrt{s} = 13$ TeV, *arXiv*:2503.16699.
- [28] H. Georgi, H. R. Quinn, and S. Weinberg, Hierarchy of interactions in unified gauge theories, *Phys. Rev. Lett.* **33**, 451 (1974).
- [29] G. 't Hooft, Dimensional regularization and the renormalization group, *Nucl. Phys. B* **61**, 455 (1973).
- [30] J. C. Collins and A. J. Macfarlane, New methods for the renormalization group, *Phys. Rev. D* **10**, 1201 (1974).
- [31] R. N. Mohapatra and J. C. Pati, A natural left-right symmetry, *Phys. Rev. D* **11**, 2558 (1975).
- [32] M. E. Machacek and M. T. Vaughn, Two loop renormalization group equations in a general quantum field theory. 1. Wave function renormalization, *Nucl. Phys. B* **222**, 83 (1983); Two loop renormalization group equations in a general quantum field theory. 2. Yukawa couplings, *Nucl. Phys. B* **236**, 221 (1984).
- [33] I. Jack and H. Osborn, General background field calculations with fermion fields, *Nucl. Phys. B* **249**, 472 (1985).
- [34] A. G. M. Pickering, J. A. Gracey, and D. R. T. Jones, Three loop gauge beta function for the most general single gauge coupling theory, *Phys. Lett. B* **510**, 347 (2001); **535**, 377(E) (2002).
- [35] A. Bednyakov and A. Pikelner, Four-loop gauge and three-loop Yukawa beta functions in a general renormalizable theory, *Phys. Rev. Lett.* **127**, 041801 (2021).
- [36] F. Staub, SARAH 4: A tool for (not only SUSY) model builders, *Comput. Phys. Commun.* **185**, 1773 (2014).
- [37] A. E. Thomsen, Introducing RGBeta: A mathematica package for the evaluation of renormalization group β -functions, *Eur. Phys. J. C* **81**, 408 (2021).
- [38] B. A. Kniehl, A. F. Pikelner, and O. L. Veretin, mr: A C++ library for the matching and running of the standard model parameters, *Comput. Phys. Commun.* **206**, 84 (2016).
- [39] G. Huang and S. Zhou, Precise values of running quark and lepton masses in the standard model, *Phys. Rev. D* **103**, 016010 (2021).
- [40] S. Navas *et al.* (Particle Data Group), Review of particle physics, *Phys. Rev. D* **110**, 030001 (2024).
- [41] M. J. Dolan, T. P. Dutka, and R. R. Volkas, Lowering the scale of Pati-Salam breaking through seesaw mixing, *J. High Energy Phys.* **05** (2021) 199.
- [42] G. Valencia and S. Willenbrock, Quark—lepton unification and rare meson decays, *Phys. Rev. D* **50**, 6843 (1994).
- [43] L. Calibbi, A. Crivellin, and T. Li, Model of vector leptoquarks in view of the B -physics anomalies, *Phys. Rev. D* **98**, 115002 (2018); S. Iguro, J. Kawamura, S. Okawa, and Y. Omura, TeV-scale vector leptoquark from Pati-Salam unification with vectorlike families, *Phys. Rev. D* **104**, 075008 (2021); Importance of vector leptoquark-scalar box diagrams in Pati-Salam unification with vectorlike families, *J. High Energy Phys.* **07** (2022) 022.
- [44] F. Jegerlehner, Electroweak effective couplings for future precision experiments, *Nuovo Cimento Soc. Ital. Fis.* **34C**, 31 (2011).
- [45] C. Giunti, C. W. Kim, and U. W. Lee, Running coupling constants and grand unification models, *Mod. Phys. Lett. A* **06**, 1745 (1991).