

Unimodular Jackiw-Teitelboim gravity and de Sitter quantum cosmology

Bruno Alexandre^{1,*}, Altay Etkin^{2,†} and Farbod-Sayyed Rassouli^{3,‡}

¹*Abdus Salam Centre for Theoretical Physics, Imperial College London,
Prince Consort Road, London SW7 2AZ, United Kingdom*

²*School of Mathematical Sciences and STAG Research Centre, University of Southampton,
Highfield Campus, Southampton SO17 1BJ, United Kingdom*

³*School of Physics and Astronomy and Nottingham Centre of Gravity, University of Nottingham,
University Park, Nottingham NG7 2RD, United Kingdom*



(Received 13 March 2025; accepted 10 June 2025; published 25 June 2025)

In this work, we show that a gauge-theoretic description of Jackiw-Teitelboim (JT) gravity naturally yields a Henneaux-Teitelboim (HT) unimodular gravity via a central extension of its isometry group, valid for both flat and curved two-dimensional spacetimes. HT gravity introduces a unimodular time canonically conjugate to the cosmological constant, serving as a physical time in quantum cosmology. By studying the minisuperspace reduction of HT_2 gravity, the Wheeler-DeWitt equation becomes a Schrödinger-like equation, giving a consistent and unitary quantum theory. Analysis of the wave function's probability density reveals a quantum distribution for the scale factor a , offering a quantum perspective on the expansion and contraction of the Universe. In this perspective, the possibility of reaching the singular point $a = 0$ signals that topology change could occur. Finally, we give a consistent quantum description of unimodular time that aligns seamlessly with Page-Wootters formulation of quantum mechanics, where quantum correlations between unimodular time and JT gravity are studied in HT_2 quantum cosmology.

DOI: [10.1103/cmbb-cwlj](https://doi.org/10.1103/cmbb-cwlj)

I. INTRODUCTION

One of the profound challenges in theoretical physics is developing a quantum-mechanical framework that encompasses the entire Universe. Indeed, the incompatibility between general relativity (GR) and quantum mechanics (QM) is particularly evident in the well-known problem of time, which arises due to the fundamentally different treatment of time in both theories [1–5]. In GR, time is treated as part of the spacetime fabric and, therefore, the theory is diffeomorphism invariant. Conversely, in QM, time is treated as an external, absolute parameter used to measure the evolution of a quantum system. Indeed, the Hamiltonian constraint of a gravitational system vanishes, therefore leading to the absence of unitary evolution in terms of quantum states. Significant efforts have focused on the role of time in quantum gravity, with three main approaches emerging: solving canonical constraints before

quantization, thus defining time classically; quantizing first and identifying time afterward through the Wheeler-DeWitt equation; or treating time purely phenomenologically, without fundamental quantum status.

A more recent and promising avenue is Henneaux-Teitelboim (HT) gravity where, by converting the constants of nature to dynamical variables, one gets physical relational times which are the canonical conjugates of the dynamical “constants” [6–19]. Therefore, we end up having a physical time variable conjugate to a dynamical cosmological constant [20–26], which preserves unitarity and allows the construction of normalizable wave packets [27–29]. This not only converts the Wheeler-DeWitt equation into a Schrödinger-like equation, but also resolves the apparent unitarity problem and gives conserved probabilities.

In this regard, we study Jackiw-Teitelboim (JT) gravity as a two-dimensional toy model for quantum gravity and quantum cosmology. Ordinarily, in JT gravity the Hamiltonian constraint is found to vanish, resulting in “frozen” quantum states. In turn, this exacerbates the problem of the absence of unitary time evolution. The missing of an external time in (de Sitter) JT gravity has been extensively discussed in the context of the Chandrasekaran-Longo-Penington-Witten (CLPW) framework of algebras of observables [30–34]. In this framework, observers are treated as clocks within the gravitating system, whose role is to fix the time translation symmetry.

*Contact author: bruno.alexandre20@imperial.ac.uk

†Contact author: a.etkin@soton.ac.uk

‡Contact author: sayyed.rassouli@nottingham.ac.uk

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Put simply, an observer is any system capable of measuring time. Recently, it has also been demonstrated that the CLPW algebras of observables is equivalent to the Page-Wootters (PW) approach [35–41] where the total system is described by correlations between the observer and the gravitational system.

In this paper, we show that a physical clock naturally emerges from an HT formulation of JT gravity, which in turn stems from a gauge-theoretic approach—a connection previously unrecognized in the literature [such as the extended Callan-Giddings-Harvey-Strominger (CGHS) model discussed in two-dimensional flat holography] [42–51]. In this scenario, introducing unimodular time allows the PW formalism to reach its full potential, providing a robust framework in which a genuine notion of time evolution can be defined. Moreover, the ability to define such a physical clock in a controlled, lower-dimensional model of gravity opens new avenues for studying quantum cosmology. In particular, it is especially enticing that, for de Sitter space, one can study time evolution of closed universes and topology change at early times.

The paper is organized as follows: in Sec. II, we show how HT gravity naturally emerges from a gauge formulation of JT gravity via central extensions of the isometry group. In Sec. III, we explore this formulation in minisuperspace (MSS), where we quantize the theory and build normalizable unimodular wave packets. We study the probability density associated with such wave packets in Sec. IV and comment on the global topology of the spacetime. In Sec. V, we introduce the Page-Wootters formalism of QM and apply it to the work done in the previous sections. We conclude with some final remarks in Sec. VI.

II. HENNEAUX-TEITELBOIM GRAVITY FROM CENTRAL EXTENSIONS OF JT GRAVITY

It is well established that GR admits a volume-preserving diffeomorphism-invariant formulation, commonly referred to as unimodular gravity [6–19]. This formulation is classically equivalent to GR and is achieved by imposing the gauge fixing condition $\sqrt{-g} = 1$. The seminal work of Henneaux and Teitelboim extended this framework by restoring full diffeomorphism invariance in unimodular gravity through the promotion of constants of nature to dynamical variables [20–26]. This parametrization becomes evident when the unimodular condition is modified to $\sqrt{-g} = \partial_\mu T^\mu$, treated as an on-shell expression, with action given by

$$S = \frac{1}{2} \int d^4x \sqrt{-g} (R - 2\Lambda) + \int d^4x \Lambda \partial_\mu T^\mu, \quad (1)$$

where T^μ is an auxiliary vector density, and Λ is the vacuum energy field (also known as the cosmological constant) promoted to an off-shell variable and on-shell constant. This




FIG. 1. Diagram illustrating various reductions of the general HT₂ gravity theory. The double arrows indicate the gauge formulations of the KSY and CJ gravity models.

theory yields equations of motion identical to those of the standard Einstein-Hilbert action.

Recent work [52,53] has demonstrated that an Abelian reduction based on the gauge group U(1) yields an HT gravity theory, where the HT term is replaced by the Chern-Pontryagin topological term. The resultant action is

$$S = \frac{1}{2} \int d^4x \sqrt{-g} (R - 2\Lambda) - \frac{1}{8\pi^2} \int \Lambda F \wedge F, \quad (2)$$

with $F = dA$ the curvature 2-form associated with an Abelian U(1) gauge field A and with the vector density T^μ identified as the Chern-Simons current.

Although typically studied in four dimensions, HT gravity also emerges in certain two-dimensional contexts, though not recognized as such. In particular, various formulations of JT gravity—such as those employed in flat holography [54–61]—are revealed to be HT gravity theories [42–51,62,63] in this paper.

This structure arises naturally when JT gravity is formulated from a gauge-theoretic perspective (see Fig. 1). The standard JT action, without the inclusion of boundary terms, is given by

$$S_{\text{JT}} = \frac{1}{2} \int d^2x \sqrt{-g} \phi (R - 2\lambda), \quad (3)$$

where λ is the “cosmological constant” that ensures that, upon variation of the dilaton field ϕ , we have the on-shell expression $R = 2\lambda$. The dilaton equation of motion is obtained from the metric variation of the action

$$(\nabla_\mu \partial_\nu - g_{\mu\nu} \square) \phi = \lambda g_{\mu\nu} \phi. \quad (4)$$

The global spacetime geometry depends on the value of λ : it is de Sitter (dS) for $\lambda > 0$ and anti-de Sitter for $\lambda < 0$.

Of particular interest is the JT gravity action (3) with nonvanishing λ , which possesses a gauge-theoretical formulation based on the $\mathfrak{sl}(2, \mathbb{R})$ algebra [64–67],

$$\begin{aligned} [P_a, P_b] &= 2\lambda J_{ab} \equiv -\lambda \epsilon_{ab} \tilde{J}, \\ [P_a, \tilde{J}] &= -\epsilon_a{}^b P_b. \end{aligned} \quad (5)$$

The resulting first-order formulation is a BF theory with action given by

$$S = \int \langle \mathbf{B}, \mathbf{F} \rangle = \int B_a T^a + \tilde{B}(\tilde{R} - \lambda \tilde{\Sigma}), \quad (6)$$

where $\mathbf{B} = B^a P_a + \tilde{B} \tilde{J}$ is a 0-form Lagrange multiplier, and

$$\mathbf{F} = d\mathbf{A} + \mathbf{A} \wedge \mathbf{A} = T^a P_a + \tilde{R} \tilde{J} \quad (7)$$

is the curvature 2-form associated with the connection 1-form $\mathbf{A} = e^a P_a + \tilde{\omega} \tilde{J}$ for the $\mathfrak{sl}(2, \mathbb{R})$ algebra. Here, the Einstein-Cartan (EC) variables are the zweibein e^a and dual spin connection $\tilde{\omega} = \frac{1}{2} \epsilon^{ab} \omega_{ab}$, with associated torsion $T^a = De^a$ and dual curvature $\tilde{R} = \frac{1}{2} \epsilon^{ab} R_{ab} = d\tilde{\omega}$. The dual area 2-form is $\tilde{\Sigma} = \frac{1}{2} \epsilon_{ab} \Sigma^{ab}$, where $\Sigma^{ab} = e^a \wedge e^b$. In this context, one can show that the on-shell action reduces to an Einstein-Cartan action for JT gravity

$$S_{\text{JT}} = \int \phi(\tilde{R} - \lambda \tilde{\Sigma}), \quad (8)$$

with vanishing torsion imposed by the Lagrange multiplier B^a and $\tilde{B}$ identified with the dilaton ϕ .

Building on top of the requirements for an HT gravity theory, one can formulate a two-dimensional unimodular JT gravity model (HT₂), described by the action

$$S_{\text{HT}_2} := \frac{1}{2} \int d^2x \sqrt{-g} (\phi(R - 2\lambda) - 2\Lambda) + \int d^2x \Lambda \partial_\mu T^\mu. \quad (9)$$

The distinction between λ and Λ is crucial: λ determines the global spacetime geometry on shell, while Λ serves as the vacuum energy field, similar to its role in four-dimensional HT gravity. In this section, we demonstrate that the action (9) emerges naturally from a gauge-invariant formulation of JT gravity and study its features.

A. HT₂ as a gauge-invariant formulation of JT gravity

We want to show that a gauge formulation of (9) arises from a BF formulation, akin to JT gravity. For that, we follow the Kim-Soh-Yee (KSY) model [47,48], where they consider an extension of JT gravity via a vacuum cosmological constant term,

$$S_{\text{KSY}} = \frac{1}{2} \int d^2x \sqrt{-g} (\phi(R - 2\lambda) - 2\Lambda). \quad (10)$$

The on-shell expressions are obtained by varying the action with respect to ϕ and $g_{\mu\nu}$, respectively,

$$R = 2\lambda, \quad (11)$$

$$(\nabla_\mu \partial_\nu - g_{\mu\nu} \square) \phi = (\lambda \phi + \Lambda) g_{\mu\nu}. \quad (12)$$

A gauge-invariant formulation of this model with $\lambda \neq 0$ is based on an Abelian central extension of the isometry group of spacetime, that is a BF theory with Lie algebra based on the extended de Sitter algebra,

$$\widehat{\mathfrak{sl}}(2, \mathbb{R}) \cong \mathfrak{sl}(2, \mathbb{R}) \oplus \mathfrak{u}(1), \quad (13)$$

and commutation relations given by

$$[P_a, P_b] = -\lambda \epsilon_{ab} \tilde{J} + \epsilon_{ab} I, \quad (14)$$

$$[P_a, \tilde{J}] = -\epsilon_a{}^b P_b, \quad (15)$$

where I is the central element given by the $\mathfrak{u}(1)$ subalgebra. We want to note that this central extension is trivial as it is merely a direct sum construction.

With this algebra, one can construct an $\widehat{\mathfrak{sl}}(2, \mathbb{R})$ -valued connection 1-form,

$$\mathbf{A} = e^a P_a + \tilde{\omega} \tilde{J} + AI, \quad (16)$$

and associated curvature 2-form given by

$$\mathbf{F} = T^a P_a + (\tilde{R} - \lambda \tilde{\Sigma}) \tilde{J} + (F + \tilde{\Sigma}) I. \quad (17)$$

Here, $F = dA$ is the $\mathfrak{u}(1)$ -valued curvature 2-form with associated connection 1-form A . Using the above curvature $\mathbf{F}$ and defining the Lagrange multiplier 0-form $\mathbf{B} = B^a P_a + \tilde{B} \tilde{J} + BI$, the BF action reduces to

$$S = \int B_a T^a + \phi(\tilde{R} - \lambda \tilde{\Sigma}) - \Lambda(F + \tilde{\Sigma}), \quad (18)$$

where we set $B = \phi + \Lambda$ and $\tilde{B} = -\lambda \phi - \Lambda$ using the trace identities induced from the Casimir invariant (see [50]). The vanishing of torsion gives the on-shell action, corresponding to an EC formulation of a gauge-invariant first-order action for the KSY model given by

$$S_{\text{HT}_2} = \int \phi(\tilde{R} - \lambda \tilde{\Sigma}) - \Lambda(F + \tilde{\Sigma}). \quad (19)$$

Transitioning to the second-order formulation, we recover the same action as in (9), which is

$$S_{\text{HT}_2} = \frac{1}{2} \int d^2x \sqrt{-g} \phi(R - 2\lambda) + \int d^2x \Lambda (\epsilon^{\mu\nu} \partial_\mu A_\nu - \sqrt{-g}), \quad (20)$$

where the vector density $\mathcal{T}^\mu$ is identified with the dual of the Abelian gauge field, $\mathcal{T}^\mu = \epsilon^{\mu\nu} A_\nu$. Here, $\epsilon^{\mu\nu}$ denotes the Levi-Civita symbol.¹

The variation of the action with respect to ϕ , $g_{\mu\nu}$, $\mathcal{T}^\mu$, and Λ , respectively, yields the following equations of motion:

$$R = 2\lambda, \quad (21)$$

$$(\nabla_\mu \partial_\nu - g_{\mu\nu} \square) \phi = (\lambda \phi + \Lambda) g_{\mu\nu}, \quad (22)$$

$$\partial_\mu \Lambda = 0, \quad (23)$$

$$\sqrt{-g} = \partial_\mu \mathcal{T}^\mu. \quad (24)$$

From the first three equations of motion, the on-shell dynamics of HT₂ match those of the KSY model, provided that the unimodular condition (24) is satisfied. Therefore, the classical dynamics remain unchanged and they describe the same physics [8,14]. Indeed, Λ is promoted to a phase-space variable, canonically conjugate to the zero mode of $\mathcal{T}^0$, which is referred to as unimodular time [8]

$$T_\Lambda(t) := \int_{\Sigma_t} d\Sigma \mathcal{T}^0 \Big|_{\Sigma_t} = - \int_{\Sigma_t} d\Sigma A_1 \Big|_{\Sigma_t}, \quad (25)$$

where Σ_t are constant time hypersurfaces and $\mathcal{T}^0 = -A_1$. This definition of unimodular time guarantees that it represents a genuine, physical “flow of time.” Identifying ΔT_Λ with the spacetime volume lying in the past of the leaf Σ_t —measured relative to a conventional initial leaf Σ_0 —one finds the on-shell relation

$$\Delta T_\Lambda = \int_{\Sigma_0}^{\Sigma_t} d^2x \sqrt{-g} =: \text{Vol}(\Sigma_t \cup \Sigma_0) \quad (26)$$

as established in [20–26]. Because the spacetime volume swept out between the two hypersurfaces is strictly positive, T_Λ is necessarily monotonic, assuming nonsingular manifolds.

Simple counting of constraints shows that the theory has no new local degrees of freedom, justifying the appearance of the topological term $\partial_\mu \mathcal{T}^\mu$. There is an additional U(1) gauge symmetry of the HT₂ action: a gauge transformation of $\mathcal{T}^\mu$,

$$\mathcal{T}^\mu \rightarrow \mathcal{T}^\mu + \chi^\mu, \quad (27)$$

with $\partial_\mu \chi^\mu = 0$ ensuring that there is one gauge degree of freedom left. This implies the invariance of the action (20).

It is important to notice that, unlike its four-dimensional cousin, the U(1) reduction of HT gravity (2) in two dimensions is naturally emergent from the extended

algebra. The Chern-Pontryagin term $F \wedge F$ in four dimensions is therefore naturally replaced in two dimensions with the first Chern class F .

B. Maxwell algebra and CGHS model as flat HT₂ gravity

In this section, we note that setting $\lambda = 0$ in the KSY model (10) yields what is known in the literature as the Cangemi-Jackiw (CJ) gravity model [42], which can also be viewed as a Weyl rescaling of the Callan-Giddings-Harvey-Strominger (CGHS) model [62]. Recently, growing interest in two-dimensional flat holography called the CGHS model and its close connection to the complex Sachdev-Ye-Kitaev model [54–57,59–61,68] have reignited attention toward this model. Indeed, to study CJ theory at different temperatures, it is necessary to allow the vacuum cosmological constant Λ to vary—precisely as achieved by the unimodular extension—which controls the Rindler inverse temperature $\beta \sim \frac{1}{\Lambda}$.

The CJ action is given by

$$S_{\text{CJ}} = \frac{1}{2} \int d^2x \sqrt{-g} (\phi R - 2\Lambda), \quad (28)$$

with on-shell expressions given by

$$R = 0, \quad (29)$$

$$(\nabla_\mu \partial_\nu - g_{\mu\nu} \square) \phi = \Lambda g_{\mu\nu}. \quad (30)$$

A consistent gauge construction in terms of Einstein-Cartan variables does exist for the CJ model [42–45,49]. For this, we take the gauge group as the isometry group of flat two-dimensional spacetime, $\text{ISO}^+(1, 1)$, and centrally extend it via U(1). The resulting Maxwell algebra $\widehat{\mathfrak{iso}}(1, 1)$ has corresponding commutation relations given by the $\lambda = 0$ limit of (14),

$$[P_a, P_b] = \epsilon_{ab} I, \quad [P_a, \tilde{J}] = -\epsilon_a^b P_b. \quad (31)$$

With the algebra defined as above, one can construct a gauge theory based on $\widehat{\mathfrak{iso}}(1, 1)$. Proceeding in a similar manner to the previous section, the $\lambda = 0$ curvature $\mathbf{F}$ is

$$\mathbf{F} = T^a P_a + \tilde{R} \tilde{J} + (F + \tilde{\Sigma}) I. \quad (32)$$

The BF action (18) reduces to

$$S = \int B_a T^a + \phi \tilde{R} - \Lambda (F + \tilde{\Sigma}), \quad (33)$$

where now $B = \phi + \Lambda$ and $\tilde{B} = -\Lambda$. Therefore, the on-shell action is given by the flat EC formulation of CJ gravity model,

¹We denote $\epsilon^{\mu\nu}$ by the Levi-Civita tensor given by $\epsilon^{\mu\nu} = \frac{\epsilon^{\mu\nu}}{\sqrt{-g}}$

$$S_{\text{CGHS}} = \int \phi \tilde{R} - \Lambda(F + \tilde{\Sigma}) = \frac{1}{2} \int d^2x \sqrt{-g} [\phi R + 2\Lambda(\epsilon^{\mu\nu} \partial_\mu A_\nu - 1)]. \quad (34)$$

Variation with respect to the dilaton and the metric yields the same classical dynamics as CJ gravity, establishing that flat HT₂ ($\lambda = 0$) and CJ gravity are on-shell equivalent theories. Moreover, the unimodular condition for flat HT₂ gravity remains unchanged from the $\lambda \neq 0$ HT₂ gravity, given by $\partial_\mu T^\mu = \sqrt{-g}$.

III. QUANTUM COSMOLOGY IN DE SITTER HT₂ GRAVITY

As an application of HT₂ gravity, we explore the quantum cosmology of closed universes in de Sitter space. To begin, we introduce the MSS metric of dS in KSY gravity and highlight the differences arising from the HT₂ model. Subsequently, we perform a canonical quantization of both theories, with a particular focus on the implications for the probability amplitudes associated with the wave function of the universe.

A. Minisuperspace KSY gravity

To construct the MSS KSY gravity action, the starting point is through its associated EC formulation (19) with constant Λ ,

$$S_{\text{KSY}} = \int \phi(\tilde{R} - \tilde{\Sigma}) - \Lambda \tilde{\Sigma}, \quad (35)$$

where Λ is now just the vacuum cosmological constant and $\lambda = 1$ gives the globally de Sitter spacetime geometry. Note that we are using the more general KSY model—which contains the cosmological constant term—rather than standard JT gravity, in order to introduce the HT term in the next section. In two dimensions, the MSS metric takes a rather simple form,

$$ds^2 = -N^2 dt^2 + a^2 d\Sigma^2, \quad (36)$$

where $N(t)$ is the lapse function (usually gauge fixed to unity) and $a(t)$ is the scale factor that often appears in cosmological models. $d\Sigma^2$ is the metric on the homogeneous and isotropic, constant-time slice, spatial hypersurface Σ .

From the on-shell condition $R = 2$, we obtain a global de Sitter geometry. Consequently, the natural choice of coordinates is global dS₂, where spatial hypersurfaces are closed universes described by circles S^1 : we can identify $d\Sigma = d\theta$, where $\theta \in S^1$. The metric for a closed universe in global dS₂ coordinates is therefore given by

$$ds^2 = -dt^2 + \cosh^2(t) d\theta^2. \quad (37)$$

Geometrically, this describes a universe that is contracting at early times and expanding at later times, with a minimal length $a(0) = 1$ reached at $t = 0$, as the scale factor is simply $a(t) = \cosh(t)$. On the other hand, the equation of motion for the dilaton field will give the solution

$$\phi(t) = \phi_0 \sinh(t) - \Lambda. \quad (38)$$

Both the scale factor and dilaton expressions derived above are solutions to the Friedmann equations in a global de Sitter space.

Performing straightforward calculations, one finds that the dual area form is

$$\tilde{\Sigma} = Na dt \wedge d\theta, \quad (39)$$

and that the dual curvature form is given by

$$\tilde{R} = \dot{b} dt \wedge d\theta, \quad (40)$$

where $b = \dot{a}/N$ is written for convenience. Since nothing in the action depends on the angular parameter θ , we define the comoving volume as $v_c = \int d\theta$. Note that, in units of $v_c = 1$ (which we adopt in this paper), the scale factor a and the coordinate volume $v = av_c$ are the same. After integration by parts, the action (35) in MSS reduces to

$$S_{\text{KSY}} = - \int dt \left(\frac{\dot{\phi} \dot{a}}{N} + Na(\phi + \Lambda) \right), \quad (41)$$

where the boundary term cancels exactly with the Gibbon-Hawking-York (GHY) term

$$S_{\text{GHY}} = - \int_{\partial\mathcal{M}} d\theta \sqrt{|h|} \phi_b K = -\phi_b \frac{\dot{a}}{N} \Big|_{\partial\mathcal{M}}, \quad (42)$$

with ϕ_b the boundary value of the dilaton and K the extrinsic curvature.

The canonical momenta are easy to read off, each associated to the dilaton and scale factor, respectively,

$$p_\phi = -\frac{\dot{a}}{N}, \quad p_a = -\frac{\dot{\phi}}{N}. \quad (43)$$

Consequently, the Hamiltonian H of the MSS KSY gravity takes the form

$$H = -N p_a p_\phi + Na(\phi + \Lambda) \equiv aNC, \quad (44)$$

where C is the KSY gravity Hamiltonian constraint given by

$$C = -p_a \left(\frac{1}{a} \right) p_\phi + \phi + \Lambda. \quad (45)$$

The action in canonical form is therefore rewritten as

$$S_{\text{KSY}} = \int dt (p_\phi \dot{\phi} + p_a \dot{a} - aNC). \quad (46)$$

The Hamiltonian constraint equation $\mathcal{C} = 0$ is obtained as an on-shell expression when the action is varied with respect to the lapse function,

$$\mathcal{C} = -p_a \left(\frac{1}{a} \right) p_\phi + \phi + \Lambda = 0, \quad (47)$$

where we kept the order of the variables $p_a a^{-1} p_\phi$ so that when promoted to operators we ensure that the canonical quantization of the theory yields the Hartle-Hawking solution for the wave function of the universe, as shown below.

Starting from the canonical pairs, the equal-time Poisson brackets are given by

$$\{\phi, p_\phi\} = \{a, p_a\} = 1. \quad (48)$$

With the phase-space variables promoted to linear operators acting on the KSY Hilbert space, we have the following prescription on the momentum observables:

$$p_\phi \rightarrow \hat{p}_\phi = -i \frac{\partial}{\partial \phi}, \quad p_a \rightarrow \hat{p}_a = -i \frac{\partial}{\partial a}. \quad (49)$$

Under this quantization scheme, the Hamiltonian constraint is transformed into the Wheeler-DeWitt (WdW) equation for MSS KSY gravity,

$$\left[\partial_a \left(\frac{1}{a} \partial_\phi \right) + \frac{1}{2} U'(\phi) \right] \psi(a, \phi, \Lambda) = 0, \quad (50)$$

where we define the potential $U(\phi) = \phi^2 + 2\phi\Lambda$ and introduce ψ as the wave function of the universe.

A solution to the WdW equation described above is the so-called Hartle-Hawking (HH) wave function [69–81]. We shall consider here both the expanding and contracting branches of the HH wave function, in the asymptotic limit with $a\phi \gg 1$. The solution then becomes a linear superposition of plane waves

$$\psi(a, \phi, \Lambda) = \alpha_- \psi_-(a, \phi, \Lambda) + \alpha_+ \psi_+(a, \phi, \Lambda), \quad (51)$$

where $\psi_\pm(a, \phi, \Lambda) = e^{\pm ia\sqrt{U(\phi)}}$ corresponds to the contracting and expanding wave functions for plus and minus, respectively.

Note that from the start Λ is a fixed constant (off shell and on shell). In HT₂ gravity, this fixed constant is promoted to an off-shell variable, but on-shell free constant. In addition, one has the freedom to consider solutions of the form $\Psi(a, \phi, \Lambda) = \mathcal{A}(\Lambda)\psi(a, \phi, \Lambda)$ for some normalizable amplitude $\mathcal{A}(\Lambda)$, which also solves the WdW equation. An

issue involving the Hartle-Hawking solution is that it is not square integrable with respect to the DeWitt norm (a variant of the Klein-Gordon norm) [69,75,77,78]. This is partially solved by introducing normalizable amplitudes [74,81]. However, this requires a summation (or integration) over Λ , which is *a priori* fixed in JT gravity. Furthermore, the theory fails to evolve unitarily, as pointed out in [30–34], an issue tied to the absence of an observer in a closed universe. In the next section, we show that both problems are resolved by transitioning from the KSY model to HT₂ gravity.

An alternative approach to obtain the HH wave functions is through a semiclassical WKB approximation, where the wave function is given by

$$\psi(a, \phi, \Lambda) \sim e^{iS_{\text{on shell}}}, \quad (52)$$

where $S_{\text{on shell}}$ is the on-shell action for the KSY model. In this context, boundary terms play a crucial role, particularly the inclusion of the GHY term (and other possible holographic counterterms),

$$S_\partial = S_{\text{GHY}} + S_{\text{ct}}.$$

The presence of a spacelike boundary at time infinity is especially significant for a holographic treatment of the theory—for instance, the Schwarzschild modes residing at future infinity. However, such considerations lie beyond the scope of this paper. Accordingly, we shall disregard the boundary dynamics in our analysis and instead focus on investigating the bulk properties of MSS HT₂ gravity.

B. Quantization of MSS HT₂

As before, the Einstein-Cartan action for the two-dimensional Henneaux-Teitelboim gravity theory is given by (19), which in MSS takes the rather simple form:

$$S_{\text{HT}_2} = - \int dt \left(\frac{\dot{\phi} \dot{a}}{N} - \Lambda \dot{T}_\Lambda + Na(\phi + \Lambda) \right). \quad (53)$$

From the perspective of gauge theory, we have in the action $\dot{T}_\Lambda = -\dot{A}_\theta$. This is apparent in global dS₂ coordinates for a closed spatial slice S^1 , where we have the unimodular time explicitly written as

$$T_\Lambda = - \int_{S^1} A \Big|_{S^1} = -v_c A_\theta, \quad (54)$$

where in units of $v_c = 1$, $T_\Lambda = -A_\theta$. Throughout this section, we shall keep the notation T_Λ to denote the physical time variable.

In addition to the canonical momenta of the KSY model given in (43), we gain an extra canonical momentum conjugate to the unimodular time, as expected from HT₂ gravity,

$$p_{T_\Lambda} = \Lambda, \quad (55)$$

with equal-time Poisson bracket given by

$$\{T_\Lambda, \Lambda\} = 1. \quad (56)$$

The Hamiltonian of HT₂ is identical to that of KSY gravity. The action in its canonical form has the additional canonical momenta conjugate to the unimodular time,

$$S_{\text{HT}_2} = \int dt (p_\phi \dot{\phi} + p_a \dot{a} + p_{T_\Lambda} \dot{T}_\Lambda - aNC). \quad (57)$$

However, due to the fact that there exists a canonical momentum Λ conjugate to T_Λ , one has the on-shell expression

$$\mathcal{C} = C_{\text{JT}} + p_{T_\Lambda} = C_{\text{JT}} + \Lambda = 0, \quad (58)$$

where C_{JT} is the JT gravity Hamiltonian constraint,

$$C_{\text{JT}} = -p_a \left(\frac{1}{a} \right) p_\phi + \phi, \quad (59)$$

where we use again the correct ordering of the phase-space variables to reproduce the HH wave function solutions of the universe for HT₂ gravity.

Similar to the quantization of the MSS KSY model, we shall perform a canonical quantization of HT₂. We demonstrate how unitary evolution and a well-defined normalizable inner product emerge within HT₂ gravity, thus resolving the aforementioned issues in JT gravity.

We start by first using the prescription that the canonical momenta are replaced by derivatives with respect to their conjugate pairs,

$$p_\phi \longrightarrow \hat{p}_\phi = -i \frac{\partial}{\partial \phi}, \quad p_a \longrightarrow \hat{p}_a = -i \frac{\partial}{\partial a}, \quad (60)$$

$$p_{T_\Lambda} \longrightarrow \hat{p}_{T_\Lambda} = -i \frac{\partial}{\partial T_\Lambda}. \quad (61)$$

From (58), the Hamiltonian constraint then becomes in the T_Λ representation

$$\left[\partial_a \left(\frac{1}{a} \partial_\phi \right) + \phi - i \frac{\partial}{\partial T_\Lambda} \right] \Psi(a, \phi; T_\Lambda) = 0. \quad (62)$$

This is indeed a Schrödinger-like equation with Hamiltonian given by $\hat{C}_{\text{JT}}$,

$$i \frac{\partial}{\partial T_\Lambda} \Psi(a, \phi; T_\Lambda) = \hat{C}_{\text{JT}} \Psi(a, \phi; T_\Lambda). \quad (63)$$

Notice that, unlike the WdW equation in JT gravity, we now have a time-dependent wave function $\Psi(a, \phi; T_\Lambda)$. To

obtain this wave function, we need to start by determining the time-independent $\psi(a, \phi, \Lambda)$ [or $\Psi(a, \phi, \Lambda) = \mathcal{A}(\Lambda) \psi(a, \phi, \Lambda)$ since it also solves the WdW equation]. This has been solved in the previous section due to the fact that the time-independent solution satisfies the WdW equation of KSY gravity. We thus need to be in the $p_{T_\Lambda} = \Lambda$ representation, that is writing an eigenfunction equation

$$\begin{aligned} \hat{C}_{\text{JT}} \Psi(a, \phi, \Lambda) &= \left[\partial_a \left(\frac{1}{a} \partial_\phi \right) + \phi \right] \Psi(a, \phi, \Lambda) \\ &= -\Lambda \Psi(a, \phi, \Lambda). \end{aligned} \quad (64)$$

This tells us that the energy eigenvalues for the JT gravity Hamiltonian constraint are simply given by the vacuum cosmological constant Λ .

Once again, in the context of HT gravity, Λ is a free parameter that is constant on shell. This allows the possibility of having a superposition of Λ states, something that is not possible for KSY (and JT) gravity alone. Indeed, the time-dependent wave function can be written as a superposition of these wave functions, multiplied by some amplitude. That is, we have solutions to the Schrödinger equation given by

$$\Psi(a, \phi; T_\Lambda) = \int \frac{d\Lambda}{\sqrt{2\pi}} e^{i\Lambda T_\Lambda} \mathcal{A}(\Lambda) \psi(a, \phi, \Lambda). \quad (65)$$

Here, we have used unitary evolution of the wave function $\Psi(a, \phi; T_\Lambda)$, dictated by Schrödinger dynamics. We can now invert this formula to obtain the amplitude $\mathcal{A}(\Lambda)$ in terms of the wave function $\Psi(a, \phi; T_\Lambda)$,

$$\mathcal{A}(\Lambda) = \int da e^{-i\Lambda T_\Lambda} \psi^*(a, \phi, \Lambda) \Psi(a, \phi; T_\Lambda). \quad (66)$$

To define a physical Hilbert space, we introduce the inner product on the space of solutions to the time-dependent wave functions $\Psi(a, \phi; T_\Lambda)$ to be

$$\langle \Psi_1 | \Psi_2 \rangle := \int d\Lambda \mathcal{A}_1^*(\Lambda) \mathcal{A}_2(\Lambda), \quad (67)$$

with norm simply given by the modulus square of the amplitudes,

$$\langle \Psi | \Psi \rangle = \int d\Lambda |\mathcal{A}(\Lambda)|^2. \quad (68)$$

This product is automatically conserved with respect to unimodular time, i.e., unitarity is satisfied, since it is defined in terms of time-independent amplitudes.

To construct the time-independent part of the wave function as a superposition of wave packets, we take the amplitude $\mathcal{A}(\Lambda)$ to be a Gaussian distribution centered at Λ_0 with standard deviation σ_{T_Λ} ,

$$\mathcal{A}(\Lambda) = \frac{1}{(2\pi\sigma_{T_\Lambda}^2)^{\frac{1}{4}}} \exp\left(-\frac{(\Lambda - \Lambda_0)^2}{4\sigma_{T_\Lambda}^2}\right). \quad (69)$$

This assumption is justified by the fact that δ function amplitudes do not produce wave packet solutions to the Schrödinger equation and, moreover, they have infinite norm, which is a undesired feature. Given that we have unitarity, it is reasonable to define physical states as any state derived from a normalized Gaussian amplitude, which satisfies $\langle\P|\Psi\rangle = 1$.

Notably, in JT gravity, the inner product is defined by the DeWitt inner product [69,75,77,78]. This resembles the Klein-Gordon inner product, as the WdW equation can be viewed as a Klein-Gordon equation in the light cone coordinates (a, ϕ) . This similarity underscores the absence of standard unitary evolution for wave functions in JT gravity, precluding the definition of a conventional Schrödinger inner product. However, we highlight that such a unitary evolution becomes feasible when unimodular time is introduced, enabling the definition of the physical inner product, as given by (67).

This construction mirrors the four-dimensional scenarios explored in [22,23,27,28], where toy models of the universe involving matter, radiation, and dark energy have been investigated. In many cases, the presence of multiple fluid components leads to distinct relational time variables, each prevailing in its respective cosmological epoch. Building unimodular wave packets in this work is directly analogous to the present work, arising as a fundamental consequence of the underlying Schrödinger equation.

IV. TOWARD UNITARY QUANTUM COSMOLOGY AND TOPOLOGY CHANGE IN HT₂

Consider the contracting and expanding branches of the HH wave function solution in the KSY model, separately,

$$\Psi_{\pm}(a, \phi, \Lambda) = \alpha_{\pm} \mathcal{A}(\Lambda) \psi_{\pm}(a, \phi, \Lambda). \quad (70)$$

Then, using the normalized Gaussian amplitude (69) and integrating over Λ in (65), we find the asymptotic solution in the limit $a\phi \gg 1$,

$$\Psi_{\pm}(a, \phi; T_{\Lambda}) \approx \left(\frac{8\sigma_{T_{\Lambda}}^2}{\pi}\right)^{\frac{1}{4}} e^{-\sigma_{T_{\Lambda}}^2 (T_{\Lambda} \pm a)^2 + i\Lambda_0 (T_{\Lambda} \pm a)} \psi_{\pm}(a, \phi), \quad (71)$$

where $\psi_{\pm}(a, \phi) = e^{\pm ia\phi}$ is the HH solution for MSS JT gravity. It can be further checked that this gives the correct norm as expected, i.e., from the Plancherel theorem we get

$$\langle\P|\Psi\rangle = \int_{-\infty}^{\infty} d\Lambda |\mathcal{A}(\Lambda)|^2 \quad (72)$$

$$= \int_0^{\infty} da |\Psi_{\pm}(a, \phi; T_{\Lambda})|^2 = 1. \quad (73)$$

Therefore, it is quite straightforward to see that the probability density given by $|\Psi_{\pm}(a, \phi; T_{\Lambda})|^2$ is found to follow a Gaussian distribution for a contracting and expanding closed universe, with its peak corresponding to the semiclassical WKB value.

However, we are also interested in examining linear combinations of the two branches, akin to (51). That is, consider the following time-dependent wave function solution

$$\Psi(a, \phi; T_{\Lambda}) = \int \frac{d\Lambda}{\sqrt{4\pi}} \mathcal{A}(\Lambda) e^{i\Lambda T_{\Lambda}} (\psi_{-} + i\psi_{+}). \quad (74)$$

Inverting the wave function gives an inversion formula for the amplitudes that now reads

$$\mathcal{A}(\Lambda) = \int \frac{da}{\sqrt{4\pi}} e^{-i\Lambda T_{\Lambda}} \Psi(a, \phi; T_{\Lambda}) (\psi_{+} - i\psi_{-}). \quad (75)$$

Consequently, we anticipate the emergence of interference terms in the probability density $|\Psi(a, \phi; T_{\Lambda})|^2$, arising from the cross terms $\psi_{\pm}^*(a, \phi) \psi_{\mp}(a, \phi)$. Indeed, we define the probability density as

$$\mathcal{P}(a, \phi, T_{\Lambda}) = |\Psi(a, \phi; T_{\Lambda})|^2, \quad (76)$$

where $\Psi(a, \phi; T_{\Lambda})$ is obtained by substituting (51) in (65).

This is plotted in Fig. 2. The interference between the two quantum states becomes noticeable when $|T_{\Lambda}| \sim \sigma_{T_{\Lambda}}$; however, at later times, as $|T_{\Lambda}| \gg \sigma_{T_{\Lambda}}$, the states separate and evolve into independent, sharply peaked semiclassical WKB states that continue to narrow as unimodular time progresses. Thus, significant physics occurs around $|T_{\Lambda}| \sim \sigma_{T_{\Lambda}}$ (with $T_{\Lambda} = 0$ in the plots). This probability density integrates to 1, as it should according to (72) for all times, confirming unitarity in the theory.

This is analogous to the four-dimensional case involving Hartle-Hawking wave packets [27], where interference arises between incident and reflected waves associated with contracting and expanding de Sitter universes. However, key differences should be emphasized. First, our construction begins from a metric space formulation, more natural in two dimensions, rather than from a connection-based approach. Second, we analyze the evolution of a instead of a^2 , which in four dimensions corresponds canonically to the connection variable, a correspondence absent in two dimensions.

Classically, the universe exhibits a bouncing de Sitter geometry in unimodular time, as shown in Fig. 3. To see why, consider the global de Sitter metric with scale factor $a(t) = \cosh(t)$. Imposing the on-shell unimodular condition

$$\sqrt{-g} = \dot{T}^0 \Rightarrow a(t) = \dot{T}_{\Lambda}, \quad (77)$$

we find that on shell, unimodular time is




FIG. 2. Probability $\mathcal{P}(a, \phi, T_\Lambda)$, according to Eq. (76), for times $T_\Lambda = 0, 100, 120$, and 200 , with $\sigma_T = 0.01$ and $\phi = 100$. For $|T_\Lambda| \sim \sigma_T$ the probability has many oscillations inside the envelope, but these disappear as $|T_\Lambda|$ increases, as illustrated. We see that, for increasing $|T_\Lambda| \gg \sigma_T$, the width of the probability decreases; that is, the uncertainty in a decreases.




FIG. 3. Schematic drawing for the on-shell unitary time evolution of global de Sitter geometry in unimodular time.

$$T_\Lambda(t) = \int_{\Sigma_t} dt a(t) = \sinh(t), \quad \forall t \in \mathbb{R}. \quad (78)$$

Since T_Λ explicitly depends on the coordinate time t , we observe that $T_\Lambda = 0$ at $t = 0$. Thus, for vanishing unimodular time, the scale factor reaches the bouncing point $a = 1$.

At early unimodular times $|T_\Lambda| \sim \sigma_{T_\Lambda}$, due to quantum interference of the contracting incident and expanding reflected waves, the classical de Sitter geometry can undergo quantum deformations. In particular, the universe may evolve toward the singular point $a = 0$, enabling a transition from a circular spatial topology to a point and back again. One can interpret this as a crunching universe, followed by the creation of an expanding universe at $a = 0$ with a bang. Therefore in this scenario, the universe undergoes a topology change process.

A preliminary interpretation of the results goes as follows: we extend JT gravity theory by allowing the vacuum cosmological constant Λ to vary off shell through the introduction of unimodular time. This modification allows the wave function to be in different energy eigenstates of Λ within the quantum theory. Consequently, quantum deformations of the global de Sitter geometry at early unimodular times can be viewed as the universe occupying various Λ -dependent energy levels.

Additionally, we find that the Hartle-Hawking “no-boundary proposal” does not appear to be applicable in this context. The no-boundary proposal might be relevant

in scenarios where the universe is exclusively expanding or contracting, with the wave function peaking at all unimodular times and without the presence of any quantum interference effects. However, these statements should be treated with caution, as a comprehensive path-integral formulation in this framework is needed to better illuminate these situations. Although these features warrant further investigation, a detailed discussion is beyond the scope of this paper.

V. PAGE-WOOTTERS QUANTIZATION OF HT₂ GRAVITY

We have seen that two-dimensional Henneaux-Teitelboim gravity offers a consistent classical resolution to the problem of time by introducing unimodular time, which is canonically conjugate to the vacuum cosmological constant Λ . Additionally, we have developed a unitary quantum cosmological description of two-dimensional de Sitter spacetime, incorporating both the time-dependent and time-independent Hartle-Hawking wave functions. A key objective is to give a proper quantization scheme that aligns with the results presented in the previous section. Here, we achieve this by integrating unimodular gravity with the Page-Wootters formalism of quantum mechanics.

A. Clocks, systems, and the universe in the PW formalism

In quantum mechanics, time T appears as a classical parameter in the Schrödinger equation, leading to time-dependent quantum states $|\psi(T)\rangle$, subject to unitary evolution and probability conservation. This notion of a “physical time” seems to be missing within quantum gravity, where quantum states are annihilated by constraint operators, indicating the absence of both unitary evolution and probability conservation, as the quantum states appear to be “frozen in time.” It is thus of great importance to be able to give a fully quantum description of time.

The starting point of PW formalism [35–39] is to consider time as an external quantum degree of freedom by assigning to it its own “clock” Hilbert space $\mathcal{H}_C$. One can then consider clock states $|T\rangle$ in $\mathcal{H}_C$, defined via the time operator in its spectral decomposition

$$\hat{T} = \int dT T |T\rangle \langle T|. \quad (79)$$

Throughout the paper, we shall assume that time is described by idealized clocks: the clock states are taken to satisfy the orthogonality condition

$$\langle T|T'\rangle = \delta(T - T'). \quad (80)$$

With that being said, the clock Hilbert space can be viewed as the space of square-integrable function $\mathcal{H}_C \cong L^2(\mathbb{R})$.

Since time has an associated canonically conjugate momenta H_C and satisfies the commutation relation on $\mathcal{H}_C$

$$[\hat{T}, \hat{H}_C] = i\hat{1}_C, \quad (81)$$

we have the clock Hamiltonian operator $\hat{H}_C$ defined as

$$\hat{H}_C = \int d\mathcal{E}_C \mathcal{E}_C |\mathcal{E}_C\rangle \langle \mathcal{E}_C|. \quad (82)$$

Here, $|\mathcal{E}_C\rangle$ are clock energy eigenstates that live in the clock Hilbert space, which also satisfies an orthogonality relation. Furthermore, the time and clock Hamiltonian eigenstates are Fourier duals of one another,

$$|\mathcal{E}_C\rangle = \int \frac{dT}{\sqrt{2\pi}} e^{i\mathcal{E}_C T} |T\rangle. \quad (83)$$

The two eigenstates are thus related by exponentiation

$$\langle T|\mathcal{E}_C\rangle = \frac{e^{i\mathcal{E}_C T}}{\sqrt{2\pi}}. \quad (84)$$

As a consequence, the clock Hamiltonian operator $\hat{H}_C$ in the time T representation satisfies

$$\hat{H}_C |T\rangle = -i \frac{d}{dT} |T\rangle. \quad (85)$$

This will play a pivotal role in defining the Schrödinger equation below.

To further define a quantum state of an explicit physical system, we introduce the “system” Hilbert space $\mathcal{H}_S$. Quantum states associated with the system are $|\psi\rangle$ and an associated Hamiltonian operator $\hat{H}_S$ acting on it

$$\hat{H}_S |\psi\rangle = E_S |\psi\rangle, \quad (86)$$

where E_S is the energy (eigenvalue) of the system. The system has an associated phase space with generalized coordinates $Q = \{q\}$ and conjugate generalized momenta $P_Q = \{p_q\}$. When promoted to linear operators acting on $\mathcal{H}_S$, they satisfy the spectral decomposition,

$$\hat{Q} |Q\rangle = Q |Q\rangle, \quad (87)$$

$$\hat{P}_Q |Q\rangle = -i\partial_Q |Q\rangle, \quad (88)$$

where $|Q\rangle$ is an eigenstate of the system coordinate operator. Any state $|\psi\rangle \in \mathcal{H}_S$ can be written as a linear superposition of $|Q\rangle$ states. The specific form of Q depends on the physical system under consideration. This will be explicitly given when considering the MSS JT gravity model in the next section.

To define correlations among the clock and the system, we construct the so-called “universe” Hilbert space $\mathcal{H} = \mathcal{H}_C \otimes \mathcal{H}_S$. This is described by the noninteracting composite quantum system formed by the clock and system Hilbert spaces. The associated universe quantum states are then denoted by $|\Psi\rangle$. The full Hamiltonian of the universe is thus given by

$$\hat{H} = \hat{H}_C \otimes \hat{1}_S + \hat{1}_C \otimes \hat{H}_S, \quad (89)$$

where the resolution of the identity on the respective subspaces are

$$\hat{1}_C := \int dT |T\rangle\langle T| = \int d\Lambda |\Lambda\rangle\langle \Lambda|, \quad (90)$$

$$\hat{1}_S := \int dQ |Q\rangle\langle Q|. \quad (91)$$

So far, we have given the kinematical Hilbert space. To define the physical Hilbert space of the universe, we identify physical states on $\mathcal{H}$ as those annihilated by the universe Hamiltonian,

$$\hat{H}|\Psi\rangle = 0. \quad (92)$$

This is akin to the Wheeler-DeWitt equation of the universe. Note that since the universe Hamiltonian annihilates physical states of the universe, they are not subject to unitary time evolution. This is what is meant by frozen (or static) quantum states.

In order to get time-dependent quantum states subject to the standard Schrödinger equation, we demand the system to be time dependent. In PW formalism, this is achieved by projecting the clock to the system: the conditioned quantum states denoted $|\Psi(T)\rangle \in \mathcal{H}_{S|T}$ are then defined as follows:

$$|\Psi(T)\rangle := \langle T| \otimes \hat{1}_S |\Psi\rangle. \quad (93)$$

These states are defined on a subspace of the system Hilbert space and are subject to time evolution. This can be seen by taking the Wheeler-DeWitt equation of the universe and projecting it to $\mathcal{H}_{S|C}$,

$$(\langle T| \otimes \hat{1}_S) \hat{H} |\Psi\rangle = 0 \Rightarrow i \frac{d}{dT} |\Psi(T)\rangle = \hat{H}_S |\Psi(T)\rangle. \quad (94)$$

We therefore get unitary time evolution of system quantum states according to

$$|\Psi(T)\rangle = e^{-iT\hat{H}_S} |\Psi(0)\rangle, \quad (95)$$

where $|\Psi(0)\rangle \equiv |\Psi(T=0)\rangle$ satisfies the time-independent Schrödinger equation

$$\hat{H}_S |\Psi(0)\rangle = E_S |\Psi(0)\rangle. \quad (96)$$

It can be seen from (83) that inverting the Fourier transform gives

$$|0\rangle = \int \frac{d\mathcal{E}_C}{\sqrt{2\pi}} |\mathcal{E}_C\rangle. \quad (97)$$

We can therefore write the time-independent quantum state $|\Psi(0)\rangle$ in terms of $|\Psi(H_C)\rangle$, that is,

$$|\Psi(0)\rangle = \int \frac{d\mathcal{E}_C}{\sqrt{2\pi}} |\Psi(\mathcal{E}_C)\rangle. \quad (98)$$

We can also condition states with the clock Hamiltonian eigenkets $|\mathcal{E}_C\rangle$ and obtain the “dual” Schrödinger equation

$$(\langle \mathcal{E}_C| \otimes \hat{1}_S) \hat{H} |\Psi\rangle = 0 \Rightarrow \hat{H}_S |\Psi(\mathcal{E}_C)\rangle = -\mathcal{E}_C |\Psi(\mathcal{E}_C)\rangle. \quad (99)$$

It can be read off that the energy of the system is given by $E_S = -\mathcal{E}_C$, in agreement with the fact that the classical Hamiltonian of the universe vanishes.

To obtain wave functions of the form $\Psi(Q; T)$, we perform the bracket operation $\langle Q| \Psi(T)\rangle$. The time-dependent wave function solution can thus be written as a superposition of energy wave functions,

$$\Psi(Q; T) = \int \frac{d\mathcal{E}_C}{\sqrt{2\pi}} e^{i\mathcal{E}_C T} \Psi(Q, \mathcal{E}_C). \quad (100)$$

One can also assume that the time-independent solutions are given via a (possibly Gaussian) amplitude $\mathcal{A}(\mathcal{E}_C)$, such that $\Psi(Q, \mathcal{E}_C) = \mathcal{A}(\mathcal{E}_C) \psi(Q, \mathcal{E}_C)$,

$$\Psi(Q; T) = \int \frac{d\mathcal{E}_C}{\sqrt{2\pi}} e^{i\mathcal{E}_C T} \mathcal{A}(\mathcal{E}_C) \psi(Q, \mathcal{E}_C). \quad (101)$$

These are also solved by the Schrödinger equation. In this case, we obtain wave packets $\Psi(Q; T)$, whereas for $\psi(Q, \mathcal{E}_C)$, these are not necessarily wave packets (for instance, it can be a plane wave solution).

We would like to end this section by mentioning the inner product on $\mathcal{H}$. Since we want normalizable wave functions on the system that are subject to probability conservation, we consider the physical inner product on the universe quantum states to be given by

$$\langle \Psi_1 | \Psi_2 \rangle := \langle \Psi_1 | (\langle T| \otimes \hat{1}_S) | \Psi_2 \rangle. \quad (102)$$

Using the fact that the universe quantum state can be written as an entangled state of the clock and system states,

$$|\Psi\rangle = \int dT |T\rangle |\Psi(T)\rangle, \quad (103)$$

the physical inner product is written as

$$\begin{aligned} \langle \Psi_1 | \Psi_2 \rangle &= \langle \Psi_1(T) | \Psi_2(T) \rangle \\ &= \int d\mathcal{E}_C \mathcal{A}_1^*(\mathcal{E}_C) \mathcal{A}_2(\mathcal{E}_C). \end{aligned} \quad (104)$$

Thus, the physical norm is simply

$$\langle \Psi | \Psi \rangle = \int d\mathcal{E}_C |\mathcal{A}(\mathcal{E}_C)|^2. \quad (105)$$

One can also write this as an integral over Q and obtain an expression for the probability density

$$\mathcal{P}(Q, T) = |\Psi(Q; T)|^2. \quad (106)$$

Indeed, from (105), if the amplitudes are normalized Gaussian distributions, then it is clear that the norm is 1 and thus all the probabilities add up to 1.

B. PW quantization of HT₂ minisuperspace

With the PW formalism established, we can now consider its application to two-dimensional Henneaux-Teitelboim gravity. For that, we first define the universe Hilbert space in terms of correlations between the clock and system subspaces. The universe being de Sitter HT₂ gravity in minisuperspace, the system is going to be JT gravity and the clock is given by unimodular time T_Λ .

The Hilbert space of HT₂ is

$$\mathcal{H}_{\text{HT}_2} = \mathcal{H}_{T_\Lambda} \otimes \mathcal{H}_{\text{JT}}. \quad (107)$$

In the unimodular time subspace $\mathcal{H}_{T_\Lambda}$, time is an operator that satisfies $\hat{T}_\Lambda |T_\Lambda\rangle = T_\Lambda |T_\Lambda\rangle$, where unimodular time eigenstates are orthogonal, realizing idealized clocks. Similarly, the clock Hamiltonian is given by the canonically conjugate momentum of unimodular time $\hat{p}_{T_\Lambda} = \hat{\Lambda}$ that satisfies $\hat{\Lambda} |\Lambda\rangle = \Lambda |\Lambda\rangle$, with orthogonal Λ eigenstates. One can therefore write (103) as

$$|\Psi\rangle = \int dT_\Lambda |T_\Lambda\rangle |\Psi(T_\Lambda)\rangle = \int d\Lambda |\Lambda\rangle |\Psi(\Lambda)\rangle, \quad (108)$$

where we have the conditional states on the JT gravity Hilbert subspace,

$$|\Psi(T_\Lambda)\rangle = \langle T_\Lambda | \otimes \hat{1}_{\text{JT}} |\Psi\rangle, \quad (109)$$

$$|\Psi(\Lambda)\rangle = \langle \Lambda | \otimes \hat{1}_{\text{JT}} |\Psi\rangle. \quad (110)$$

The relation (108) shows that $|\Psi\rangle$ provides a complete description of the temporal evolution of the JT gravity system by representing it in terms of correlations between the latter and the degree of freedom of the unimodular time.

From the resolution of the identity

$$\hat{1}_{T_\Lambda} = \int dT_\Lambda |T_\Lambda\rangle \langle T_\Lambda| = \int d\Lambda |\Lambda\rangle \langle \Lambda|, \quad (111)$$

and the fact that $\langle T_\Lambda | \Lambda \rangle = \frac{1}{\sqrt{2\pi}} e^{i\Lambda T_\Lambda}$, the Fourier duality expression (83) simply becomes

$$|\Lambda\rangle = \int \frac{dT_\Lambda}{\sqrt{2\pi}} e^{i\Lambda T_\Lambda} |T_\Lambda\rangle. \quad (112)$$

Indeed, at time $T_\Lambda = 0$, we have that

$$|0\rangle = \int \frac{d\Lambda}{\sqrt{2\pi}} |\Lambda\rangle. \quad (113)$$

Finally, the action of $\hat{\Lambda}$ on unimodular time eigenstates is given by the derivative expression

$$\hat{\Lambda} |T_\Lambda\rangle = -i \frac{d}{dT_\Lambda} |T_\Lambda\rangle. \quad (114)$$

Going back to the Hilbert space of JT gravity, we first define the generalized coordinates as the coordinates on the minisuperspace: this is given by the scale factor $a(t)$ and the dilaton $\phi(t)$. Thus $Q = \{a, \phi\}$, with associated momenta $P_Q = \{p_a, p_\phi\}$. We define on $\mathcal{H}_{\text{JT}}$ the following resolution of the identity:

$$\hat{1}_{\text{JT}} := \int da |a, \phi\rangle \langle a, \phi|, \quad (115)$$

where we take constant- ϕ slices (corresponding to constant-coordinate time t slice). Indeed, these eigenstates satisfy the orthogonality relation at equal $\phi = \phi'$,

$$\langle a, \phi | a', \phi \rangle = \delta(a - a'). \quad (116)$$

The choice of orthogonality is motivated from the fact that we care about the size of each slice (given by the scale factor). We thus look for differently sized universes for a given constant- ϕ slice.

Promoting the phase-space variables to operators acting on $\mathcal{H}_{\text{JT}}$, we have the following eigenstate equations of relevance:

$$\hat{a} |a, \phi\rangle = a |a, \phi\rangle, \quad \hat{p}_a |a, \phi\rangle = -i \partial_a |a, \phi\rangle, \quad (117)$$

$$\hat{\phi} |a, \phi\rangle = \phi |a, \phi\rangle, \quad \hat{p}_\phi |a, \phi\rangle = -i \partial_\phi |a, \phi\rangle. \quad (118)$$

States $|\psi\rangle \in \mathcal{H}_{\text{JT}}$ can be written as a superposition of eigenstates, provided the state is subject to Hamiltonian dynamics.

This is the case when we take the full Hilbert space of Henneaux-Teitelboim gravity. One can define the

Hamiltonian constraint and promoting it to an operator $\hat{C}$, annihilating physical states $|\Psi\rangle$,

$$\hat{C}|\Psi\rangle = 0, \quad (119)$$

where the Hamiltonian operator of the universe is written in terms of $\hat{C}_{JT}$ (the JT gravity Hamiltonian) (59) and the unimodular time Hamiltonian (55). For that, we take the general form of the Hamiltonian of the universe (89) and write it as

$$\hat{C} = \hat{\Lambda} \otimes \hat{1}_{JT} + \hat{1}_{T_\Lambda} \otimes \hat{C}_{JT}. \quad (120)$$

The WdW equation can be conditioned to be that of the Schrödinger equation by projecting the universe states to states living in JT gravity Hilbert space,

$$(\langle T_\Lambda | \otimes \hat{1}_{JT}) \hat{C} |\Psi\rangle = 0 \Rightarrow i \frac{d}{dT_\Lambda} |\Psi(T_\Lambda)\rangle = \hat{C}_{JT} |\Psi(T_\Lambda)\rangle. \quad (121)$$

The quantum states $|\Psi(T_\Lambda)\rangle$ are thus subject to unitary time evolution in unimodular time

$$|\Psi(T_\Lambda)\rangle = e^{-iT_\Lambda \hat{C}_{JT}} |\Psi(0)\rangle. \quad (122)$$

Using (113), we have a relation between the time-dependent and time-independent JT gravity quantum states,

$$\begin{aligned} |\Psi(T_\Lambda)\rangle &= \int \frac{d\Lambda}{\sqrt{2\pi}} e^{-iT_\Lambda \hat{C}_{JT}} |\Psi(\Lambda)\rangle \\ &= \int \frac{d\Lambda}{\sqrt{2\pi}} e^{i\Lambda T_\Lambda} |\Psi(\Lambda)\rangle, \end{aligned} \quad (123)$$

where we used the dual Schrödinger equation (99),

$$\hat{C}_{JT} |\Psi(\Lambda)\rangle = -\Lambda |\Psi(\Lambda)\rangle \Rightarrow E_{JT} = -\Lambda. \quad (124)$$

Assuming a Gaussian state for the energy eigenstates written as $|\Psi(\Lambda)\rangle = \mathcal{A}(\Lambda) |\psi(\Lambda)\rangle$, we have recovered a quantum state version of (65),

$$|\Psi(T_\Lambda)\rangle = \int \frac{d\Lambda}{\sqrt{2\pi}} e^{i\Lambda T_\Lambda} \mathcal{A}(\Lambda) |\psi(\Lambda)\rangle. \quad (125)$$

Note that the wave function expression (65) is recovered upon performing $\langle a, \phi | \Psi(T_\Lambda) \rangle = \Psi(a, \phi; T_\Lambda)$.

The final thing to check is whether the physical inner products match in both approaches. Indeed, using the PW inner product given in (102), we find that

$$\begin{aligned} \langle \Psi_1 | \Psi_2 \rangle &= \langle \Psi_1(T_\Lambda) | \Psi_2(T_\Lambda) \rangle \\ &= \int d\Lambda \mathcal{A}_1^*(\Lambda) \mathcal{A}_2(\Lambda) \equiv \langle \Psi_1 | \Psi_2 \rangle. \end{aligned} \quad (126)$$

This matches the inner product defined in (67) and the norm of PW also agrees,

$$\langle \Psi | \Psi \rangle \equiv \langle \Psi_1 | \Psi_2 \rangle = \int d\Lambda |\mathcal{A}(\Lambda)|^2, \quad (127)$$

which is unity for normalized Gaussian amplitudes.

Some comments are in order, especially regarding quantum interference effects within the formalism. Indeed, as discussed in the previous section, the WdW equation solves both for the expanding and contracting wave functions. That is, there is the freedom of considering their quantum superposition. In that case, we expect the emergence of interference terms at the level of probability densities. This is translated in the PW formalism by first defining the time-independent states on $\mathcal{H}_{JT}$ to be

$$|\psi(\Lambda)\rangle = \frac{|\psi_-(\Lambda)\rangle + i|\psi_+(\Lambda)\rangle}{\sqrt{2}}. \quad (128)$$

Therefore, the time-dependent states are now written as

$$|\Psi(T_\Lambda)\rangle = \int \frac{d\Lambda}{\sqrt{4\pi}} e^{i\Lambda T_\Lambda} \mathcal{A}(\Lambda) (|\psi_-(\Lambda)\rangle + i|\psi_+(\Lambda)\rangle). \quad (129)$$

The inner product is known to be time independent, hence respecting unitarity; that is,

$$\langle \Psi_1(T_\Lambda) | \Psi_2(T_\Lambda) \rangle = \langle \Psi_1(0) | \Psi_2(0) \rangle. \quad (130)$$

The norm is therefore also conserved and so is the total probability. The interference terms arise at the level of the probability amplitude, which are obtained from

$$\begin{aligned} \langle \Psi(0) | \Psi(0) \rangle &= \int_0^\infty da \langle \Psi(0) | a, \phi \rangle \langle a, \phi | \Psi(0) \rangle \\ &= \int_0^\infty da |\Psi(a, \phi, \Lambda)|^2. \end{aligned} \quad (131)$$

This is normalizable if the amplitudes $\mathcal{A}(\Lambda)$ are Gaussian amplitudes.

VI. DISCUSSIONS

The common theme of this work is that a gauge-invariant formulation of Jackiw-Teitelboim gravity with a vacuum cosmological constant Λ naturally emerges as a two-dimensional Henneaux-Teitelboim gravity theory.

By gauging the isometry group of two-dimensional spacetime and introducing an Abelian U(1) central

extension, one obtains HT gravity, which shares the same on-shell dynamics—and therefore the same physical content—as JT gravity. In this framework, Λ is promoted to an off-shell variable that becomes fixed on shell. Furthermore, the unimodular condition introduces a natural physical time called unimodular time that is canonically conjugate to Λ . Moreover, the flat-space analog of this formulation, known as CJ (or Weyl-rescaled CGHS) gravity, admits a similar central extension and retains the core features of “flat” HT gravity (or the $\widehat{\text{CGHS}}$ model).

With HT_2 gravity in hand, we illustrate its utility by performing a minisuperspace reduction in a two-dimensional quantum cosmology framework. In particular, introducing unimodular time transforms the Wheeler-DeWitt equation into a Schrödinger-like equation, thereby ensuring unitary evolution of quantum states. Consequently, the inner product is well defined, leading to a consistent quantum-mechanical description of two-dimensional de Sitter gravity. By solving the Schrödinger equation, we obtain explicit normalizable and unitary time-dependent wave functions for the global dS_2 geometry, represented by expanding and contracting branches described through the unimodular Hartle-Hawking wave packets.

Analyzing the probability density shows that, at different unimodular times T_Λ , the scale factor a follows a probability distribution, thus providing a quantum account of the Universe’s expansion and contraction. Notably, at early unimodular times, quantum interference between the expanding and contracting branches dominates. In particular, at $T_\Lambda = 0$, this interference persists beyond the classical region, leading to the possibility of the global de Sitter geometry to be deformed and undergo topology change at the singular point $a = 0$.

The study of quantum correlations between external physical time (or clocks in terms of unimodular time) and JT gravity leads to a Page-Wootters formulation of HT_2 quantum gravity. Promoting the HT gravity Hamiltonian constraint to a linear operator that annihilates physical

states yields the Schrödinger equation by T_Λ -state conditioning. This gives unitary evolution and probability conservation of quantum states, as expected.

A natural next step is to investigate how HT_2 gravity behaves within a holographic framework, especially through its path-integral formulation and potential connections to random matrix theory. Studying these aspects could uncover new insights into how HT_2 gravity fits into broader holographic paradigms. Another promising avenue involves exploring topology change involving higher topologies and two-dimensional flat holography, given the extensive existing literature on flat holography in two dimensions [54–57, 59–61]. Establishing concrete links between the HT_2 formalism and holographic approaches may not only deepen our understanding of low-dimensional quantum gravity, but also offer novel perspectives on the role of topology in holographic setups. Finally, it would be intriguing to explore whether in higher dimensions (such as three and four dimensions), HT gravity theories naturally emerge from a gauge-theoretic formulation akin to the approach developed in this work.

ACKNOWLEDGMENTS

We would like to thank A. Blommaert, A. S. Garrido, O. Gould, F.M. Haehl, R. Isichei, J. Magueijo, I. Matthaikakis, B. Muntz, B. Oguz, P. Saffin, and B. Tekin for helpful discussions and comments. We also thank A. Roy for assistance with the plots. This work was supported by FCT Grant No. 2021.05694.BD (B. A.), UK Research and Innovation (UKRI) under the UK government’s Horizon Europe funding Guarantee EP/X030334/1 (A.E. funded by F.M.H.), and Royal Society Dorothy Hodgkin Fellowship (F.S.R. funded by O. G.).

DATA AVAILABILITY

No data were created or analyzed in this study.

-
- [1] K. V. Kuchar, *Phys. Rev. D* **43**, 3332 (1991).
 - [2] L. Bombelli, W. E. Couch, and R. J. Torrence, *Phys. Rev. D* **44**, 2589 (1991).
 - [3] A. Daughton, J. Louko, and R. D. Sorkin, in *Proceedings of the 5th Canadian Conference on General Relativity and Relativistic Astrophysics (5CCGRR)* (1993).
 - [4] R. D. Sorkin, *Int. J. Theor. Phys.* **33**, 523 (1994).
 - [5] A. Daughton, J. Louko, and R. D. Sorkin, *Phys. Rev. D* **58**, 084008 (1998).
 - [6] A. Einstein, Do gravitational fields play an essential part in the structure of the elementary particles of matter? in *The*

- Principle of Relativity*, edited by A. Einstein (Dover Books on Physics, New York, 1952), pp. 189–198.
- [7] S. Weinberg, *Rev. Mod. Phys.* **61**, 1 (1989).
- [8] M. Henneaux and C. Teitelboim, *Phys. Lett. B* **222**, 195 (1989).
- [9] W. G. Unruh, *Phys. Rev. D* **40**, 1048 (1989).
- [10] L. Smolin, *Phys. Rev. D* **80**, 084003 (2009).
- [11] L. Smolin, *Phys. Rev. D* **84**, 044047 (2011).
- [12] N. Kaloper and A. Padilla, *Phys. Rev. Lett.* **112**, 091304 (2014).
- [13] J. M. Fernández Cristóbal, *Ann. Phys. (N.Y.)* **350**, 441 (2014).

- [14] A. Padilla and I. D. Saltas, *Eur. Phys. J. C* **75**, 561 (2015).
- [15] R. Bufalo, M. Oksanen, and A. Tureanu, *Eur. Phys. J. C* **75**, 477 (2015).
- [16] N. Kaloper, A. Padilla, D. Stefanyshyn, and G. Zahariade, *Phys. Rev. Lett.* **116**, 051302 (2016).
- [17] R. Percacci, *Found. Phys.* **48**, 1364 (2018).
- [18] L. Lombriser, *Phys. Lett. B* **797**, 134804 (2019).
- [19] R. Carballo-Rubio, L. J. Garay, and G. García-Moreno, *Classical Quantum Gravity* **39**, 243001 (2022).
- [20] S. Alexander, J. Magueijo, and L. Smolin, *Symmetry* **11**, 1130 (2019).
- [21] J. Magueijo, T. Zlosnik, and S. Speziale, *Phys. Rev. D* **102**, 064006 (2020).
- [22] J. Magueijo, *Phys. Lett. B* **820**, 136487 (2021).
- [23] J. Magueijo, *Phys. Rev. D* **106**, 084021 (2022).
- [24] P. Jiroušek and A. Vikman, *J. Cosmol. Astropart. Phys.* **04** (2019) 004.
- [25] P. Jiroušek, K. Shimada, A. Vikman, and M. Yamaguchi, *J. Cosmol. Astropart. Phys.* **04** (2021) 028.
- [26] A. Vikman, in *Proceedings of the 55th Rencontres de Moriond on Gravitation* (2021).
- [27] B. Alexandre and J. a. Magueijo, *Phys. Rev. D* **107**, 063501 (2023).
- [28] B. Alexandre, R. Isichei, and J. a. Magueijo, *Phys. Rev. D* **108**, 023526 (2023).
- [29] J. Magueijo and R. Isichei, *Phys. Rev. D* **107**, 023526 (2023).
- [30] V. Chandrasekaran, R. Longo, G. Penington, and E. Witten, *J. High Energy Phys.* **02** (2023) 082.
- [31] E. Witten, *Proc. Symp. Pure Math.* **107**, 247 (2024).
- [32] C. Gomez, [arXiv:2302.14747](#).
- [33] C. Gomez, [arXiv:2304.11845](#).
- [34] E. Witten, *J. High Energy Phys.* **03** (2024) 077.
- [35] D. N. Page and W. K. Wootters, *Phys. Rev. D* **27**, 2885 (1983).
- [36] W. K. Wootters, *Int. J. Theor. Phys.* **23**, 701 (1984).
- [37] V. Giovannetti, S. Lloyd, and L. Maccone, *Phys. Rev. D* **92**, 045033 (2015).
- [38] A. R. H. Smith and M. Ahmadi, *Quantum* **3**, 160 (2019).
- [39] P. A. Hohn, A. R. H. Smith, and M. P. E. Lock, *Phys. Rev. D* **104**, 066001 (2021).
- [40] J. De Vuyst, S. Eccles, P. A. Hohn, and J. Kirklin, [arXiv:2405.00114](#).
- [41] J. De Vuyst, S. Eccles, P. A. Hohn, and J. Kirklin, [arXiv:2412.15502](#).
- [42] D. Cangemi and R. Jackiw, *Phys. Rev. Lett.* **69**, 233 (1992).
- [43] R. Jackiw, in *Proceedings of the 19th International Colloquium on Group Theoretical Methods in Physics* (1992).
- [44] D. Cangemi, *Phys. Lett. B* **297**, 261 (1992).
- [45] G. Grignani and G. Nardelli, *Nucl. Phys.* **B412**, 320 (1994).
- [46] D. Cangemi and R. Jackiw, *Phys. Lett. B* **299**, 24 (1993).
- [47] S. K. Kim, K. S. Soh, and J.-H. Yee, *Phys. Rev. D* **47**, 4433 (1993).
- [48] S. K. Kim, K. S. Soh, and J.-H. Yee, *Phys. Lett. B* **300**, 223 (1993).
- [49] D. Cangemi and R. Jackiw, *Ann. Phys. (N.Y.)* **225**, 229 (1993).
- [50] D. Cangemi and G. V. Dunne, *Phys. Rev. D* **48**, 5721 (1993).
- [51] R. Jackiw, in *International Conference on Interface Between Physics and Mathematics (IPM 93) (Part 1: Plenary Session, 6-10 Sep 1993; Part 2: Tutorial Lectures, 13-17 Sep 1993)* (1993), pp. 116–129.
- [52] K. Hammer, P. Jirousek, and A. Vikman, [arXiv:2001.03169](#).
- [53] A. Etkin, J. Magueijo, and F.-S. Rassouli, *Phys. Lett. B* **855**, 138810 (2024).
- [54] H. A. González, D. Grumiller, and J. Salzer, *J. High Energy Phys.* **05** (2018) 083.
- [55] H. R. Afshar, *J. High Energy Phys.* **02** (2020) 126.
- [56] H. Afshar, H. A. González, D. Grumiller, and D. Vassilevich, *Phys. Rev. D* **101**, 086024 (2020).
- [57] V. Godet and C. Marteau, *J. High Energy Phys.* **07** (2021) 138.
- [58] H. Afshar and B. Oblak, *J. High Energy Phys.* **11** (2022) 172.
- [59] A. Kar, L. Lamprou, C. Marteau, and F. Rosso, *J. High Energy Phys.* **03** (2023) 249.
- [60] F. Rosso, *J. High Energy Phys.* **02** (2023) 037.
- [61] H. Afshar and N. Aghamir, *J. High Energy Phys.* **03** (2023) 009.
- [62] C. G. Callan, Jr., S. B. Giddings, J. A. Harvey, and A. Strominger, *Phys. Rev. D* **45**, R1005 (1992).
- [63] V. Godet and C. Marteau, *J. High Energy Phys.* **12** (2020) 020.
- [64] C. Teitelboim, *Phys. Lett. B* **126**, 41 (1983).
- [65] R. Jackiw, *Nucl. Phys.* **B252**, 343 (1985).
- [66] K. Isler and C. A. Trugenberger, *Phys. Rev. Lett.* **63**, 834 (1989).
- [67] A. H. Chamseddine and D. Wyler, *Phys. Lett. B* **228**, 75 (1989).
- [68] A. Kar, L. Lamprou, C. Marteau, and F. Rosso, *Phys. Rev. Lett.* **129**, 201601 (2022).
- [69] J. Maldacena, G. J. Turiaci, and Z. Yang, *J. High Energy Phys.* **01** (2021) 139.
- [70] J. Cotler and K. Jensen, *J. High Energy Phys.* **12** (2021) 089.
- [71] J. Cotler, K. Jensen, and A. Maloney, *J. High Energy Phys.* **06** (2020) 048.
- [72] U. Moitra, S. K. Sake, and S. P. Trivedi, *J. High Energy Phys.* **06** (2022) 138.
- [73] J. Cotler and K. Jensen, *Phys. Rev. Lett.* **131**, 211601 (2023).
- [74] K. K. Nanda, S. K. Sake, and S. P. Trivedi, *J. High Energy Phys.* **02** (2024) 145.
- [75] J. Cotler and K. Jensen, *J. High Energy Phys.* **12** (2024) 016.
- [76] D. Anninos, C. Baracco, and B. Mühlmann, *J. Cosmol. Astropart. Phys.* **10** (2024) 031.
- [77] J. Held and H. Maxfield, [arXiv:2410.14824](#).
- [78] W. Buchmuller, A. Hebecker, and A. Westphal, [arXiv:2412.09211](#).
- [79] M. Honda, H. Matsui, K. Numajiri, and K. Okabayashi, [arXiv:2412.20398](#).
- [80] N. Iizuka and S. K. Sake, [arXiv:2501.02614](#).
- [81] I. Dey, K. K. Nanda, A. Roy, S. K. Sake, and S. P. Trivedi, [arXiv:2501.03148](#).