

Lepton number violation at the LHC in radiative neutrino mass models with leptoquarks

K. S. Babu^{1,*}, Rahool Kumar Barman^{2,†} and Dorival Gonçalves^{1,‡}

¹*Department of Physics, Oklahoma State University, Stillwater, Oklahoma, 74078, USA*

²*Kavli Institute for the Physics and Mathematics of the Universe (WPI), The University of Tokyo Institutes for Advanced Study, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan*



(Received 15 February 2026; accepted 1 September 2026; published 21 September 2026)

We investigate the prospects for observing lepton number violation (LNV) by two units, $|\Delta L| = 2$, at the LHC within the leptoquark variant of the Zee Model, where Majorana neutrino masses arise radiatively at one loop. The model features an $SU(2)_L$ doublet and singlet leptoquarks, whose interactions produce a distinctive same-sign dilepton plus jets signature, $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$. Taking into account current experimental constraints, we identify the dominant production channels for this LNV signal and perform a detailed collider analysis. We find that the HL-LHC can probe leptoquark masses up to $m_{LQ} \sim 1.5$ TeV with this process. Observation of this signal would provide a direct test of LNV and would unambiguously establish the Majorana nature of neutrinos.

DOI: 10.1103/vbbz-98kg

I. INTRODUCTION

Majorana neutrino masses can naturally arise from loop-induced processes, and a wide range of radiative mechanisms has been proposed in the literature at one-loop [1,2], two-loop [3–5], and three-loop levels [6]. For reviews, see Refs. [7,8]. Among the minimal one-loop realizations is the Zee model [1], which, due to the electroweak s -channel production of its charged scalars, offers promising sensitivity to lepton number violating (LNV) signature at the high-luminosity LHC (HL-LHC) [9].

A central theoretical implication of observing LNV at colliders is rooted in the so-called *black-box theorem* for neutrino mass [10]. Originally formulated in the context of neutrinoless double beta decay ($0\nu\beta\beta$) [11], the theorem states that any process that violates lepton number by two units, $|\Delta L| = 2$, necessarily induces a nonzero Majorana mass for neutrinos, irrespective of the underlying mechanism. For a recent evaluation of the quantitative impact of the theorem, see Ref. [12]. The theorem can be generalized to high-energy collider processes [9], establishing that the observation of a $|\Delta L| = 2$ final state at the LHC would likewise constitute direct evidence for the Majorana nature

of neutrinos. Unlike $0\nu\beta\beta$ searches, which only probe effective LNV interactions with two electrons e^-e^- , the LHC can access effective interactions with electrons and muons in multiple combinations: $e^\pm e^\pm$, $\mu^\pm \mu^\pm$, and $e^\pm \mu^\pm$. This greatly broadens the scope of experimental tests of LNV, providing complementary and, in some cases, unique avenues to probe the Majorana nature of neutrinos.

The LNV signal $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ has been extensively studied in several theoretical frameworks. This includes the left-right symmetric model (LRSM) [13–31], type-I seesaw framework [32–44], and more recently type-II seesaw and Zee models [9,45]. The complementarity between neutrinoless double beta decay and LNV searches at the LHC with $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ has also been investigated in simplified model frameworks [46–52]. Experimental searches for the same-sign dilepton plus jets signature have been performed by ATLAS [53] and CMS [54] in the context of the LRSM and the type-I seesaw, respectively. Related new physics signatures have also been studied at future lepton colliders [55–62], offering a complementary perspective to the same-sign dilepton plus jets channel searches at the LHC. Complementary studies have also been carried out at LHCb [63], focusing on LNV signal from B -meson decays.

Motivated by the interplay between neutrino mass generation and collider observables, we investigate the leptoquark analog of the Zee model in which the original scalar fields are replaced by leptoquarks. This variant preserves the radiative origin of Majorana neutrino masses while benefiting from the sizable QCD production of leptoquarks. Within this framework, we investigate the projected HL-LHC sensitivity to the LNV same-sign

*Contact author: babu@okstate.edu

†Contact author: rahoolbarman@physics.iitm.ac.in

‡Contact author: dorival@okstate.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

dilepton plus jets signature, $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$. The model is further constrained by neutrino mass measurements [64], charged lepton flavor violation [8], electroweak precision observables [65], and existing leptoquark searches [66,67]. We evaluate their impact to map out the currently allowed parameter space, before assessing the collider reach for the lepton number violating signature.

The remainder of this paper is organized as follows. In Sec. II, we present the model framework. Constraints from neutrino mass measurements, low energy measurements, electroweak precision observables, and leptoquark searches at the LHC are discussed in Sec. III. In Sec. IV, we analyze the relevant production channels leading to the LNV same-sign dilepton plus jets signature. In Sec. V, we present a detector level collider analysis, where we evaluate the discovery potential at the HL-LHC. We summarize and conclude in Sec. VI.

II. LEPTOQUARK VARIANT OF THE ZEE MODEL

The Leptoquark variant of the Zee model generates small neutrino masses radiatively at the one-loop level [1,68]. It preserves the Standard Model (SM) gauge structure $SU(3)_C \times SU(2)_L \times U(1)_Y$ and extends the SM particle content by introducing a $SU(2)_L$ doublet leptoquark (LQ) $\Omega(3, 2, 1/6) = (\omega^{2/3}, \omega^{-1/3})$ and a singlet LQ $\chi^{-1/3}(3, 1, -1/3)$. The Yukawa Lagrangian for the LQs can be written as

$$\mathcal{L}_{\text{Yuk}} \supset \lambda_{\alpha\beta} L_\alpha^i d_\beta^c \Omega^j \epsilon_{ij} + \lambda'_{\alpha\beta} L_\alpha^i Q_\beta^j \chi^* \epsilon_{ij} + \text{H.c.}, \quad (2.1)$$

where $\{\alpha, \beta\}$ denote the generation indices and $\{i, j\}$ are the $SU(2)_L$ indices. The couplings λ and λ' are the Yukawa coupling constants for the Ω and χ fields. L and Q are, respectively, the left-handed lepton and quark doublets, and d^c represent the conjugate of the right-handed down-type fields.

To ensure proton stability, we assume a global baryon number symmetry under which both Ω and $\chi^{-1/3}$ carry baryon number $-1/3$. In the absence of this symmetry, dangerous baryon number violating operators of the type $QQ\chi$ and $u^c d^c \chi^*$ would be allowed, leading to rapid proton decay. This model can be viewed as a special case of R -parity violating supersymmetric model [69], where the fields Ω and $\chi^{-1/3}$ are identified as the superpartners of the

quark doublet and d^c fields respectively (i.e., $\Omega \sim \tilde{Q}$ and $\chi^{*1/3} \sim \tilde{d}^c$).

The Higgs potential contains nontrivial interaction terms given by

$$V \supset \mu(\Omega^\dagger H \chi + (\Omega^\dagger H \chi)^*) + \lambda_7(H^\dagger \Omega)(\Omega^\dagger H). \quad (2.2)$$

Here, we have made the complex parameter μ real by a field redefinition. This term in the Higgs potential generates mixing between the $\omega^{-1/3}$ and $\chi^{-1/3}$ fields after electroweak symmetry breaking, with the mass-squared matrix given by

$$\mathcal{M}_{\omega^{-1/3}-\chi^{-1/3}}^2 = \begin{pmatrix} M_{\omega^{2/3}}^2 + \frac{\lambda_7}{2} v^2 & \sqrt{2}\mu v \\ \sqrt{2}\mu v & m_\chi^2 \end{pmatrix}, \quad (2.3)$$

where $v = 246$ GeV is the vacuum expectation value of the neutral component of the SM Higgs doublet H , $M_{\omega^{2/3}}^2$ is the physical mass squared of $\omega^{2/3}$, which includes a contribution from a $\lambda_4(H^\dagger H)(\Omega^\dagger \Omega)$ term, and m_χ^2 is the bare mass term plus the contribution from the $\lambda_5(H^\dagger H)(\chi^\dagger \chi)$ term. The mixing between $\omega^{-1/3}$ and $\chi^{-1/3}$ results in the mass eigenstates X_1 and X_2 ,

$$X_1 = \cos \theta_{\text{LQ}} \omega^{-1/3} + \sin \theta_{\text{LQ}} \chi^{-1/3}, \quad (2.4)$$

$$X_2 = -\sin \theta_{\text{LQ}} \omega^{-1/3} + \cos \theta_{\text{LQ}} \chi^{-1/3}. \quad (2.5)$$

The mixing angle θ_{LQ} is given by $\tan 2\theta_{\text{LQ}} = 2\sqrt{2}\mu v / (m_\omega^2 + \lambda_7 v^2 / 2 - m_\chi^2)$. The squared masses of the physical eigenstates X_1 and X_2 are given by

$$M_{X_{1,2}}^2 = \frac{1}{2} \left[M_{\omega^{2/3}}^2 + \frac{\lambda_7}{2} v^2 + m_\chi^2 \mp \sqrt{\left(M_{\omega^{2/3}}^2 + \frac{\lambda_7}{2} v^2 - m_\chi^2 \right)^2 + 8\mu^2 v^2} \right]. \quad (2.6)$$

The Yukawa Lagrangian in Eq. (2.1) can be expressed in terms of the physical mass eigenstates as

$$\begin{aligned} \mathcal{L}_{\text{Yuk}} = & -\lambda_{\alpha\beta} \ell_\alpha d_\beta^c \omega^{2/3} - \lambda'_{\alpha\beta} V_{\text{CKM}}^\dagger \sin \theta_{\text{LQ}} \ell_\alpha u_\beta X_1 - \lambda'_{\alpha\beta} V_{\text{CKM}}^\dagger \cos \theta_{\text{LQ}} \ell_\alpha u_\beta X_2 \\ & + (\lambda_{\alpha\beta} \cos \theta_{\text{LQ}} \nu_\alpha d_\beta^c + \lambda'_{\alpha\beta} \sin \theta_{\text{LQ}} \nu_\alpha d_\beta^c) X_1 - (\lambda_{\alpha\beta} \sin \theta_{\text{LQ}} \nu_\alpha d_\beta^c - \lambda'_{\alpha\beta} \cos \theta_{\text{LQ}} \nu_\alpha d_\beta^c) X_2 + \text{H.c.}, \end{aligned} \quad (2.7)$$

where ℓ and ν are the charged and neutral-lepton components of the lepton doublet, and u is the left-handed up-type quark. The Yukawa terms in Eq. (2.7), together with the LQ mixing induced by

the cubic scalar term in Eq. (2.2), lead to LNV by two units, $|\Delta L| = 2$. This allows neutrino mass generation at one loop as illustrated with the Feynman diagram in Fig. 1.




FIG. 1. Feynman diagram representing neutrino mass generation at one-loop in the Leptoquark variant of the Zee model.

III. CONSTRAINTS

The considered model is sensitive to a range of experimental measurements, including neutrino oscillation data, low energy observables, precision electroweak measurements, and leptoquark searches at the LHC. In this section, we examine the impact of these constraints on the model parameter space and identify the currently allowed region, which will serve as the foundation for our study of LNV signatures at the HL-LHC.

A. Neutrino mass constraints

Neutrino oscillation data place strong constraints on the mass-squared differences [64]

$$\begin{aligned}\Delta m_{21}^2 &= 7.49_{-0.19}^{+0.19} \times 10^{-5} \text{ eV}^2, \\ \Delta m_{32}^2 &= 2.513_{-0.019}^{+0.021} \times 10^{-3} \text{ eV}^2.\end{aligned}\quad (3.1)$$

In addition, kinematic measurements of tritium beta decay, ${}^3\text{H} \rightarrow {}^3\text{He} + e^- + \bar{\nu}_e$, at the KATRIN experiment constrain the effective electron neutrino mass to $m_{\nu_e} < 0.45 \text{ eV}$ at 90% CL [70].¹ Recent results from NuFit Collaboration [64], performing a global analysis of the neutrino data, imply that for normal ordering the sum of neutrino masses must lie within

$$0.058 \text{ eV} \lesssim \sum m_i \lesssim 1.2 \text{ eV} \quad \text{at 95\% CL.} \quad (3.2)$$

In this work, we adopt normal ordering and fix the lightest neutrino mass to $m_1 = 0.04 \text{ eV}$. The remaining masses are then determined from the measured mass-squared differences: $m_2 = \sqrt{m_1^2 + \Delta m_{21}^2}$ and $m_3 = \sqrt{m_2^2 + \Delta m_{32}^2}$. Using the values in Eq. (3.1), this yields $\sum m_i \simeq 0.15 \text{ eV}$.

In the leptoquark variant of the Zee model, the neutrino mass matrix in the flavor basis, M_ν , arises radiatively at one loop and takes the form

¹The parameter m_{ν_e} is defined as $m_{\nu_e}^2 \equiv \sum_i m_i^2 |U_{ei}|^2$, where U is the 3×3 unitary leptonic Pontecorvo–Maki–Nakagawa–Sakata (PMNS) mixing matrix.

$$M_\nu = U^\dagger \text{diag}(m_1, m_2, m_3) U^\dagger = \kappa (\lambda M_d \lambda'^T + \lambda' M_d \lambda^T), \quad (3.3)$$

where the loop-induced coefficient is

$$\kappa = (3/32\pi^2) \sin 2\theta_{\text{LQ}} \ln(m_{X_1}^2/m_{X_2}^2). \quad (3.4)$$

Here, M_d is the down-type quark mass matrix, and m_{X_1} , m_{X_2} denote the masses of the leptoquark fields X_1 and X_2 , respectively. We adopt the standard parametrization for the 3×3 PMNS matrix, U , from Ref. [71] and use the matrix element values from the NuFit-6.0 global analysis [64]. For simplicity, we assume that the Yukawa matrices are related by a constant scaling factor c :

$$\lambda_{ij} = c \lambda'_{ij}, \quad (3.5)$$

which reduces Eq. (3.3) to a relation involving a single unknown matrix. In our analysis, for each choice of leptoquark masses and mixing angle, we solve this equation to determine the elements of the 3×3 Yukawa matrices.

B. Low energy measurements

Various low energy measurements, including rare decays of charged leptons and mesons, place relevant constraints on the parameter space of interest. For instance, leptoquarks can induce charged lepton flavor violating (cLFV) decays of the form $\ell_\alpha \rightarrow \ell_\beta + \gamma$ at one loop, where the γ is radiated either from the leptoquark or the quark lines in the loop. In this class of processes, the contributions mediated by the ω leptoquark are subject to a Glashow–Iliopoulos–Maiani-like cancellation [8,65,72,73], resulting in relatively weak constraints on the λ couplings. The surviving terms in the amplitude are suppressed by a factor of order (m_d^2/m_ω^2) . In contrast, the processes mediated by X_1 and X_2 , owing to their $\chi^{-1/3}$ admixture, can introduce strong constraints on the λ' couplings. The branching ratio for the $\ell_\alpha \rightarrow \ell_\beta + \gamma$ process mediated by X_1 and X_2 is given by

$$\begin{aligned}\mathcal{BR}(\ell_\alpha \rightarrow \ell_\beta + \gamma) &= \frac{9\alpha_{\text{em}} m_{\ell_\alpha}^5}{576(16\pi^2)^2 \Gamma(\ell_\alpha)} |\lambda'_{\beta d} \lambda'^{\star}_{ad}|^2 \\ &\times \left(\frac{\sin^2 \theta_{\text{LQ}}}{m_{X_1}^2} + \frac{\cos^2 \theta_{\text{LQ}}}{m_{X_2}^2} \right)^2.\end{aligned}\quad (3.6)$$

In Table I, we present the resulting constraints on $|\lambda'_{\beta d} \lambda'^{\star}_{ad}|$ for $\mu \rightarrow e\gamma$, $\tau \rightarrow e\gamma$, and $\tau \rightarrow \mu\gamma$, as a function of θ_{LQ} and leptoquark masses m_{X_1} and m_{X_2} .

Another rare cLFV process that probes both λ and λ' couplings is coherent μ – e conversion in the nuclei, $\mu N \rightarrow e N$, which occurs at tree level in these models. The corresponding branching ratio can be expressed as [8,65]

TABLE I. Summary of constraints from low energy measurements on the leptoquark Yukawa couplings λ and λ' as a function of leptoquark masses $\{m_{\omega^{2/3}}, m_{X_1}, m_{X_2}\}$ [8,65,74–86]. In the expression for the $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ constraint, $\hat{\lambda}$ and $\hat{\lambda}'$ are defined as $\hat{\lambda} = \lambda^T U$ and $\hat{\lambda}' = \lambda'^T U$ [85], where U is the PMNS matrix (see Sec. III A for details). The indices $m, n = 1, 2, 3$ appearing in the last line denote the neutrino states in the mass basis, following the convention of [85].

Process	Current limit	Bounds (TeV) ⁻²
$\mu \rightarrow e\gamma$	$\mathcal{BR} < 1.5 \times 10^{-13}$ [74]	$ \lambda'_{ed}\lambda_{\mu d}^* < 8.6 \times 10^{-4} \left(\frac{s_\theta^2}{m_{X_1}^2} + \frac{c_\theta^2}{m_{X_2}^2} \right)^{-1}$
$\tau \rightarrow e\gamma$	$\mathcal{BR} < 3.3 \times 10^{-8}$ [75]	$ \lambda'_{ed}\lambda_{\tau d}^* < 1.6 \left(\frac{s_\theta^2}{m_{X_1}^2} + \frac{c_\theta^2}{m_{X_2}^2} \right)^{-1}$
$\tau \rightarrow \mu\gamma$	$\mathcal{BR} < 4.2 \times 10^{-8}$ [76]	$ \lambda_{\mu d}^*\lambda'_{\tau d} < 1.8 \left(\frac{s_\theta^2}{m_{X_1}^2} + \frac{c_\theta^2}{m_{X_2}^2} \right)^{-1}$
$\mu N \rightarrow eN$		
$^{48}_{22}\text{Ti}$	$\mathcal{BR} < 6.1 \times 10^{-13}$	$\frac{ \lambda_{ed}^*\lambda_{\mu d} }{m_{\omega^{2/3}}^2} + \lambda_{ed}^*\lambda'_{\mu d} \left(\frac{s_\theta^2}{m_{X_1}^2} + \frac{c_\theta^2}{m_{X_2}^2} \right) < 4.30 \times 10^{-6}$
$^{197}_{79}\text{Au}$	$\mathcal{BR} < 7.0 \times 10^{-13}$ [77]	$\frac{ \lambda_{ed}^*\lambda_{\mu d} }{m_{\omega^{2/3}}^2} + \lambda_{ed}^*\lambda'_{\mu d} \left(\frac{s_\theta^2}{m_{X_1}^2} + \frac{c_\theta^2}{m_{X_2}^2} \right) < 4.29 \times 10^{-6}$
$^{208}_{82}\text{Pb}$	$\mathcal{BR} < 4.6 \times 10^{-11}$ [78]	$\frac{ \lambda_{ed}^*\lambda_{\mu d} }{m_{\omega^{2/3}}^2} + \lambda_{ed}^*\lambda'_{\mu d} \left(\frac{s_\theta^2}{m_{X_1}^2} + \frac{c_\theta^2}{m_{X_2}^2} \right) < 3.56 \times 10^{-5}$
$K_L^0 \rightarrow e^+e^-$	$\mathcal{BR} < 8.7^{+5.7}_{-4.1} \times 10^{-12}$ [79]	$ \lambda_{ed}^*\lambda_{es} \left(\frac{1}{m_{\omega^{2/3}}^2} \right) < 1.9 \times 10^{-3}$
$K_L^0 \rightarrow \mu^+\mu^-$	$\mathcal{BR} < (6.84 \pm 0.11) \times 10^{-9}$ [80]	$ \lambda_{\mu d}^*\lambda_{\mu s} \left(\frac{1}{m_{\omega^{2/3}}^2} \right) < 4.7 \times 10^{-5}$
$K_L^0 \rightarrow e^\pm\mu^\mp$	$\mathcal{BR} < 4.7 \times 10^{-12}$ [81]	$ \lambda_{es}\lambda_{\mu d}^* + \lambda_{ed}\lambda_{\mu s}^* \left(\frac{1}{m_{\omega^{2/3}}^2} \right) < 1.9 \times 10^{-5}$
$K^+ \rightarrow \pi^+e^+e^-$	$\mathcal{BR} < (3.00 \pm 0.09) \times 10^{-7}$ [80]	$ \lambda_{ed}^*\lambda_{es} \left(\frac{1}{m_{\omega^{2/3}}^2} \right) < 2.3 \times 10^{-2}$
$K^+ \rightarrow \pi^+\mu^+\mu^-$	$\mathcal{BR} < (9.15 \pm 0.08) \times 10^{-8}$ [82]	$ \lambda_{\mu d}^*\lambda_{\mu s} \left(\frac{1}{m_{\omega^{2/3}}^2} \right) < 1.85 \times 10^{-2}$
$K^+ \rightarrow \pi^+\mu^+e^-$	$\mathcal{BR} < 1.3 \times 10^{-11}$ [83]	$ \lambda_{ed}^*\lambda_{\mu s} \left(\frac{1}{m_{\omega^{2/3}}^2} \right), \lambda_{es}\lambda_{\mu d} ^* \left(\frac{1}{m_{\omega^{2/3}}^2} \right) < 1.9 \times 10^{-4}$
$K^+ \rightarrow \pi^+\nu\bar{\nu}$	$\mathcal{BR} < 9.6^{+1.9}_{-1.8} \times 10^{-11}$ [84]	$\sum_{m \neq n} \left \left(\frac{s_\theta^2}{m_{X_1}^2} + \frac{c_\theta^2}{m_{X_2}^2} \right) \hat{\lambda}'_{dn}\hat{\lambda}'_{sm} - \left(\frac{c_\theta^2}{m_{X_1}^2} + \frac{s_\theta^2}{m_{X_2}^2} \right) \hat{\lambda}'_{sn}\hat{\lambda}'_{dm} \right < 3.1 \times 10^{-4}$

$$\begin{aligned}
&\mathcal{BR}(\mu N \rightarrow eN) \\
&= \frac{|\vec{p}_e \cdot E_e m_\mu^3 G_F^2 \alpha_{\text{em}}^4 Z_{\text{eff}}^4 F_p^2|}{64 \pi^2 Z \Gamma_N} (2A - Z)^2 \\
&\times \left(\frac{|\lambda_{ed}^*\lambda_{\mu d}|}{m_{\omega^{2/3}}^2} + |\lambda_{ed}^*\lambda'_{\mu d}| \left(\frac{\sin^2 \theta_{\text{LQ}}}{m_{X_1}^2} + \frac{\cos^2 \theta_{\text{LQ}}}{m_{X_2}^2} \right) \right)^2, \quad (3.7)
\end{aligned}$$

where $\vec{p}_e$ and E_e denote the momentum and energy of the outgoing electron, Z_{eff} is the effective atomic number, F_p is the nuclear matrix element, and A and Z are the mass number and atomic number of the nucleus, respectively. Γ_N is the rate of capture of muons by the nucleus N . Using the nuclear and model input parameters detailed in Ref. [8], we derive upper limits on the Yukawa couplings from current measurements of $\mathcal{BR}(\mu N \rightarrow eN)$ for $^{48}_{22}\text{Ti}$, $^{197}_{79}\text{Au}$ and $^{208}_{82}\text{Pb}$, as summarized in Table I. Rare kaon decays provide additional probes of the flavor structure of leptoquark interactions, constraining the product of leptoquark

Yukawa couplings involving the first and second generation down-type quarks. The adopted limits are summarized in Table I. We find that $\mu \rightarrow e\gamma$ and $\mu N \rightarrow eN$ processes are the most stringent among the cLFV and meson decay constraints on our parameter space. A detailed overview of these processes and their impact on the LQ variant of the Zee model can be found in Refs. [8,65,86].

C. Electroweak precision constraints

Precision electroweak observables provide additional restrictions via the oblique parameters S , T , and U [8,87,88]. In our model, they take the form

$$\begin{aligned}
S &= \frac{1}{12\pi} \left(\sin^2 \theta_{\text{LQ}} \ln \frac{m_{X_1}^2}{m_{\omega^{2/3}}^2} + \cos^2 \theta_{\text{LQ}} \ln \frac{m_{X_2}^2}{m_{\omega^{2/3}}^2} \right. \\
&\quad \left. + 3 \cos^2 \theta_{\text{LQ}} \sin^2 \theta_{\text{LQ}} \mathcal{G} \left(\frac{m_{X_1}^2}{m_{X_2}^2} \right) \right), \quad (3.8)
\end{aligned}$$

$$T = \frac{3}{8\pi^2\alpha_{\text{em}}v^2} \left(\cos^2\theta_{\text{LQ}}\mathcal{F}(m_{\omega^{2/3}}^2, m_{\tilde{X}_1}^2) + \sin^2\theta_{\text{LQ}}\mathcal{F}(m_{\omega^{2/3}}^2, m_{\tilde{X}_2}^2) - \cos^2\theta_{\text{LQ}}\sin^2\theta_{\text{LQ}}\mathcal{F}(m_{\tilde{X}_1}^2, m_{\tilde{X}_2}^2) \right), \quad (3.9)$$

$$U = \frac{1}{4\pi} \left(\sin^2\theta_{\text{LQ}}\mathcal{G}\left(\frac{m_{\omega^{2/3}}^2}{m_{\tilde{X}_1}^2}\right) + \cos^2\theta_{\text{LQ}}\mathcal{G}\left(\frac{m_{\omega^{2/3}}^2}{m_{\tilde{X}_2}^2}\right) - \cos^2\theta_{\text{LQ}}\sin^2\theta_{\text{LQ}}\mathcal{G}\left(\frac{m_{\tilde{X}_1}^2}{m_{\tilde{X}_2}^2}\right) \right), \quad (3.10)$$

with

$$\mathcal{F}(a, b) = \frac{1}{2}(a + b) - \frac{ab}{a - b} \ln\left(\frac{a}{b}\right), \quad (3.11)$$

$$\mathcal{G}(c) = \frac{c^3 - 3c^2 - 3c + 1}{(c - 1)^3} \ln c - \frac{5c^2 - 22c + 5}{3(c - 1)^2}. \quad (3.12)$$

The current data yield the following 2σ ranges [71]:

$$\begin{aligned} -0.24 < \Delta S < 0.26, \quad -0.23 < \Delta T < 0.25, \\ \text{and} \quad -0.19 < \Delta U < 0.17, \end{aligned} \quad (3.13)$$

which we impose in our parameter scans. While ΔT is controlled by isospin-breaking effects and therefore typically favors small mass splittings among the leptoquark mass eigenstates, ΔS and ΔU depend logarithmically on the leptoquark mass ratios and vanish in the degenerate mass limit. In practice, the most stringent constraint among the three generally arises from ΔT , as we demonstrate explicitly in Sec. III E.

D. LHC bounds

Leptoquark searches at the LHC have been performed in both single and pair production channels [66,89–94]. The pair production mode is dominated by gluon-gluon and quark-antiquark annihilation, with a cross section that is nearly independent of the leptoquark Yukawa couplings and depends primarily on the leptoquark mass. The ATLAS and CMS Collaborations have derived upper limits on the pair production cross section multiplied by the branching fraction as a function of the leptoquark mass, which can be directly interpreted in the parameter space relevant to our analysis. In Fig. 2, we show the corresponding upper limits from recent searches at the $\sqrt{s} = 13$ TeV LHC using $\mathcal{L} \sim 139 \text{ fb}^{-1}$ and 138 fb^{-1} of data collected by ATLAS and CMS, respectively [66,67]. While the observation of one or more of these signals would provide unambiguous evidence of physics beyond the Standard Model, it would not indicate LNV, as events in these searches are selected by




FIG. 2. Upper limits on the leptoquark pair production cross section times branching fraction, $pp \rightarrow \text{LQ}\bar{\text{LQ}} \rightarrow X$, as a function of leptoquark mass m_{LQ} , from searches in the $X = \{qe qe, q\mu q\mu, ce ce, c\mu c\mu, be be, b\mu b\mu\}$ channels by ATLAS [66] and CMS [67] at the $\sqrt{s} = 13$ TeV LHC.

requiring an oppositely charged electron or muon, resulting in final states with zero net lepton number.

E. Combined constraints

In Fig. 3, we show the region of parameter space in the $m_{\omega^{2/3}} - m_{\tilde{X}_1}$ plane that is consistent with current constraints. Among the three oblique parameters, ΔT imposes the most stringent constraint, allowing only a narrow band of small




FIG. 3. Parameter space is shown in the $m_{\omega^{2/3}} - m_{\tilde{X}_1}$ plane. The blue shaded region is excluded by constraints from ΔT at 95% CL, while the yellow shaded region is ruled out by leptoquark pair production searches at the $\sqrt{s} = 13$ TeV LHC [66,67]. The green shaded region denotes the parameter space consistent with current constraints. Results are shown for the benchmark scenario $\{\theta_{\text{LQ}} = 0.2, \lambda_7 = 0.5, c \equiv \lambda_{ij}/\lambda'_{ij} = 0.13, m_1 = 0.04 \text{ eV}\}$.

mass splittings, $\Delta m = m_{\omega^{2/3}} - m_{X_1}$. For illustration, the region excluded by the 95% CL constraint on ΔT is displayed as a blue-shaded area in Fig. 3 for a representative benchmark point with $\theta_{LQ} = 0.2$, $\lambda_7 = 0.5$, $c \equiv \lambda_{ij}/\lambda'_{ij} = 0.13$, and $m_1 = 0.04$ eV. By contrast, the constraints from ΔS and ΔU , as well as those from low energy observables, are significantly weaker and do not restrict the scanned parameter region. The stronger sensitivity of ΔT arises because it is particularly sensitive to isospin-breaking effects, i.e., mass splittings within weak multiplets, whereas ΔS and ΔU are less affected by such splittings.

The region allowed by ΔT is further constrained by leptoquark searches at the LHC. Existing upper limits on the leptoquark pair production cross section times branching fraction exclude leptoquark masses $m_{LQ} \lesssim 1.4$ TeV. The parameter region consistent with all the current constraints, including neutrino mass bounds, low energy measurements, oblique parameters, and LHC limits, is illustrated as the green shaded region in Fig. 3. In the following, we focus on this allowed region, characterized by small mass splittings, $\Delta m = m_{\omega^{2/3}} - m_{X_1}$, and leptoquark masses $m_{LQ} \gtrsim 1.4$ TeV, to explore the potential for observing a signature of lepton number violation at the HL-LHC.

IV. LNV SIGNATURE AT THE LHC

The characteristic signature of LNV by two units, $|\Delta L| = 2$, at the LHC is the production of two same-sign charged leptons accompanied by jets, $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$. In the parameter space of interest, this final state can arise from cascade decays of both pair and singly produced leptoquarks. We classify the production mechanisms into three categories: QCD pair production, electroweak (EW) pair production, and single leptoquark production, as illustrated in Fig. 4.

The theoretical importance of such a $|\Delta L| = 2$ signal lies in its intimate connection to Majorana neutrino mass generation. The generalized form of the black-box theorem implies that any mechanism producing LNV at colliders necessarily feeds into a Majorana mass term for neutrinos [9,10]. In contrast to neutrinoless double beta decay searches, which probe only the electron channel, collider experiments can access a wide range of lepton-flavor combinations, providing complementary and potentially unique insights into the origin of neutrino mass. A schematic illustration of this correspondence between LNV collider signatures and Majorana mass generation is shown in Figs. 4 and 5. In the following, we discuss each of the three leading mechanisms for LNV production at the LHC.

(1) QCD pair production of leptoquarks

$$pp \rightarrow \omega^{2/3} \omega^{-2/3} \rightarrow (\omega^{2/3} \rightarrow \ell^+ j)(\omega^{-2/3} \rightarrow (X_1^* \rightarrow \ell^+ j)(W^- \rightarrow jj)), \quad (4.1)$$

$$pp \rightarrow X_2 X_1^* \rightarrow (X_2 \rightarrow (\omega^{2/3} \rightarrow \ell^+ j)(W^- \rightarrow jj))(X_1^* \rightarrow \ell^+ j), \quad (4.2)$$

$$pp \rightarrow X_2 X_2^* \rightarrow (X_2 \rightarrow (\omega^{2/3} \rightarrow \ell^+ j)(W^- \rightarrow jj))(X_2^* \rightarrow \ell^+ j). \quad (4.3)$$

In this production mode, QCD pair production of leptoquarks is followed by leptoquark decays through different channels to yield the $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ signature. A representative Feynman diagram for this channel is shown in Fig. 4 (left panel). For instance, in the case of $pp \rightarrow \omega^{2/3} \omega^{-2/3}$, $\omega^{2/3}$ may decay into $\omega^{2/3} \rightarrow \ell^+ j$, while its antiparticle $\omega^{-2/3}$ can undergo either on shell decay $\omega^{-2/3} \rightarrow X_1^* W^-$, if $m_{\omega^{2/3}} - m_{X_1} \geq m_W$, followed by $X_1^* \rightarrow \ell^+ j$ and

$W^- \rightarrow jj$, or three-body decay $\omega^{-2/3} \rightarrow X_1^* jj$, with $X_1^* \rightarrow \ell^+ j$. In our analysis presented in Sec. V, we account for the full matrix element generation for all relevant signal channels, incorporating both on shell and off shell contributions.

(2) Electroweak pair production of leptoquarks

$$pp \rightarrow \omega^{2/3} X_{1,2}^* \rightarrow (\omega^{2/3} \rightarrow \ell^+ j)(X_{1,2}^* \rightarrow \ell^+ j). \quad (4.4)$$




FIG. 4. Representative Feynman diagrams illustrating the QCD pair production of leptoquarks (left), electroweak pair production (center), and single leptoquark production (right), leading to LNV signatures $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ at the LHC.




FIG. 5. Diagrams illustrating the connection between Majorana neutrino mass generation and LNV signatures at the LHC [9,10]. The left, center, and right diagrams for neutrino mass generation correspond, respectively, to the LHC processes shown in the left, center, and right panels of Fig. 4.

This process is generated by quark-quark scattering, $q\bar{q}' \rightarrow \omega^{2/3} X_{1,2}^*$, mediated by s-channel W boson exchange. A representative Feynman diagram is shown in Fig. 4 (central panel). The relevant vertices $W^- \omega^{2/3} X_1$ and $W^- \omega^{2/3} X_2$ originate from the Higgs trilinear coupling, with interaction strengths proportional to $\cos \theta_{LQ}$ and $\sin \theta_{LQ}$, respectively. Further model details are provided in Sec. II.

To realize the LNV signature, both leptoquarks, $\omega^{2/3}$ and $X_{1,2}^*$, must decay into “visible” final states: $\omega^{2/3} \rightarrow \ell^+ j$ and $X_{1,2}^* \rightarrow \ell^+ j$. However, when the mass splitting $m_{\omega^{2/3}} - m_{X_{1,2}}$ exceeds the W boson mass, the decay $\omega^{2/3} \rightarrow W^+ X_{1,2}$ can become dominant. For smaller mass splittings $m_{\omega^{2/3}} - m_{X_{1,2}} < m_W$, the three-body decays $\omega^{2/3} \rightarrow jj X_1$ can still have appreciable rates for small values of λ . The visible decay mode $X_1 \rightarrow \ell^- u$, controlled by $\lambda' \sin \theta_{LQ}$, competes with invisible channels such as $X_1 \rightarrow \nu d$ and $X_1 \rightarrow \nu u$, which are sensitive to $\lambda' \sin \theta_{LQ}$ and $\lambda \cos \theta_{LQ}$, respectively. Since neutrino final states obscure the lepton number information, the LNV signal is appreciable when visible modes dominate, i.e., when $\lambda' \sin \theta_{LQ} > \lambda \cos \theta_{LQ}$. The case of X_2 is qualitatively similar to X_1 , with the favorable region for the visible decay channel being $\lambda' \cos \theta_{LQ} > \lambda \sin \theta_{LQ}$.

(3) Single leptoquark production

$$pp \rightarrow \omega^{-2/3} \ell^+ \rightarrow (\omega^{-2/3} \rightarrow (X_1^* \rightarrow \ell^+ j)(W^- \rightarrow jj)) \ell^+, \quad (4.5)$$

$$pp \rightarrow X_2 \ell^+ \rightarrow (X_2 \rightarrow (\omega^{2/3} \rightarrow \ell^+ j)(W^- \rightarrow jj)) \ell^+, \quad (4.6)$$

$$pp \rightarrow X_{1,2}^* \ell^+ j \rightarrow (X_{1,2}^* \rightarrow \ell^+ j) \ell^+ j, \quad (4.7)$$

$$pp \rightarrow \omega^{2/3} \ell^+ j \rightarrow (\omega^{2/3} \rightarrow \ell^+ j) \ell^+ j. \quad (4.8)$$

These processes involve leptoquark production in association with charged leptons and jets, primarily via quark-quark and quark-gluon fusion.

We show a representative Feynman diagram in Fig. 4 (right panel).

In the radiative neutrino mass scenario considered here, the flavor structure of the LNV signal is not uniquely determined by neutrino oscillation data. Unlike tree-level scenarios such as the type-II seesaw model, where the neutrino mass matrix and collider observables are controlled by the same Yukawa structure, the neutrino mass matrix shown in Eq. (3.3) involves two different Yukawa matrices, λ and λ' . In our analysis we assume the relation $\lambda_{ij} = c \lambda'_{ij}$, which reduces the number of free parameters but does not lead to unique predictions for flavor-specific collider observables.

Furthermore, the flavor composition of the LNV signal depends on the benchmark parameters entering the neutrino mass fit, including the leptoquark masses, mixing angle, and the proportionality constant c . Hence, different benchmark points that are equally consistent with neutrino oscillation data can lead to different flavor compositions of the $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ signal. Since the goal of the present work is to establish the inclusive LNV discovery potential at the HL-LHC, we focus on the flavor-inclusive signal rather than flavor-dependent observables.

V. ANALYSIS

The various leptoquark production modes and decay channels leading to the LNV signature $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ at the LHC are summarized in Table II. We simulate these LNV signal events using MadGraph5_aMC@NLO [95], interfaced with Pythia8 [96] for parton showering and hadronization. The leptoquark variant of the Zee Model is implemented in FeynRules [97]. Detector effects are simulated with Delphes3 [98], employing the default HL-LHC detector card.

We compute cross sections for the LNV signature $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ arising from the QCD pair production, EW pair production, and single leptoquark production modes at $\sqrt{s} = 13$ TeV. To preserve the lepton number information in the final state, we restrict our analysis to regions of parameter space where the visible decay mode of the lightest leptoquark $X_1 \rightarrow \ell^- j$ dominates or is at least comparable to the neutrino-containing decay channel. From the Yukawa Lagrangian in Eq. (2.7), this condition

TABLE II. Summary of QCD and EW mediated leptoquark pair production, as well as single leptoquark production channels, within the leptoquark extension of the Zee model, together with their decay topologies that give rise to the LNV, $|\Delta L| = 2$, same-sign dilepton plus jets final state, $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$, at the LHC.

Production modes		LNV signature: $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$	
QCD	$\omega^{2/3} \omega^{-2/3}$	$\omega^{2/3} \rightarrow \ell^+ j$	$\omega^{-2/3} \rightarrow (X_1^* \rightarrow \ell^+ j)(W^- \rightarrow \text{jets})$
		$\omega^{-2/3} \rightarrow \ell^- j$	$\omega^{2/3} \rightarrow (X_1 \rightarrow \ell^- j)(W^+ \rightarrow \text{jets})$
	$X_2 X_1^*$	$X_1^* \rightarrow \ell^+ j$	$X_2 \rightarrow (\omega^{2/3} \rightarrow \ell^+ j)(W^- \rightarrow \text{jets})$
	$X_1 X_2^*$	$X_1 \rightarrow \ell^- j$	$X_2^* \rightarrow (\omega^{-2/3} \rightarrow \ell^- j)(W^+ \rightarrow \text{jets})$
	$X_2 X_2^*$	$X_2^* \rightarrow \ell^+ j$	$X_2 \rightarrow (\omega^{2/3} \rightarrow \ell^+ j)(W^- \rightarrow \text{jets})$
		$X_2 \rightarrow \ell^- j$	$X_2^* \rightarrow (\omega^{-2/3} \rightarrow \ell^- j)(W^+ \rightarrow \text{jets})$
EW	$\omega^{2/3} X_1^*$	$\omega^{2/3} \rightarrow \ell^+ j$	$X_1^* \rightarrow \ell^+ j$
	$\omega^{-2/3} X_1$	$\omega^{-2/3} \rightarrow \ell^- j$	$X_1 \rightarrow \ell^- j$
	$\omega^{2/3} X_2^*$	$\omega^{2/3} \rightarrow \ell^+ j$	$X_2^* \rightarrow \ell^+ j$
	$\omega^{-2/3} X_2$	$\omega^{-2/3} \rightarrow \ell^- j$	$X_2 \rightarrow \ell^- j$
Single-LQ	$\omega^{-2/3} \ell^+$	$\omega^{-2/3} \rightarrow (X_1^* \rightarrow \ell^+ j)(W^- \rightarrow \text{jets})$	
	$\omega^{2/3} \ell^-$	$\omega^{2/3} \rightarrow (X_1 \rightarrow \ell^- j)(W^+ \rightarrow \text{jets})$	
	$X_2 \ell^+$	$X_2 \rightarrow (\omega^{2/3} \rightarrow \ell^+ j)(W^- \rightarrow \text{jets})$	
	$X_2^* \ell^-$	$X_2^* \rightarrow (\omega^{-2/3} \rightarrow \ell^- j)(W^+ \rightarrow \text{jets})$	
	$X_{1,2}^* \ell^+ j$	$X_{1,2}^* \rightarrow \ell^+ j$	
	$X_{1,2} \ell^- j$	$X_{1,2} \rightarrow \ell^- j$	
	$\omega^{2/3} \ell^+ j$	$\omega^{2/3} \rightarrow \ell^+ j$	
	$\omega^{-2/3} \ell^- j$	$\omega^{-2/3} \rightarrow \ell^- j$	

translates to $\lambda' \sin \theta_{LQ} \gtrsim \lambda \cos \theta_{LQ}$. For the production rates, we compute the leading order (LO) parton level cross sections using MadGraph5_aMC@NLO and account for higher-order effects through an overall next-to-leading-order (NLO) K factor. The NLO K factor for the QCD pair production channel is obtained by comparing the LO cross-sections with the NLO prediction [99]. For the electroweak pair production and single leptoquark production channels, we restrict ourselves to LO predictions. Given that these two channels are typically suppressed by 2 orders of magnitude relative to the QCD production mode in the relevant parameter regions, as shown later in this section, neglecting their NLO corrections has a negligible effect on our HL-LHC projections.

In Fig. 6, we present the stacked cross sections for $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ mediated by QCD pair production, EW pair production, and single leptoquark production as a function of the leptoquark mass $m_{\omega^{2/3}}$. The QCD pair production mode dominates over the entire mass range, yielding cross sections several orders of magnitude larger than those from EW pair or single leptoquark production for both benchmark scenarios. This enhancement arises from the purely QCD-driven nature of that production mode, larger gluon parton distribution functions in the proton, and the sizable color factor associated with gluon-gluon initiated channels. In contrast, EW leptoquark pair production is suppressed due to their dependence on electroweak couplings and quark-initiated parton luminosities. Single leptoquark production is further reduced, as it

requires a leptoquark Yukawa coupling insertion already at the production level.² For instance, the $qg \rightarrow \omega^{-2/3} \ell^+$ channel is suppressed by a factor of $\sim |\lambda|^2$. In the allowed parameter space, the Yukawa couplings to the valence quarks are typically small $|\lambda_{ed}|, |\lambda_{ud}| \lesssim 10^{-3}$, resulting in subdominant contributions from the single leptoquark channel, as observed in Fig. 6. Despite this suppression, we account for all three production mechanisms in our signal simulation for completeness.

It is worth emphasizing that a sizable LNV signal in this model does not imply an enhanced neutrino mass contribution. The neutrino mass matrix is generated radiatively and is constrained to reproduce the observed neutrino masses and mixing parameters through Eq. (3.3). By contrast, the dominant contribution to the LNV signal at the LHC arises from QCD-mediated leptoquark pair production, whose cross section depends primarily on the leptoquark masses and is largely independent of the Yukawa couplings entering the neutrino mass matrix. The Yukawa couplings mainly affect the subsequent decay branching fractions of the leptoquarks. Thus, an observable LNV signal at the HL-LHC is fully compatible with the small neutrino masses required by current neutrino oscillation data.

²This explains why the QCD-driven channel in Fig. 4 (left panel) yields the largest rate among the three LNV production mechanisms, even though the corresponding neutrino-mass diagram arises only at higher loop order, as shown in Fig. 5.




FIG. 6. Stacked cross section for the LNV process, $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$, mediated by QCD $pp \rightarrow \text{LQ LQ}$ (yellow), EW $pp \rightarrow \text{LQ LQ}$ (blue), and single leptoquark production $pp \rightarrow \text{LQ} \ell$ (gray) at $\sqrt{s} = 13$ TeV, assuming leptoquark mixing angle $\theta_{\text{LQ}} = 0.2$, $\Delta m \equiv m_{\omega^{2/3}} - m_{X_1} = 10$ GeV, $\lambda_7 = 0.5$, $c \equiv \lambda_{ij}/\lambda'_{ij} = 0.13$, and $m_1 = 0.04$ eV. The 95% confidence level limits are shown for an integrated luminosity of 137 fb^{-1} (red dashed) and for the HL-LHC projection with 3 ab^{-1} (red solid).

The LNV signal $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ is subject to substantial background arising from nonprompt leptons, which are challenging to simulate reliably. Therefore, we adopt the background estimates reported in the CMS same-sign dilepton plus jets search with 137 fb^{-1} of data [100], and rescale them to 3 ab^{-1} for HL-LHC projections. Following the CMS event selection criteria in [100], we require signal events to contain exactly two same-sign isolated leptons with transverse momentum $p_{T\ell} > 25$ GeV and pseudorapidity $|\eta_\ell| < 2.5$ for electrons and $|\eta_\mu| < 2.4$ for muons. Events must also include at least two jets, reconstructed using the anti- k_T jet algorithm with radius $R = 0.4$, satisfying $p_{Tj} > 40$ GeV and $|\eta_j| < 2.4$. To suppress background contributions, events with same-flavor dileptons having invariant mass $m_{\ell\ell} < 12$ GeV, or different-flavor leptons with $m_{\ell\ell'} < 8$ GeV, are vetoed. Events are also

required to have only small missing energy $\cancel{E}_T < 50$ GeV and a large scalar sum of the transverse momenta $H_T > 1300$ GeV.

Using these event selection criteria and background estimates from the CMS analysis in Ref. [100], we evaluate the projected sensitivity to LNV at the HL-LHC. To illustrate the potential reach, we consider a representative benchmark point in the currently allowed parameter space: $\theta_{\text{LQ}} = 0.2$, $m_{\omega^{2/3}} = 1.4$ TeV, $\Delta m = 10$ GeV, $\lambda_7 = 0.5$, $c \equiv \lambda_{ij}/\lambda'_{ij} = 0.13$, and $m_1 = 0.04$ eV. The signal significance is quantified using $S/\sqrt{B}$, where S is the expected number of signal events and B is the luminosity-scaled projected background rate derived from the CMS analysis [100].

In Fig. 7 (left panel), we show the signal significance as a function of the integrated luminosity $\mathcal{L}$. For the representative parameter choices, a 2σ sensitivity to LNV can be achieved around $\mathcal{L} \sim 790 \text{ fb}^{-1}$. In Fig. 7 (right panel), the signal significance is shown as a function of leptoquark mass $m_{\omega^{2/3}}$, assuming a mass splitting $\Delta m = m_{\omega^{2/3}} - m_{X_1} = 10$ GeV. We observe that the HL-LHC with $\mathcal{L} = 3 \text{ ab}^{-1}$ will be able to probe the LNV signal for leptoquark masses up to approximately $m_{\omega^{2/3}} \sim 1.5$ TeV.

VI. CONCLUSION

Determining whether a lepton number is a conserved symmetry in nature remains a fundamental question, closely linked to the Majorana versus Dirac character of neutrinos. Observation of lepton number violation by two units at colliders would provide a direct and complementary probe to neutrinoless double beta decay, opening a window into the mechanism behind neutrino mass generation. Motivated by this, we have investigated the potential for detecting the LNV process $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ at the HL-LHC within the leptoquark variant of the Zee model.




FIG. 7. Signal significance $S/\sqrt{B}$ for the LNV process $pp \rightarrow \ell^\pm \ell'^\pm + \text{jets}$ at the LHC with $\sqrt{s} = 13$ TeV. Left: Significance as a function of integrated luminosity. Right: Significance at the HL-LHC with $\mathcal{L} = 3 \text{ ab}^{-1}$ as a function of the leptoquark mass $m_{\omega^{2/3}}$, assuming $\theta_{\text{LQ}} = 0.2$, $\Delta m = 10$ GeV, $\lambda_7 = 0.5$, $c \equiv \lambda_{ij}/\lambda'_{ij} = 0.13$, and $m_1 = 0.04$ eV.

In this scenario, neutrino masses are generated radiatively at one loop, while the same interactions give rise to distinct LNV signatures at colliders. After applying current experimental constraints, including neutrino mass bounds, charged lepton flavor violation limits, electroweak precision tests, and existing LHC searches for leptoquarks, we identified the allowed parameter space and the dominant production mechanisms for the $|\Delta L| = 2$ final state. We divide these channels into three classes: QCD leptoquark pair production, electroweak pair production, and single leptoquark production. Among these, QCD pair production overwhelmingly dominates the signal across the relevant parameter space.

Based on background estimates from the CMS same-sign dilepton plus jets analysis [100], we performed a collider sensitivity study for the HL-LHC. Our results indicate that leptoquark masses up to $m_{LQ} \sim 1.5$ TeV could be probed through this LNV channel. These results

demonstrate that the same-sign dilepton plus jets signature is a promising probe of lepton number violation at the HL-LHC, providing a direct test of leptoquark-induced radiative neutrino mass generation mechanism.

ACKNOWLEDGMENTS

The work of K. S. B. and D. G. is supported by the U.S. Department of Energy under Grant No. DE-SC0016013. The work of R. K. B. is supported by the World Premier International Research Center Initiative (WPI), MEXT, Japan, and by JSPS KAKENHI Grant No. JP24K22876. This work used computing resources at Kavli IPMU.

DATA AVAILABILITY

All results can be reproduced using the publicly available packages cited in the text.

-
- [1] A. Zee, A theory of lepton number violation, neutrino majorana mass, and oscillation, *Phys. Lett.* **93B**, 389 (1980); *Phys. Lett.* **95B**, 461(E) (1980).
 - [2] L. J. Hall and M. Suzuki, Explicit R -parity breaking in supersymmetric models, *Nucl. Phys.* **B231**, 419 (1984).
 - [3] A. Zee, Quantum numbers of majorana neutrino masses, *Nucl. Phys.* **B264**, 99 (1986).
 - [4] K. S. Babu, Model of “Calculable” majorana neutrino masses, *Phys. Lett. B* **203**, 132 (1988).
 - [5] E. Carquin, N. A. Neill, J. C. Helo, and M. Hirsch, Exotic colored fermions and lepton number violation at the LHC, *Phys. Rev. D* **99**, 115028 (2019).
 - [6] L. M. Krauss, S. Nasri, and M. Trodden, A model for neutrino masses and dark matter, *Phys. Rev. D* **67**, 085002 (2003).
 - [7] Y. Cai, T. Han, T. Li, and R. Ruiz, Lepton number violation: Seesaw models and their collider tests, *Front. Phys.* **6**, 40 (2018).
 - [8] K. S. Babu, P. S. B. Dev, S. Jana, and A. Thapa, Non-standard interactions in radiative neutrino mass models, *J. High Energy Phys.* **03** (2020) 006.
 - [9] K. S. Babu, R. K. Barman, D. Gonçalves, and A. Ismail, Probing lepton number violation and Majorana nature of neutrinos at the LHC, *J. High Energy Phys.* **06** (2024) 132.
 - [10] J. Schechter and J. Valle, Neutrinoless Double beta decay in $SU(2) \times U(1)$ theories, *Phys. Rev. D* **25**, 2951 (1982).
 - [11] W. H. Furry, On transition probabilities in double beta-deintegration, *Phys. Rev.* **56**, 1184 (1939).
 - [12] M. Duerr, M. Lindner, and A. Merle, On the quantitative impact of the Schechter-Valle theorem, *J. High Energy Phys.* **06** (2011) 091.
 - [13] W.-Y. Keung and G. Senjanovic, Majorana neutrinos and the production of the right-handed charged gauge boson, *Phys. Rev. Lett.* **50**, 1427 (1983).
 - [14] S. N. Gninenko, M. M. Kirsanov, N. V. Krasnikov, and V. A. Matveev, Detection of heavy Majorana neutrinos and right-handed bosons, *Phys. At. Nucl.* **70**, 441 (2007).
 - [15] A. Maiezza, M. Nemevsek, F. Nesti, and G. Senjanovic, Left-right symmetry at LHC, *Phys. Rev. D* **82**, 055022 (2010).
 - [16] M. Nemevsek, F. Nesti, G. Senjanovic, and Y. Zhang, First limits on left-right symmetry scale from LHC data, *Phys. Rev. D* **83**, 115014 (2011).
 - [17] C.-Y. Chen and P. S. B. Dev, Multi-lepton collider signatures of heavy Dirac and majorana neutrinos, *Phys. Rev. D* **85**, 093018 (2012).
 - [18] J. Chakraborty, J. Gluza, R. Sevilano, and R. Szafron, Left-right symmetry at LHC and precise 1-loop low energy data, *J. High Energy Phys.* **07** (2012) 038.
 - [19] J. A. Aguilar-Saavedra and F. R. Joaquim, Measuring heavy neutrino couplings at the LHC, *Phys. Rev. D* **86**, 073005 (2012).
 - [20] T. Han, I. Lewis, R. Ruiz, and Z.-g. Si, Lepton number violation and W' Chiral couplings at the LHC, *Phys. Rev. D* **87**, 035011 (2013); *Phys. Rev. D* **87**, 039906(E) (2013).
 - [21] C.-Y. Chen, P. S. B. Dev, and R. N. Mohapatra, Probing heavy-light neutrino mixing in left-right seesaw models at the LHC, *Phys. Rev. D* **88**, 033014 (2013).
 - [22] P. S. B. Dev, A. Pilaftsis, and U.-k. Yang, New production mechanism for heavy neutrinos at the LHC, *Phys. Rev. Lett.* **112**, 081801 (2014).
 - [23] B. Dutta, R. Eusebi, Y. Gao, T. Ghosh, and T. Kamon, Exploring the doubly charged Higgs boson of the left-right symmetric model using vector boson fusionlike events at the LHC, *Phys. Rev. D* **90**, 055015 (2014).
 - [24] J. Gluza and T. Jeliński, Heavy neutrinos and the $pp \rightarrow lljj$ CMS data, *Phys. Lett. B* **748**, 125 (2015).

- [25] J. N. Ng, A. de la Puente, and B. W.-P. Pan, Search for heavy right-handed neutrinos at the LHC and beyond in the same-sign same-flavor leptons final state, *J. High Energy Phys.* **12** (2015) 172.
- [26] A. Maiezza, M. Nemevšek, and F. Nesti, Lepton number violation in Higgs decay at LHC, *Phys. Rev. Lett.* **115**, 081802 (2015).
- [27] F. F. Deppisch, P. S. Bhupal Dev, and A. Pilaftsis, Neutrinos and collider physics, *New J. Phys.* **17**, 075019 (2015).
- [28] C. Degrande, O. Mattelaer, R. Ruiz, and J. Turner, Fully-automated precision predictions for heavy neutrino production mechanisms at hadron colliders, *Phys. Rev. D* **94**, 053002 (2016).
- [29] P. S. B. Dev, R. N. Mohapatra, and Y. Zhang, Probing the Higgs sector of the minimal left-right symmetric model at future hadron colliders, *J. High Energy Phys.* **05** (2016) 174.
- [30] A. Roitgrund, Search for like-sign dileptons plus two jets signal in the framework of the manifest left-right symmetric model, *J. High Energy Phys.* **01** (2021) 031.
- [31] M. Nemevšek, F. Nesti, and G. Popara, Keung-Senjanović process at the LHC: From lepton number violation to displaced vertices to invisible decays, *Phys. Rev. D* **97**, 115018 (2018).
- [32] D. A. Dicus, D. D. Karatas, and P. Roy, Lepton non-conservation at supercollider energies, *Phys. Rev. D* **44**, 2033 (1991).
- [33] A. Datta, M. Guchait, and A. Pilaftsis, Probing lepton number violation via majorana neutrinos at hadron supercolliders, *Phys. Rev. D* **50**, 3195 (1994).
- [34] A. Ali, A. V. Borisov, and N. B. Zamorin, Majorana neutrinos and same sign Dilepton production at LHC and in rare meson decays, *Eur. Phys. J. C* **21**, 123 (2001).
- [35] T. Han and B. Zhang, Signatures for Majorana neutrinos at hadron colliders, *Phys. Rev. Lett.* **97**, 171804 (2006).
- [36] J. Kersten and A. Y. Smirnov, Right-handed neutrinos at CERN LHC and the mechanism of neutrino mass generation, *Phys. Rev. D* **76**, 073005 (2007).
- [37] F. del Aguila, J. A. Aguilar-Saavedra, and R. Pittau, Heavy neutrino signals at large hadron colliders, *J. High Energy Phys.* **10** (2007) 047.
- [38] A. Atre, T. Han, S. Pascoli, and B. Zhang, The search for heavy majorana neutrinos, *J. High Energy Phys.* **05** (2009) 030.
- [39] D. Alva, T. Han, and R. Ruiz, Heavy Majorana neutrinos from $W\gamma$ fusion at hadron colliders, *J. High Energy Phys.* **02** (2015) 072.
- [40] A. Das and N. Okada, Improved bounds on the heavy neutrino productions at the LHC, *Phys. Rev. D* **93**, 033003 (2016).
- [41] A. Das, P. Konar, and S. Majhi, Production of Heavy neutrino in next-to-leading order QCD at the LHC and beyond, *J. High Energy Phys.* **06** (2016) 019.
- [42] A. Das, P. Konar, and A. Thalapillil, Jet substructure shedding light on heavy Majorana neutrinos at the LHC, *J. High Energy Phys.* **02** (2018) 083.
- [43] M. Drewes, J. Klarić, and P. Klose, On lepton number violation in heavy neutrino decays at colliders, *J. High Energy Phys.* **11** (2019) 032.
- [44] B. Fuks, J. Neundorff, K. Peters, R. Ruiz, and M. Saimpert, Majorana neutrinos in same-sign $W^\pm W^\pm$ scattering at the LHC: Breaking the TeV barrier, *Phys. Rev. D* **103**, 055005 (2021).
- [45] P. D. Bolton, J. Kriewald, M. Nemevšek, F. Nesti, and J. C. Vasquez, Hadron colliders signatures of lepton number violation in the type II seesaw model, *Phys. Rev. D* **111**, 035016 (2025).
- [46] J. C. Helo, M. Hirsch, S. G. Kovalenko, and H. Pas, Neutrinoless double beta decay and lepton number violation at the LHC, *Phys. Rev. D* **88**, 011901 (2013).
- [47] J. C. Helo, M. Hirsch, H. Päs, and S. G. Kovalenko, Short-range mechanisms of neutrinoless double beta decay at the LHC, *Phys. Rev. D* **88**, 073011 (2013).
- [48] F. del Aguila, M. Chala, A. Santamaria, and J. Wudka, Discriminating between lepton number violating scalars using events with four and three charged leptons at the LHC, *Phys. Lett. B* **725**, 310 (2013).
- [49] F. del Aguila and M. Chala, LHC bounds on lepton number violation mediated by doubly and singly-charged scalars, *J. High Energy Phys.* **03** (2014) 027.
- [50] T. Peng, M. J. Ramsey-Musolf, and P. Winslow, TeV lepton number violation: From neutrinoless double- β decay to the LHC, *Phys. Rev. D* **93**, 093002 (2016).
- [51] J. Harz, M. J. Ramsey-Musolf, T. Shen, and S. Urrutia-Quiroga, TeV-scale lepton number violation: Connecting leptogenesis, neutrinoless double beta decay, and colliders, *Phys. Rev. D* **110**, 035024 (2024).
- [52] M. L. Graesser, G. Li, M. J. Ramsey-Musolf, T. Shen, and S. Urrutia-Quiroga, Uncovering a chirally suppressed mechanism of $0\nu\beta\beta$ decay with LHC searches, *J. High Energy Phys.* **10** (2022) 034.
- [53] M. Aaboud *et al.* (ATLAS Collaboration), Search for heavy Majorana or Dirac neutrinos and right-handed W gauge bosons in final states with two charged leptons and two jets at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **01** (2019) 016.
- [54] A. M. Sirunyan *et al.* (CMS Collaboration), Search for heavy Majorana neutrinos in same-sign dilepton channels in proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **01** (2019) 122.
- [55] T. G. Rizzo, Doubly charged Higgs bosons and lepton number violating processes, *Phys. Rev. D* **25**, 1355 (1982); *Phys. Rev. D* **27**, 657(A) (1983).
- [56] A. Crivellin, M. Ghezzi, L. Panizzi, G. M. Pruna, and A. Signer, Low- and high-energy phenomenology of a doubly charged scalar, *Phys. Rev. D* **99**, 035004 (2019).
- [57] T. Li, C.-Y. Yao, and M. Yuan, Searching for heavy neutral lepton and lepton number violation through VBS at high-energy muon colliders, *J. High Energy Phys.* **09** (2023) 131.
- [58] O. Mikulenko and M. Marinichenko, Measuring lepton number violation in heavy neutral lepton decays at the future muon collider, *J. High Energy Phys.* **01** (2024) 032.
- [59] S. P. Maharathy and M. Mitra, Type-II see-saw at $\mu^+\mu^-$ collider, *Phys. Lett. B* **844**, 138105 (2023).
- [60] C. H. de Lima, D. McKeen, J. N. Ng, M. Shamma, and D. Tuckler, Probing lepton number violation at same-sign lepton colliders, *Phys. Rev. D* **111**, 075002 (2025).

- [61] Q.-H. Cao, K. Cheng, and Y. Liu, Distinguishing Dirac from majorana heavy neutrino at future lepton colliders, *Phys. Rev. Lett.* **134**, 021801 (2025).
- [62] F.-X. Yang, F.-L. Shao, Z.-L. Han, F. Huang, Y. Jin, and H. Li, Lepton number violation Higgs decay at muon collider, *J. High Energy Phys.* **09** (2025) 200.
- [63] R. Aaij *et al.* (LHCb Collaboration), Search for Majorana neutrinos in $B^- \rightarrow \pi^+ \mu^- \mu^-$ decays, *Phys. Rev. Lett.* **112**, 131802 (2014).
- [64] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, J. a. P. Pinheiro, and T. Schwetz, NuFit-6.0: Updated global analysis of three-flavor neutrino oscillations, *J. High Energy Phys.* **12** (2024) 216.
- [65] K. S. Babu and J. Julio, Two-loop neutrino mass generation through leptoquarks, *Nucl. Phys.* **B841**, 130 (2010).
- [66] G. Aad *et al.* (ATLAS Collaboration), Search for pairs of scalar leptoquarks decaying into quarks and electrons or muons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector, *J. High Energy Phys.* **10** (2020) 112.
- [67] A. Hayrapetyan *et al.* (CMS Collaboration), Search for pair production of scalar and vector leptoquarks decaying to muons and bottom quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Rev. D* **109**, 112003 (2024).
- [68] D. Aristizabal Sierra, M. Hirsch, and S. G. Kovalenko, Leptoquarks: Neutrino masses and accelerator phenomenology, *Phys. Rev. D* **77**, 055011 (2008).
- [69] D. Dercks, H. Dreiner, M. E. Krauss, T. Opferkuch, and A. Reinert, R -parity violation at the LHC, *Eur. Phys. J. C* **77**, 856 (2017).
- [70] M. Aker *et al.* (KATRIN Collaboration), Direct neutrino-mass measurement based on 259 days of KATRIN data, *Science* **388**, adq9592 (2025).
- [71] S. Navas *et al.* (Particle Data Group), Review of particle physics, *Phys. Rev. D* **110**, 030001 (2024).
- [72] L. Lavoura, General formulae for $f_1 \rightarrow f_2 \gamma$, *Eur. Phys. J. C* **29**, 191 (2003).
- [73] A. Crivellin, C. Greub, D. Müller, and F. Saturnino, Scalar leptoquarks in leptonic processes, *J. High Energy Phys.* **02** (2021) 182.
- [74] K. Afanaciev *et al.* (MEG II Collaboration), New limit on the $\mu^+ \rightarrow e^+ \gamma$ decay with the MEG II experiment, *Eur. Phys. J. C* **85**, 1177 (2025).
- [75] B. Aubert *et al.* (BABAR Collaboration), Searches for lepton flavor violation in the decays $\tau^\pm \rightarrow e^\pm \gamma$ and $\tau^\pm \rightarrow \mu^\pm \gamma$, *Phys. Rev. Lett.* **104**, 021802 (2010).
- [76] A. Abdesselam *et al.* (Belle Collaboration), Search for lepton-flavor-violating tau-lepton decays to $\ell \gamma$ at Belle, *J. High Energy Phys.* **10** (2021) 19.
- [77] W. Bertl, R. Engfer *et al.*, A search for mu-e conversion in muonic gold, *Eur. Phys. J. C* **47**, 337 (2006).
- [78] W. Honecker, C. Dohmen *et al.* (SINDRUM II Collaboration), Improved limit on the branching ratio of $\mu \rightarrow e$ conversion on lead, *Phys. Rev. Lett.* **76**, 200 (1996).
- [79] D. Ambrose *et al.* (BNL E871 Collaboration), First observation of the rare decay mode $K_L^0 \rightarrow e^+ e^-$, *Phys. Rev. Lett.* **81**, 4309 (1998).
- [80] F. Takahashi *et al.* (Particle Data Group), Review of particle physics, *Int. J. Mod. Phys. A* **41**, 2630011 (2026).
- [81] D. Ambrose *et al.* (BNL Collaboration), New limit on muon and electron lepton number violation from $K_L^0 \rightarrow \mu^\pm e^\mp$ decay, *Phys. Rev. Lett.* **81**, 5734 (1998).
- [82] E. Cortina Gil *et al.* (NA62 Collaboration), A measurement of the $K^+ \rightarrow \pi^+ \mu^+ \mu^-$ decay, *J. High Energy Phys.* **11** (2022) 011.
- [83] A. Sher *et al.*, An Improved upper limit on the decay $K^+ \rightarrow \pi^+ \mu^+ e^-$, *Phys. Rev. D* **72**, 012005 (2005).
- [84] S. Alibocus *et al.* (NA62 Collaboration), Measurement of the branching ratio of the $K^+ \rightarrow \nu \bar{\nu}$ decay, *arXiv*: 2607.16413.
- [85] R. Mandal and A. Pich, Constraints on scalar leptoquarks from lepton and kaon physics, *J. High Energy Phys.* **12** (2019) 089.
- [86] K. S. Babu, P. S. B. Dev, S. Jana, and A. Thapa, Unified framework for B -anomalies, muon $g - 2$ and neutrino masses, *J. High Energy Phys.* **03** (2021) 179.
- [87] W. Grimus, L. Lavoura, O. M. Ogreid, and P. Osland, The Oblique parameters in multi-Higgs-doublet models, *Nucl. Phys.* **B801**, 81 (2008).
- [88] A. Crivellin, D. Müller, and F. Saturnino, Leptoquarks in oblique corrections and Higgs signal strength: status and prospects, *J. High Energy Phys.* **11** (2020) 094.
- [89] G. Aad *et al.* (ATLAS Collaboration), Combination of searches for pair-produced leptoquarks at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Lett. B* **854**, 138736 (2024).
- [90] G. Aad *et al.* (ATLAS Collaboration), Search for supersymmetry in final states with missing transverse momentum and charm-tagged jets using 139 fb $^{-1}$ of proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **02** (2025) 193.
- [91] G. Aad *et al.* (ATLAS Collaboration), Search for pair production of third-generation leptoquarks decaying into a bottom quark and a τ -lepton with the ATLAS detector, *Eur. Phys. J. C* **83**, 1075 (2023).
- [92] G. Aad *et al.* (ATLAS Collaboration), Search for leptoquarks decaying into the $b\tau$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **10** (2023) 001.
- [93] G. Aad *et al.* (ATLAS Collaboration), Search for leptoquark pair production decaying into $te^- \bar{t}e^+$ or $t\mu^- \bar{t}\mu^+$ in multi-lepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Eur. Phys. J. C* **84**, 818 (2024).
- [94] I. d. M. Varzielas and A. Sengupta, Constraining flavoured leptoquarks with LHC and LFV, *Nucl. Phys.* **B1001**, 116495 (2024).
- [95] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *J. High Energy Phys.* **07** (2014) 079.
- [96] T. Sjöstrand, S. Mrenna, and P. Z. Skands, A Brief Introduction to PYTHIA 8.1, *Comput. Phys. Commun.* **178**, 852 (2008).
- [97] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr, and B. Fuks, FeynRules 2.0—A complete toolbox for tree-level

- phenomenology, *Comput. Phys. Commun.* **185**, 2250 (2014).
- [98] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens, and M. Selvaggi (DELPHES 3 Collaboration), DELPHES 3, A modular framework for fast simulation of a generic collider experiment, *J. High Energy Phys.* **02** (2014) 057.
- [99] M. Kramer, T. Plehn, M. Spira, and P.M. Zerwas, Pair production of scalar leptoquarks at the CERN LHC, *Phys. Rev. D* **71**, 057503 (2005).
- [100] A.M. Sirunyan *et al.* (CMS Collaboration), Search for physics beyond the standard model in events with jets and two same-sign or at least three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **80**, 752 (2020).