

Phase-Space Microscopes for Quantum Gases: Imaging Conjugate Variables and Momentum-Weighted Densities

N. R. Cooper¹, Y. Yang¹, and C. Weitenberg²

¹*TCM Group, Cavendish Laboratory, University of Cambridge, J.J. Thomson Avenue, Cambridge, CB3 0US, United Kingdom*

²*Department of Physics, TU Dortmund University, 44227 Dortmund, Germany*



(Received 31 March 2026; revised 23 June 2026; accepted 20 August 2026; published 21 October 2026)

Quantum gas microscopes offer unprecedented insights into quantum many-body states of cold atomic gases. Here we introduce concrete protocols for extending quantum gas microscopes to measure in phase space, by mapping momentum onto auxiliary degrees of freedom and using positive operator-valued measures. We distinguish between two distinct operational modes. In the Husimi- Q phase-space microscope, position and momentum are jointly measured; in this mode, the fundamental quantum noise is distributed between position and momentum. Conversely, the averaged-mode phase space microscope extracts the spatial dependence of averages of the momentum density (and its moments); these averages can be retrieved with arbitrary spatial resolution. We illustrate the utility of these techniques in diverse physical settings.

DOI: [10.1103/12m-rlg8](https://doi.org/10.1103/12m-rlg8)

Quantum gas microscopes (QGMs) perform spatially resolved detection of the individual particles in a quantum many-body system of ultracold atoms and allow for microscopic insights into quantum phases [1]. They were first realized in lattice systems [2,3] and later in the continuum [4–7]. Projective measurements of atomic positions allow access not only to density-density correlators [8] but also to various multipoint correlators including string order [8–10]. Recent experiments went beyond detection in the position basis by measuring local currents and kinetic energy operators along certain bonds of an optical lattice [11,12], as well as local phases on the lattice sites [13]. An effective way to overcome the optical diffraction limit is the matter-wave microscope, which magnifies the density distribution prior to optical imaging using two matter-wave lenses formed by quarter-period evolutions in harmonic traps [14,15]. This method also gives access to matter-wave Fourier space and allows manipulation to provide an analog of phase-contrast imaging [13,16] or access to various long-range off-diagonal correlators [17].

Here we show how to extend these techniques to phase-space microscopes, which access atomic occupations and correlations jointly in both position and momentum. Despite being conjugate variables—and, therefore, noncommuting—joint measurements of position and momentum are permitted, albeit at the expense of introducing quantum noise via the

measurement apparatus [18,19]. We will show how this can be done for cold gases by mapping momentum (or some function of it) onto auxiliary degrees of freedom. This realizes generalized measurements, described by positive operator-valued measures (POVMs), that reflect the noncommuting observables [19]. We discuss protocols for mapping momentum space to an auxiliary dimension or to an auxiliary spin by appropriate manipulation in matter-wave Fourier space. We further show that the approach can be adapted to allow measurements of *averages* of the momentum density (or its moments). Such observables can, in principle, be measured with arbitrary spatial resolution. We illustrate the utility of phase-space microscopes for a variety of physical settings.

We first introduce the phase-space microscope for a single particle of mass M moving in a continuous system in one dimension, with system coordinate x , and using the z dimension as the auxiliary space (Fig. 1). The particle starts in the state $\Psi_i(x, z) = \psi_i(x)\phi_0(z)$, where $\psi_i(x)$ is the state we wish to characterize. We take $\phi_0(z) = e^{-z^2/2\sigma^2}/(\pi\sigma^2)^{1/4}$, assuming the particle to be in the ground state of a harmonic confinement in z with oscillator length σ . We encode the momentum p_x in z by applying a unitary operator,

$$\hat{U}_{\text{kick}} = e^{i\alpha\hat{p}_x\hat{z}/\hbar}, \quad (1)$$

that boosts the momentum in the z direction by p_x scaled by a factor α . To do so, one can use a pulsed harmonic trap in x of trapping frequency $\omega = 2\pi/T$ for a quarter-period duration $T/4$ to map momentum p_x onto position x , thereby transforming to the Fourier plane. One then applies a potential $V_{\text{kick}}(x, z) = -\hbar\kappa xz$ for a short time $\tau \ll T$, which imparts an x -dependent impulse $\hbar\kappa\tau x$ along z . Such a saddle

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.




FIG. 1. Sketch of the protocol for mapping momentum to an auxiliary dimension to realize a 1D Husimi- Q phase-space microscope. (a) Two atoms with initial positions x_1 and x_2 and momenta p_1 and p_2 in the physical x dimension. (b) A $T/4$ pulse is applied to move to the Fourier plane in the x direction: the horizontal position of a particle now encodes its initial momentum, so it is labeled by $p_{1,2}$. (c) A spatially dependent impulse maps this momentum to a momentum in the auxiliary z dimension. (d) A second $T/4$ pulse along both directions creates an image in the x direction and maps the acquired z momentum to a z displacement. The protocol gives a choice of how to distribute uncertainty between x and p_x .

potential can be realized by a magnetic field away from the Helmholtz condition [20]. Acting with another quarter-period harmonic trap in x then maps back to position space. Denoting the action of each quarter-period harmonic trap (with oscillator length $a_0 = \sqrt{\hbar/M\omega}$) by the unitary operator $\hat{U}_x$, the overall transformation is

$$\hat{U}_x e^{i\kappa\tau\hat{x}} \hat{U}_x = \hat{U}_x^2 e^{i\kappa\tau(\hat{p}_x a_0^2/\hbar)\hat{z}}, \quad (2)$$

which matches the desired form (1) with $\alpha = \kappa\tau a_0^2$. The prefactor $\hat{U}_x^2$ just reverses the spatial coordinate $x \rightarrow -x$ and adds an unimportant phase [21]. Finally, we measure the x position and p_z momentum of the particle in this final state, which we denote by $x_m \equiv -x$ (to remove the spatial reversal) and $p_m \equiv p_z/\alpha$ (to recover p_x by removing the scaling by α). The measurement of p_z can be conveniently achieved by applying a quarter-period harmonic oscillator in z simultaneously with the other directions and measuring in position space. The final positions can be imaged by capture in a lattice, where for a reasonable matter-wave magnification, the discrete position measurement from the lattice will not be limiting. The phase space can then be read out via an image of the x - z plane.

Following this protocol, the probability of measuring the particle in the state (x_m, p_m) is readily found to be [22]

$$P(x_m, p_m) = \left| \langle \psi_{\text{coh}}^{(x_m, p_m)} | \psi_i \rangle \right|^2, \quad (3)$$

where $\psi_{\text{coh}}^{(x_m, p_m)}(x) \equiv e^{-(x-x_m)^2/2\sigma_c^2 + ip_m x/\hbar}/(\pi\sigma_c^2)^{1/4}$ is the phase-space coherent state centered on (x_m, p_m) with positional uncertainty $\Delta x = \sigma_c = \alpha\sigma$. Equation (3) defines the Husimi- Q representation of the quantum state: it is related to the Wigner function by a convolution that renders it a positive-definite measure in phase space [23]. Equation (3) shows that our protocol is a POVM in the coherent state basis [19]. Indeed, it implements precisely the Arthurs-Kelly proposal for measurement of noncommuting variables [18]. Similar POVMs using two noncommuting spin observables were previously performed in trapped ions [24] and for collective observables [25], while Wigner functions of individual atoms were previously obtained by motional state tomography in tight harmonic traps [26,27]. Our Letter goes beyond these by being applicable to systems of very general form.

Joint measurements of conjugate variables [18] leave freedom to choose how to distribute quantum noise between the two variables while remaining consistent with the uncertainty principle. In our protocol, the uncertainty of the measured momentum is set to be $\Delta p_m \sim \hbar/(\alpha\sigma)$ by the size σ of $\phi_0(z)$ and the scaling α in Eq. (1). Reducing this must cause the measurement to add noise to Δx_m [18]. Here this arises because the potential $V_{\text{kick}}(x, z)$, applied in the Fourier plane, imparts an impulse $\hbar\kappa\tau z$ in x due to the necessarily bent force lines, with uncertainty $\sim \hbar\kappa\tau\sigma$ due to the spread σ in z . Moving back to real space using $\hat{U}_x$, this introduces the noise $\Delta x_m \sim (\hbar\kappa\tau\sigma)(a_0^2/\hbar) = \alpha\sigma$.

One can readily generalize this procedure [22], extending the system of interest to 2D $\mathbf{r} = (x, y)$ and imprinting a general $\hat{U}_{\text{kick}}^G = e^{iG(\hat{\mathbf{p}})\hat{z}}$, where $G(\mathbf{p})$ is a chosen scalar function of $\mathbf{p} = (p_x, p_y)$. This generalization requires an isotropic harmonic oscillator with equal frequencies in x and y , as well as imaging in a three-dimensional quantum gas microscope. Note also that the full phase space is four dimensional, so it cannot be fully encoded in these three spatial dimensions with the scalar function $G(\mathbf{p})$. One can instead use an internal spin degree of freedom as an additional auxiliary coordinate, which allows the mapping of both components of the 2D momentum onto the Bloch sphere, analogous to a stereographic projection (see Fig. 2 and [22]). Finally, although we have focused on a single particle, the method can be extended to multiparticle systems of bosons or fermions, provided interactions are sufficiently weak that scattering can be neglected during the quarter-period evolutions. It also admits extensions to spinful systems, provided internal levels can be separately measured.




FIG. 2. Sketches of protocols to map moments of the momentum density to an auxiliary spin state in averaged-mode phase-space microscopes. These are applied in the Fourier plane, so the position is labeled by momentum $\mathbf{p}$. (a) Spin rotation (i) transforms the initial $|S, S\rangle$ spin state (aligned vectors of lower layer) using a resonant Rabi pulse that circulates as $\vec{\Omega} \propto (-p_y, p_x, 0)$ (orange central layer) to map the 2D momentum space onto the Bloch sphere in the final state (vectors of upper layer), in a mapping analogous to a stereographic projection (see End Matter). (b) Spin rotation (ii) uses a uniform Rabi pulse $\vec{\Omega} \propto (1, 0, 0)$ in the presence of a quadratically varying Zeeman shift to imprint $|\mathbf{p}|^2$ in S_z . In both cases, the measured spin encodes moments of the momentum distribution.

We refer to the above protocol as a “*Husimi- Q phase-space microscope*” since it measures the wave function in phase space according to Eq. (3). We use this term broadly to include the situations in which, jointly with the atomic positions, the microscope images only a subset of the momentum components or some function of momentum.

The Husimi- Q framework also reveals a natural path to a second class of measurement. The joint position-momentum measurement has position uncertainty σ_c and momentum uncertainty $\hbar/\sigma_c$. When $\sigma_c \rightarrow 0$, such that the position becomes sharply resolved, one might expect that momentum information is lost. However, this is not the case. Instead, even if one cannot resolve the full distribution of momentum, one can still measure phase-space averages of the momentum density or its moments. For example, in the above 1D Husimi- Q microscope, if instead of measuring the momentum in the z direction one measures the probability to occupy the n th harmonic oscillator eigenstate, one finds this to be proportional to the $2n$ th moment of the local momentum distribution, even as $\sigma_c \rightarrow 0$. The resulting “*averaged-mode phase-space microscopes*” are functions only of the spatial coordinates and are not subject to Heisenberg noise.

Averages in phase space are readily obtained when one makes the auxiliary space finite: either as a finite set of states of a harmonic trap or as a spin of finite length. Consider the auxiliary space to be an internal spin S with the system initially prepared in the maximally polarized spin state, i.e., $|\Psi_i\rangle = |\psi_i\rangle \otimes |S, S\rangle$. We encode information about the system state $|\psi_i\rangle$ in the spin using a momentum-dependent

spin rotation, $\hat{U}_{\text{rot}} = \exp[-i\vec{\theta}(\hat{\mathbf{p}}) \cdot \hat{\mathbf{S}}]$. One can again do this by transforming to the Fourier plane by quarter-period evolutions in both x and y (via the product of operators $\hat{U}_x \hat{U}_y$), and then imposing a spatially dependent spin rotation $\vec{\theta}(\mathbf{r})$, before transforming back to position space with $\hat{U}_x \hat{U}_y$. Here we provide two example protocols to measure different moments of the momentum distribution.

Spin rotation (i)—We map the 2D momentum to the surface of the Bloch sphere by a spin rotation $\vec{\theta}(\mathbf{p}) = 2(\mp p_y, p_x, 0)/p_0$, where p_0 sets a characteristic momentum scale [28]. In the matter-wave Fourier plane (where $\mathbf{p}$ is transcribed to $\mathbf{r}$), this requires a resonant Rabi field $\vec{\Omega}(\mathbf{p}) \propto \vec{\theta}(\mathbf{p})$ that is circulating in the plane, Fig. 2(a). Such a spin rotation could be performed via an optical two-photon transition using Gauss-Laguerre beams with angular momenta $L = 0$ and $L = \pm 1$ [29]. Overall, this couples space and spin as

$$|\Psi_f\rangle = \hat{U}_y^2 \hat{U}_x^2 e^{-2i\frac{(\hat{p}_x \hat{S}_y \mp \hat{p}_y \hat{S}_x)}{p_0}} |\psi_i\rangle \otimes |S, S\rangle, \quad (4)$$

where $\hat{U}_y^2 \hat{U}_x^2$ effects the inversion $\mathbf{r} \rightarrow -\mathbf{r}$. This protocol can, in principle, be used to construct a POVM for the full four-dimensional phase space of (x, y) and (p_x, p_y) [22].

Here we show that projective measurements of the spin components extract moments of the momentum density. Consider projective measurements of $\hat{S}_z$. Expanding the exponential (4) for small $|\hat{\mathbf{p}}|/p_0$ gives $|\Psi_f\rangle \propto \hat{U}_y^2 \hat{U}_x^2 \sum_{j=0}^2 S \sqrt{\binom{2S}{j}} (\hat{p}_x \pm i\hat{p}_y/p_0)^j |\psi_i\rangle \otimes |S, S-j\rangle$. The particle is found in the $|S, S-1\rangle$ state at position $\mathbf{r}_m = -\mathbf{r}$ (chosen to correct for the spatial inversion) with probability density proportional to $|\langle \mathbf{r}_m | (\hat{p}_x \pm i\hat{p}_y) | \psi_i \rangle|^2 \equiv 2M [\mathcal{K} \mp \hbar \mathcal{M}]$, where

$$\mathcal{K}(\mathbf{r}_m) \equiv \frac{\hbar^2}{2M} |\nabla \psi_i(\mathbf{r}_m)|^2, \quad (5)$$

$$\mathcal{M}(\mathbf{r}_m) \equiv \frac{\hbar}{2iM} \nabla \psi_i^* \times \nabla \psi_i \cdot \hat{\mathbf{z}} \quad (6)$$

are the kinetic energy density [31] and the orbital magnetization (the curl of the current density), respectively. Thus, repeating measurements for $\Delta L = \pm 1$ allows both the kinetic energy and orbital magnetization densities to be measured. Similarly, the particle is in the $|S, S-j\rangle$ state with probability density $\propto |\langle \mathbf{r}_m | (\hat{p}_x \pm i\hat{p}_y)^j | \psi_i \rangle|^2$. Thus, the measurement of all spin states provides $2S$ moments of the momentum distribution as well as the particle density ($j = 0$). Measurements of $\hat{S}_{x,y}$ provide the local momentum density [22].

Spin rotation (ii)—In place of the circulating Rabi field, one can engineer a useful momentum-dependent coupling

using rf pulses. We impose a quadratically varying Zeeman shift in the Fourier plane, as realized in a magnetic trap [Fig. 2(b)], leading to a detuning $\delta \equiv \Omega|\mathbf{p}|^2/p_0^2$ when the rf is resonant at $|\mathbf{p}|^2 = 0$. We apply a *uniform* Rabi field $\Omega\hat{S}_x$ for a time π/Ω , which rotates a particle at $|\mathbf{p}|^2 = 0$ into $|S, -S\rangle$, in general in a multiphoton transition. A particle at $|\mathbf{p}|^2 \neq 0$ will not perform an exact π rotation, leading to

$$|\Psi_f\rangle = \hat{U}_y^2 \hat{U}_x^2 \exp[\pi i(\hat{S}_x - \hat{S}_z|\hat{\mathbf{p}}|^2/p_0^2)]|\psi_i\rangle \otimes |S, S\rangle. \quad (7)$$

Expanding in $|\hat{\mathbf{p}}|^2/p_0^2$ shows that the amplitude of $|S, -S + j\rangle$ grows as $|\hat{\mathbf{p}}|^{2j}$. In particular, the particle is in the $|S, -S + 1\rangle$ state at position $\mathbf{r}_m = -\mathbf{r}$, with probability proportional to

$$|\langle \mathbf{r}_m | |\hat{\mathbf{p}}|^2 | \psi_i \rangle|^2 = |\hbar^2 \nabla^2 \psi_i(\mathbf{r}_m)|^2, \quad (8)$$

providing a quartic moment of the momentum density for $|\mathbf{p}| \ll p_0$.

Thus, measurements of $\hat{S}_z$ return spatial maps of moments of the momentum density. Larger values of S allow access to higher moments. However, $S = 1/2$ is already sufficient for measuring the kinetic energy density and orbital magnetization [Eqs. (5), (6)] with spin rotation (i), or the quartic moment of momentum [Eq. (8)] with spin rotation (ii). As mentioned above, there is no fundamental limit to the spatial resolution of these averaged-mode measurements since they do not measure separate non-commuting operators. Rather, they measure certain momentum-dependent observables. For an N -particle system, the observables Eqs. (5), (6), and (8) correspond to

$$\hat{\rho}_{\text{KE}}(\mathbf{r}) \equiv \sum_{i=1}^N \sum_{\alpha=x,y} \frac{1}{2M} \hat{p}_i^\alpha \delta(\mathbf{r} - \hat{\mathbf{r}}_i) \hat{p}_i^\alpha, \quad (9)$$

$$\hat{\rho}_{\text{mag}}(\mathbf{r}) \equiv \nabla \times \sum_{i=1}^N \frac{1}{2} \left[\frac{\hat{\mathbf{p}}_i}{M}, \delta(\mathbf{r} - \hat{\mathbf{r}}_i) \right]_+ \cdot \hat{\mathbf{z}}, \quad (10)$$

$$\hat{\rho}_{\text{quartic}}(\mathbf{r}) \equiv \sum_{i=1}^N |\hat{\mathbf{p}}_i|^2 \delta(\mathbf{r} - \hat{\mathbf{r}}_i) |\hat{\mathbf{p}}_i|^2, \quad (11)$$

where $[\hat{A}, \hat{B}]_+$ denotes the anticommutator of $\hat{A}$ and $\hat{B}$. Averaged-mode phase-space microscopes allow measurements of these new observables and their correlations, in addition to the particle density $\hat{\rho}(\mathbf{r}) \equiv \sum_{i=1}^N \delta(\mathbf{r} - \hat{\mathbf{r}}_i)$. In effect, by making the derivatives of the wave function observables, they provide continuum realizations of protocols for measuring off-diagonal coherences on lattices, which can measure local currents and kinetic energies [11,32,33]. The freedom to control the evolution in the Fourier plane provides a very rich set of possible momentum-based observables to be realized using averaged-mode phase-space microscopes.

In practice, the spatial resolution of the measurement process will be limited by atomic shot noise. For a 2D system with mean density $\bar{\rho}$, measurements of density $\hat{\rho}(\mathbf{r})$ with spatial resolution w have an error $\delta\rho/\bar{\rho} \sim 1/\sqrt{\bar{\rho}w^2}$ due to the projection noise from the atom number in that area. Similarly, measurements of the kinetic energy density $\hat{\rho}_{\text{KE}}(\mathbf{r})$ for particles with characteristic wavelength λ will have an error $\delta\rho_{\text{KE}}/\bar{\rho}_{\text{KE}} \sim 1/\sqrt{(\bar{\rho}/\lambda^2)(\hbar^2/p_0^2)w^2}$ because the typical density of spin flips is $(\bar{\rho}/\lambda^2)(\hbar^2/p_0^2)$. Thus, to have a signal with minimal noise, one should choose $p_0 \sim \hbar/\lambda$. Note that the above protocols (4) and (7) approximate the densities (9), (10), and (11) accurately for $|\mathbf{p}| \ll p_0$, with p_0 acting as a high-momentum cutoff.

The proposed phase-space microscopes could be directly implemented in state-of-the-art quantum gas microscopes. They would significantly expand the toolbox of cold atom experiments: by exposing momentum-space features with spatial resolution, they open up many new ways to probe quantum many-body systems. Here we focus on weakly trapped gases where the continuum momentum is the relevant quantity. However, the microscope can also be used for lattice-based systems, where it can be helpful to use band mapping to recover the lattice quasimomentum [34]. In the following, we present four applications in continuum settings.

(1) Short-range external potentials: Imagine that one wishes to characterize a sharp potential step imposed at the edge of a system, leading to a wave function $\psi_{\text{edge}}(x)$ that vanishes over a small length scale δ . Measurement of the particle density $|\psi_{\text{edge}}(x)|^2$ with a point spread function in position of w can resolve down to $\delta \sim w$. However, a Husimi- Q phase-space microscope with the same positional resolution $\sigma_c = w$, evaluated at the edge, can detect a high momentum tail out to $\sim \hbar/\delta$, even for $\delta \ll w$ (see Fig. 3). While a global measurement of the momentum distribution would also depend on the sharpness of the




FIG. 3. Measurement of a sharp step in the potential, with edge thickness δ mimicked by a state $\psi_{\text{edge}}(x) = [1 - \tanh(x/\delta)]/2$. (a) Density distribution as measured by a conventional QGM with point spread function $\propto e^{-x^2/w^2}$. (b) Distribution of particles with momentum $p_m = 3\hbar/\sigma_c$ as measured by a Husimi- Q phase-space microscope with $\sigma_c = w$. The density measurement (a) becomes insensitive to δ for $\delta/w \lesssim 0.5$, while the phase-space measurement (b) continues to discriminate via the high-momentum tail.

edge, the phase-space measurement targets the local properties and can lead to a larger signal-to-noise ratio.

(2) Imaging quantum vortices: Averaged-mode phase-space microscopes allow access to the velocity field associated with the circulation around a vortex. They can be used to image the kinetic energy and magnetization densities (9) and (10), which have peaks at the vortex core, or the quartic density (11), which is a ring around the vortex core, Fig. 4. These features have sizes set by the healing length ξ , so sensitivity is maximized for $p_0 \sim \hbar/\xi$. The measurements could also be performed in 3D, using spin as the auxiliary space. Then, the paths of the vortices will appear as lines of spin flips along their cores. This may facilitate measurements of quantum turbulence [35].

(3) Thermometry: Measurements of the local kinetic energy allow access to the local temperature. This could facilitate thermometry of strongly interacting systems—e.g., by partitioning the cloud into a region of interest and a region in which the equation of state is well known. For noninteracting bosons in 2D, a thermal cloud with density $\bar{\rho}$ has a kinetic energy density $\bar{\rho}_{\text{KE}} = k_B T \bar{\rho}$, while a BEC has $\bar{\rho}_{\text{KE}} = [\pi M / 12 \hbar^2] (k_B T)^2$, both providing simple scales. (See Ref. [22] for other cases.) The protocol would also allow a better fit to bimodal profiles in a partially condensed cloud by including spatial position and kinetic energy for each particle. Such precise determination of the condensate fraction is, e.g., important for studying thermal condensate fluctuations [37–39]. The local nature also opens up the possibility of measuring temperature gradients, or indeed measuring second sound.

(4) Local Tan contact: Quite generally, for a gas with short-range interactions, the high-momentum population takes the form $\rho_p \sim C \hbar^2 / |p|^4$. Here, C is the Tan contact, which characterizes the interacting system [40]. Depending on spatial dimension and on which moment of the momentum is measured, this can dominate the signal in an averaged-mode phase-space microscope. For example,




FIG. 4. Particle density, kinetic energy density, magnetization density, and quartic momentum density around a 2D vortex of healing length ξ , computed for the ansatz [36] $\psi(r, \theta) = \sqrt{\bar{\rho}} e^{i\theta} / \sqrt{r^2 + 2\xi^2}$. The averaged-mode phase-space microscope also allows measurements of the latter three quantities.

for 2D systems, the local quartic density Eq. (11) is dominated by this contribution when $p_0 \gtrsim \hbar\sqrt{\bar{\rho}}$ [41]. Using the protocol (7) for $S = 1/2$, one finds $\langle \hat{\rho}_{\text{quartic}} \rangle = 0.14 \hbar^2 C p_0^2$. While C has been measured by other methods [42–45], the averaged-mode phase-space microscope permits measurements even in spatially inhomogeneous systems, e.g., probing its variation in vortex cores even for strongly interacting superfluids. More generally, the dependence of $\langle \hat{\rho}_{\text{quartic}} \rangle$ on p_0 would allow a detailed analysis of condensate depletion in such inhomogeneous settings.

Beyond the applications outlined above, phase-space microscopes offer even more powerful ways to probe strong correlated phases through measurements of spatial correlations of momentum-space features [34]. In the lowest Landau level regime, correlations of density and velocity have been shown to provide a way to measure the many-body interaction energy [46]. We expect that correlations in phase space also have the potential to reveal hidden order in exotic many-body phases, potentially in combination with machine learning analysis of the data [47]. The access to conjugate observables with detection of every particle might be useful for accessing entanglement structure [48] similar to the measurement of noncommuting spin observables [49]. Squeezing between particles in a quantum many-body system could directly show in the phase-space microscope via a narrower distribution in one of the conjugate variables of position and momentum.

Acknowledgments—We thank Ulrich Schneider and Zoran Hadzibabic for helpful comments on the manuscript. This work was funded by EPSRC (Grants No. EP/V062654/1 and No. EP/Y01510X/1), by a Simons Investigator Award (Grant No. 511029), and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) via Research Unit FOR 5688, Project No. 521530974.

Data availability—The data that support the findings of this article are openly available [56].

- [1] C. Gross and W. S. Bakr, *Nat. Phys.* **17**, 1316 (2021).
- [2] W. S. Bakr, A. Peng, M. E. Tai, R. Ma, J. Simon, J. I. Gillen, S. Fölling, L. Pollet, and M. Greiner, *Science* **329**, 547 (2010).
- [3] J. F. Sherson, C. Weitenberg, M. Endres, M. Cheneau, I. Bloch, and S. Kuhr, *Nature (London)* **467**, 68 (2010).
- [4] M. Holten, L. Bayha, K. Subramanian, C. Heintze, P. M. Preiss, and S. Jochim, *Phys. Rev. Lett.* **126**, 020401 (2021).
- [5] T. de Jongh, J. Verstraten, M. Dixmerias, C. Daix, B. Peaudecerf, and T. Yefsah, *Phys. Rev. Lett.* **134**, 183403 (2025).
- [6] J. Xiang, E. Cruz-Colón, C. C. Chua, W. R. Milner, J. de Hond, J. F. Fricke, and W. Ketterle, *Phys. Rev. Lett.* **134**, 183401 (2025).

- [7] R. Yao, S. Chi, M. Wang, R. J. Fletcher, and M. Zwiernlein, *Phys. Rev. Lett.* **134**, 183402 (2025).
- [8] M. Endres, M. Cheneau, T. Fukuhara, C. Weitenberg, P. Schauß, C. Gross, L. Mazza, M. C. Bañuls, L. Pollet, I. Bloch, and S. Kuhr, *Science* **334**, 200 (2011).
- [9] T. A. Hilker, G. Salomon, F. Grusdt, A. Omran, M. Boll, E. Demler, I. Bloch, and C. Gross, *Science* **357**, 484 (2017).
- [10] J. Koepsell, J. Vijayan, P. Sompet, F. Grusdt, T. A. Hilker, E. Demler, G. Salomon, I. Bloch, and C. Gross, *Nature (London)* **572**, 358 (2019).
- [11] A. Impertro, S. Karch, J. F. Wienand, S. J. Huh, C. Schweizer, I. Bloch, and M. Aidelsburger, *Phys. Rev. Lett.* **133**, 063401 (2024).
- [12] A. Impertro, S. Huh, S. Karch, J. F. Wienand, I. Bloch, and M. Aidelsburger, *Nat. Phys.* **21**, 895 (2025).
- [13] J. C. Brüggengjürgen, M. S. Fischer, and C. Weitenberg, *Science* **393**, 167 (2026).
- [14] L. Asteria, H. P. Zahn, M. N. Kosch, K. Sengstock, and C. Weitenberg, *Nature (London)* **599**, 571 (2021).
- [15] S. Brandstetter, C. Heintze, P. Hill, P. M. Preiss, M. Gałka, and S. Jochim, *Phys. Rev. Lett.* **135**, 103401 (2025).
- [16] P. A. Murthy and S. Jochim, [arXiv:1911.10824](https://arxiv.org/abs/1911.10824).
- [17] C. Weitenberg, L. Asteria, O. Carlsson, A. Bohrdt, and F. Grusdt, [arXiv:2602.12142](https://arxiv.org/abs/2602.12142).
- [18] E. Arthurs and J. L. Kelly, Jr., *Bell Syst. Tech. J.* **44**, 725 (1965).
- [19] A. Peres, *Quantum Theory: Concepts and Methods*, Fundamental Theories of Physics (Kluwer Academic Publishers, Dordrecht, 1995), Vol. 57.
- [20] This can be achieved via the coil geometry or a current imbalance, with the 1D system oriented at 45 degrees to the coil axis.
- [21] If the second quarter-period oscillator has a different length a'_0 , the transformation also provides an overall magnification $(a'_0/a_0)^2$.
- [22] See End Matter.
- [23] U. Leonhardt, *Essential Quantum Optics: From Quantum Measurements to Black Holes* (Cambridge University Press, Cambridge, England, 2010).
- [24] M. A. Rowe, D. Kielpinski, V. Meyer, C. A. Sackett, W. M. Itano, C. Monroe, and D. J. Wineland, *Nature (London)* **409**, 791 (2001).
- [25] P. Kunkel, M. Prüfer, S. Lannig, R. Rosa-Medina, A. Bonnín, M. Gärtner, H. Strobel, and M. K. Oberthaler, *Phys. Rev. Lett.* **123**, 063603 (2019).
- [26] J. F. Kanem, S. Maneshi, S. H. Myrskog, and A. M. Steinberg, *J. Opt. B* **7**, S705 (2005).
- [27] M. O. Brown, S. R. Muleady, W. J. Dworschack, R. J. Lewis-Swan, A. M. Rey, O. Romero-Isart, and C. A. Regal, *Nat. Phys.* **19**, 569 (2023).
- [28] At small momenta, this is an approximation to the stereographic projection discussed in the End Matter.
- [29] Despite the near detuning required for alkali atoms, the losses should be small, as one only needs to make a small fractional rotation of the levels, as in Ref. [30].
- [30] L. W. Cheuk, A. T. Sommer, Z. Hadzibabic, T. Yefsah, W. S. Bakr, and M. W. Zwiernlein, *Phys. Rev. Lett.* **109**, 095302 (2012).
- [31] The kinetic energy density can be written as $(-\hbar^2/2M)\psi(\mathbf{r}^*)\nabla^2\psi(\mathbf{r})$ or $(+\hbar^2/2M)|\nabla\psi(\mathbf{r})|^2$, which gives the same total kinetic energy under integration when ψ vanishes on the boundary.
- [32] M. C. Tran, D. K. Mark, W. W. Ho, and S. Choi, *Phys. Rev. X* **13**, 011049 (2023).
- [33] M. Kiser, M. McGinley, and D. Malz, *Phys. Rev. A* **114**, 013311 (2026).
- [34] Y. Yang, C. Weitenberg, and N. R. Cooper (to be published).
- [35] N. Navon, A. L. Gaunt, R. P. Smith, and Z. Hadzibabic, *Nature (London)* **539**, 72 (2016).
- [36] A. L. Fetter, *Rev. Mod. Phys.* **81**, 647 (2009).
- [37] M. A. Kristensen, M. B. Christensen, M. Gajdacz, M. Iglicki, K. Pawłowski, C. Klempt, J. F. Sherson, K. Rzażewski, A. J. Hilliard, and J. J. Arlt, *Phys. Rev. Lett.* **122**, 163601 (2019).
- [38] M. B. Christensen, T. Vibel, A. J. Hilliard, M. B. Kruk, K. Pawłowski, D. Hryniuk, K. Rzażewski, M. A. Kristensen, and J. J. Arlt, *Phys. Rev. Lett.* **126**, 153601 (2021).
- [39] Z. Jalali-Mola, N. Käming, L. Asteria, U. Bhattacharya, R. W. Chhajlany, K. Sengstock, M. Lewenstein, T. Grass, and C. Weitenberg, *Phys. Rev. Lett.* **136**, 083401 (2026).
- [40] S. Tan, *Ann. Phys. (Amsterdam)* **323**, 2971 (2008).
- [41] For a weakly interacting 2D BEC, $C \sim (\tilde{g}\tilde{\rho})^2$, so it contributes $\langle\hat{p}_{\text{quartic}}\rangle \sim \tilde{g}^2\tilde{\rho}^2 p_0^2/\hbar^2$. For $\tilde{\rho}p_0^2/\hbar^2 \gg 1$, this is larger than the contribution from the condensate $\langle\hat{p}_{\text{quartic}}\rangle \sim \tilde{\rho}(\hbar/\xi)^4$ expected in the vicinity of a vortex or edge, with healing length $\xi \sim 1/\sqrt{\tilde{g}\tilde{\rho}}$.
- [42] J. T. Stewart, J. P. Gaebler, T. E. Drake, and D. S. Jin, *Phys. Rev. Lett.* **104**, 235301 (2010).
- [43] E. D. Kuhnle, H. Hu, X.-J. Liu, P. Dyke, M. Mark, P. D. Drummond, P. Hannaford, and C. J. Vale, *Phys. Rev. Lett.* **105**, 070402 (2010).
- [44] Y.-Q. Zou, B. Bakkali-Hassani, C. Maury, É. Le Cerf, S. Nascimbène, J. Dalibard, and J. Beugnon, *Nat. Commun.* **12**, 760 (2021).
- [45] Q. Huang, H. Yao, X. Chen, and L. Sanchez-Palencia, *Sci. Adv.* **11**, eadv3727 (2025).
- [46] D. Spasic-Mlacak and N. R. Cooper, *Phys. Rev. Research* **7**, L032032 (2025).
- [47] G. Carleo, I. Cirac, K. Cranmer, L. Daudet, M. Schuld, N. Tishby, L. Vogt-Maranto, and L. Zdeborová, *Rev. Mod. Phys.* **91**, 045002 (2019).
- [48] R. Islam, R. Ma, P. M. Preiss, M. Eric Tai, A. Lukin, M. Rispoli, and M. Greiner, *Nature (London)* **528**, 77 (2015).
- [49] P. Kunkel, M. Prüfer, S. Lannig, R. Strohmaier, M. Gärtner, H. Strobel, and M. K. Oberthaler, *Phys. Rev. Lett.* **128**, 020402 (2022).
- [50] O. Elíasson, J. S. Laustsen, R. Heck, R. Müller, J. J. Arlt, C. A. Weidner, and J. F. Sherson, *Phys. Rev. A* **102**, 053311 (2020).
- [51] J. Koepsell, S. Hirthe, D. Bourgund, P. Sompet, J. Vijayan, G. Salomon, C. Gross, and I. Bloch, *Phys. Rev. Lett.* **125**, 010403 (2020).
- [52] T. Legrand, F.-R. Winkelmann, W. Alt, D. Meschede, A. Alberti, and C. A. Weidner, *Phys. Rev. A* **109**, 033304 (2024).
- [53] J. M. Radcliffe, *J. Phys. A Gen. Phys.* **4**, 313 (1971).
- [54] T. Chalopin, T. Satoor, A. Evrard, V. Makhalov, J. Dalibard, R. Lopes, and S. Nascimbene, *Nat. Phys.* **16**, 1017 (2020).
- [55] M. Saffman and K. Mølmer, *Phys. Rev. A* **78**, 012336 (2008).
- [56] N. R. Cooper, Y. Yang, and C. Weitenberg, Supporting Data, Apollo Repository, [10.17863/CAM.133500](https://doi.org/10.17863/CAM.133500).

End Matter

General analysis for auxiliary spatial coordinate—We consider a particle in an initial state $\psi_i(\mathbf{r})$ within the plane described by $\mathbf{r} = (x, y)$ and a state $\phi_0(z)$ along the perpendicular direction z . We encode the momentum $\mathbf{p} = (p_x, p_y)$ in the z momentum by applying a generalized kick,

$$\hat{U}_{\text{kick}}^G = e^{iG(\hat{\mathbf{p}})\hat{z}}, \quad (\text{A1})$$

where $G(\mathbf{p})$ is a scalar function.

We write the initial state of the particle,

$$|\Psi_i\rangle = \int d^2\mathbf{r} dz \Psi_i(\mathbf{r}, z) |\mathbf{r}, z\rangle, \quad (\text{A2})$$

with $\Psi_i(\mathbf{r}, z) = \psi_i(\mathbf{r})\phi_0(z)$. We retain (weak) confinement in z and apply a harmonic trap to both x and y for a $T/4$ pulse. We denote the oscillator length of this trap by a_0 such that this pulse transforms the wave function via the operator,

$$\hat{U}_x = \exp[-i(\hat{p}_x^2 a_0^2 / \hbar^2 + \hat{x}^2 / a_0^2) \pi / 2]. \quad (\text{A3})$$

Using the fact that $\hat{U}_x = \sum_n |n\rangle\langle n| \exp[-i(n + 1/2)\pi/2]$ and the fact that the oscillator states $|n\rangle$ take the same functional form in both position and momentum space, up to scaling, one finds that the application of quarter-period

pulses to both x and y produces the “Fourier plane” for the $\mathbf{r} = (x, y)$ coordinates, where $|\Psi_{\text{Fourier}}\rangle = \hat{U}_x \hat{U}_y |\Psi_i\rangle$. In this Fourier plane, the position $\mathbf{r}$ encodes the momentum $\mathbf{p}$ of the initial state. Formally, this arises from $\hat{U}_x^{-1} \hat{x} \hat{U}_x = -\hat{p}_x a_0^2 / \hbar$ and $\hat{U}_x^{-1} \hat{p}_x \hat{U}_x = \hat{x} \hbar / a_0^2$.

We now apply a kick in the z direction with a strength that depends on the position $\mathbf{r}$. To match the desired form (A1), this is

$$|\Psi'_{\text{Fourier}}\rangle = \exp[-iG(\hat{\mathbf{r}}\hbar/a_0^2)\hat{z}] \hat{U}_x \hat{U}_y |\Psi_{\text{initial}}\rangle. \quad (\text{A4})$$

We transform back from the Fourier plane to real space by applying the $T/4$ evolution in x, y again. Then,

$$|\Psi_f\rangle = \hat{U}_x \hat{U}_y \exp[iG(\hat{\mathbf{r}}\hbar/a_0^2)\hat{z}] \hat{U}_x \hat{U}_y |\Psi_{\text{initial}}\rangle \quad (\text{A5})$$

$$= -\hat{R}_x \hat{R}_y \exp[i\hat{z}G(\hat{\mathbf{p}})] |\Psi_{\text{initial}}\rangle, \quad (\text{A6})$$

where we have used $\hat{U}_x^2 = (-i)\hat{R}_x$, with $\hat{R}_x$ being the reflection operator, taking $|x\rangle \rightarrow |-x\rangle$.

Finally, we measure the particle to be at position $\mathbf{r}_f$ in the x - y plane and in the momentum state $p_{z,f}$ along z . (The measurement of momentum in z is conveniently performed by converting p_z to position z_f via $\hat{U}_z$, and then measuring z_f .) Introducing a complete set of states in the x - y plane, the probability of detecting the particle at $\mathbf{r}_f$ and $p_{z,f}$ is

$$|\langle \mathbf{r}_f; p_{z,f} | \Psi_f \rangle|^2 = \frac{1}{(2\pi\hbar)^3} \left| \int d^2\mathbf{p} d^2\mathbf{r} dz e^{-i\mathbf{p} \cdot (\mathbf{r} + \mathbf{r}_f) / \hbar} e^{i[G(\mathbf{p}) - p_{z,f}/\hbar]z} \Psi_{\text{initial}}(\mathbf{r}, z) \right|^2. \quad (\text{A7})$$

This general expression (A7) allows one to analyze the consequences of different kicks G in the Fourier plane. The example used in the main text is the 1D phase space (x, p) with $G(p) = \alpha p / \hbar$. The final coordinates $(x_f, p_{z,f})$ used in Eq. (A7) are expressed in terms of $x_m \equiv -x_f$ and $p_m = p_{z,f}/\alpha$ to correct for the spatial inversion and the scaling of momentum with α . Using the 1D version of Eq. (A7), this leads to the following probability:

$$\begin{aligned} P(x_m, p_m) &= \frac{\alpha}{(2\pi\hbar)^2} \left| \int d\mathbf{p} dx dz e^{i(z\alpha + x_m - x)p/\hbar - i\alpha p_m z/\hbar} \psi_i(x) \phi_0(z) \right|^2 \\ &= \left| \int dx e^{-i(x - x_m)p_m/\hbar} \frac{e^{-\frac{(x - x_m)^2}{2\alpha^2 \sigma^2}}}{(\pi\alpha^2 \sigma^2)^{1/4}} \right|^2. \end{aligned} \quad (\text{A8})$$

For 2D systems, one can envisage encoding any scalar function of $\mathbf{p}$ in the transverse momentum, for example, using $G(\mathbf{p}) \propto |\mathbf{p}|^2$. The final imaging could be performed via a spatial tomography consisting of successive images of different planes [50,51] or other techniques such as helical point spread function engineering [52].

General analysis for auxiliary spin S —The spin rotation (i) used in the main text (4) is an approximation to the stereographic projection of the 2D momentum to spin-coherent states on the Bloch sphere.

We consider an internal spin S . A stereographic projection that maps from the 2D momentum (p_x, p_y) to the

surface of a sphere (n_x, n_y, n_z) with $|\vec{n}|^2 = 1$ is described by

$$\frac{n_x + in_y}{1 - n_z} = \frac{p_x + ip_y}{p_0}, \quad (\text{B1})$$

where p_0 is a characteristic momentum scale such that $|\mathbf{p}| = p_0$ is mapped to the equator, $n_z = 0$. The key step in the mapping is to perform a unitary transformation that rotates the initial fully spin-aligned state $|S, +S\rangle$ to the spin-coherent state that is aligned with this $\vec{n}$. This is achieved by choosing a rotation $\hat{\theta}(\mathbf{p})$ such that

$$\hat{U}_{\text{rot}} = \exp[2i\hat{S} \cdot (\hat{p}_y, -\hat{p}_x, 0)/|\hat{\mathbf{p}}| \arctan(|\hat{\mathbf{p}}|/p_0)]. \quad (\text{B2})$$

A state of definite momentum (p_x, p_y) is mapped to the coherent spin state $|\chi_{\mathbf{p}}\rangle \propto \hat{U}_{\text{rot}}(\mathbf{p})|S, S\rangle$. For small momenta $|\mathbf{p}| \ll p_0$, we have $\arctan(|\hat{\mathbf{p}}|/p_0) \simeq |\hat{\mathbf{p}}|/p_0$ to get

$$|\chi_{\mathbf{p}}\rangle \simeq \exp[2i\hat{S} \cdot \mathbf{e}_z \times (\hat{p}_y, -\hat{p}_x, 0)/p_0]|S, S\rangle, \quad (\text{B3})$$

which is the form used in the main text.

The spin-coherent states are [53]

$$|\eta\rangle = \frac{1}{(1 + |\eta|^2)^S} \exp(\eta \hat{S}_-)|S, S\rangle, \quad (\text{B4})$$

with $\eta \equiv \eta_x + i\eta_y$ being a complex number. Like phase-space coherent states, spin-coherent states provide an

overcomplete basis. As such, one can imagine performing POVMs in the $|\eta\rangle$ basis. To do so would require the spin- S to be mapped onto another (higher dimensional) Hilbert space, with a spin with $S' \gg S$, in the spirit of Ref. [25].

Imagine that one were to perform the POVM in spin-coherent states. The outcome would be to measure the particle at position $\mathbf{r}_f$ and spin-coherent state $|\eta_f\rangle$ with probability $P(\mathbf{r}_f, \eta_f) \equiv |\langle \mathbf{r}_f, \eta_f | \Psi_f \rangle|^2$, where

$$\langle \mathbf{r}_f, \eta_f | \Psi_f \rangle \propto \int d^2p d^2r e^{-i\mathbf{p} \cdot (\mathbf{r} + \mathbf{r}_f)} \Psi_i(\mathbf{r}) \langle \eta_f | \chi_{\mathbf{p}} \rangle. \quad (\text{B5})$$

Using

$$\langle \eta_f | \chi_{\mathbf{p}} \rangle = \frac{(1 + \eta_f^* \chi_{\mathbf{p}})^{2S}}{(1 + |\eta_f|^2)^S (1 + |\chi_{\mathbf{p}}|^2)^S} \quad (\text{B6})$$

and assuming $|\eta_f|, |\chi_{\mathbf{p}}| \ll 1$, we obtain

$$\langle \eta_f | \chi_{\mathbf{p}} \rangle \simeq \exp[-S(|\eta_f|^2 + |\chi_{\mathbf{p}}|^2 - 2\eta_f^* \chi_{\mathbf{p}})], \quad (\text{B7})$$

with $\chi_{\mathbf{p}} \simeq (p_x + ip_y)/p_0$. The integral over momentum in Eq. (B5) can then be completed to give

$$P(\mathbf{r}_f, \eta_f) = |\langle \mathbf{r}_f, \eta_f | \Psi_f \rangle|^2 = \left| \langle \psi_{\text{coh}}^{(\mathbf{r}_f, \eta_f)} | \psi_i \rangle \right|^2, \quad (\text{B8})$$

where the position representation of the state $|\psi_{\text{coh}}^{(\mathbf{r}_f, \eta_f)}\rangle$ is

$$\langle \mathbf{r} | \psi_{\text{coh}}^{(\mathbf{r}_f, \eta_f)} \rangle = \frac{p_0}{\sqrt{2\pi S \hbar}} e^{i(\eta_{f,x}, \eta_{f,y}) \cdot (\mathbf{r} - \mathbf{r}_f) p_0 / \hbar} e^{-\frac{1}{4S} \{ [(x - x_f) p_0 / \hbar + 2S\eta_{f,y}]^2 + (y - y_f) p_0 / \hbar - 2S\eta_{f,x} \}^2}. \quad (\text{B9})$$

This is a coherent state of the *two-dimensional* phase space, centered on the position $(x_f - 2S\eta_{f,y}\hbar/p_0, y_f + 2S\eta_{f,x}\hbar/p_0)$ and momentum $(p_x, p_y) = p_0(\eta_{f,x}, \eta_{f,y})$. Thus, the measurement is a POVM in the full 2D phase space and would realize a 2D Husimi- Q phase-space microscope. The positional uncertainty is controlled by $\sigma'_c \equiv \sqrt{2S\hbar}/p_0$ and the momentum uncertainty by $\hbar/\sigma'_c$. For fixed characteristic momentum p_0 , one achieves increasingly accurate resolution in momentum by increasing the size of the spin S . A proper choice of the atomic species would allow large spin manifolds such as 17 states in dysprosium [54] or 23 states in holmium [55]. For Dy ($S = 8$), this gives $\sigma'_c = 4\hbar/p_0$ and, for Ho ($S = 11$), $\sigma'_c \approx 4.7\hbar/p_0$.

Even without full POVMs on the spin S , this mapping of momentum to spin can be readily used to construct an averaged-mode phase-space microscope using standard projective measurements of the spin components $\hat{S}_{x,y,z}$. Since $|\chi_{\mathbf{p}}\rangle$ is the spin-coherent state aligned along $\vec{n} \simeq (p_x/p_0, p_y/p_0, 1 - |\mathbf{p}|^2/(2p_0^2))$, one has

$$\langle \chi_{\mathbf{p}} | \hat{S} | \chi_{\mathbf{p}} \rangle \simeq S(p_x/p_0, p_y/p_0, 1 - |\mathbf{p}|^2/(2p_0^2)). \quad (\text{B10})$$

Measurements of the expectation values of $\hat{S}_x$ and $\hat{S}_y$ provide information on the mean momentum densities, while the expectation value of $\hat{S}_z$ gives the mean kinetic energy density as discussed in the main text.

Thermodynamics of the weakly interacting BEC—For a weakly interacting Bose gas in 2D, the low-temperature thermodynamics is dominated by Bogoliubov sound mode with $E_{\mathbf{p}} = s|\mathbf{p}|$. The speed of sound is $s = \sqrt{\tilde{g}\bar{\rho}(\hbar/M)}$, with $\tilde{g}$ being the dimensionless coupling constant, which is $\tilde{g} = \sqrt{8\pi a_s/\ell_z}$ for atoms of scattering length a_s confined to 2D by an oscillator state of size ℓ_z . Applying Bogoliubov theory, one finds

$$\bar{\rho}_{\text{KE}}(T) = \bar{\rho}_{\text{KE}}(0) + \frac{\zeta(3)}{2\pi\hbar^2 s^2} (k_B T)^3, \quad (\text{C1})$$

with $\zeta(3) \simeq 1.202$. The zero-temperature value $\bar{\rho}_{\text{KE}}(0)$ has a divergent contribution $\propto C \ln(p_{\text{max}})$, arising from the contact [40], which sets the large momentum tail of the momentum distribution, $\rho_p \sim C\hbar^2/p^4$. The momentum cutoff p_{max} is set by nonuniversal physical effects: the ramping times in the experimental dynamics [42] or, as

discussed in the main text, the characteristic scale p_0 up to which the quadratic expansion of the observable is valid. This generates a nonuniversal background value for $\bar{\rho}_{\text{KE}}$. However, the temperature can be extracted from the change $\Delta\bar{\rho}_{\text{KE}} = \bar{\rho}_{\text{KE}}(T) - \bar{\rho}_{\text{KE}}(0)$, with $T \propto (\Delta\bar{\rho}_{\text{KE}})^{1/3}$.