

Topological Arrest of Ballooning Modes in Nonaxisymmetric Toroidal Plasmas

Amitava Bhattacharjee^{*}

Department of Astrophysical Sciences, Princeton University, Princeton, New Jersey 08540, USA



(Received 4 February 2026; accepted 12 June 2026; published 1 September 2026)

Why do nonaxisymmetric stellarators avoid ballooning crashes that afflict tokamaks? Three-dimensional geometry induces Anderson localization of ballooning modes, converting a global instability into a Ginzburg-Landau network of isolated wave packets. Global stability reduces to a percolation problem: Below a critical threshold, instability is arrested; above it, a crash occurs. This explains benign stellarator saturation, predicts vulnerability in quasisymmetric designs, and introduces the critical threshold as a nonlinear stability metric for reactor optimization, pending experimental validation.

DOI: [10.1103/6m2z-wlmb](https://doi.org/10.1103/6m2z-wlmb)

Introduction—The pursuit of high-pressure plasma confinement in toroidal devices is constrained by ballooning instabilities, which can drive edge-localized modes (ELMs) in tokamaks [1] and profile-limiting density collapses in stellarators [2]. A puzzling asymmetry has emerged: Tokamaks are susceptible to ELMs that expel significant heat fluxes to the wall, while highly nonaxisymmetric stellarators such as Large Helical Device (LHD) [3] and Wendelstein 7-X (W7-X) [4] operate safely within linearly unstable regimes. Understanding this contrast is a central open problem in fusion plasma physics.

Ballooning modes are characterized by long-wavelength structures aligned with field lines and short-wavelength structures perpendicular to them. Cuthbert and Dewar [5] showed that, in general toroidal geometry, ballooning eigenfunctions are exponentially localized along field lines and interpreted this behavior in terms of Anderson localization [6], a well-known phenomenon in condensed matter physics. Redi *et al.* [7] subsequently demonstrated that stellarator ballooning spectra exhibit signatures of quantum chaos. These results suggest that three-dimensional (3D) geometry fundamentally alters the structure of ballooning modes.

In this Letter, we propose that Anderson localization converts weakly nonlinear ballooning dynamics into a Ginzburg-Landau system on a sparse network of localized structures and that global stability is governed by the connectivity of this network. We identify a dimensionless percolation parameter η and argue that global instability can be avoided when $\eta < \eta_c = 1.128$ [8]. While the framework

requires quantitative experimental validation, it suggests that magnetic aperiodicity can serve as a topological safety net against global ballooning crashes.

Anderson localization—In the high-toroidal-mode-number limit, the linearized ballooning equation can be written as [5,9]

$$\frac{d}{d\theta} \left(A(\theta) \frac{d\xi}{d\theta} \right) - [K(\theta) + \lambda N(\theta)] \xi = 0, \quad (1)$$

where ξ is related to the normal component of the plasma displacement, $A(\theta)$, $K(\theta)$, and $N(\theta)$ are functions of the coordinate θ along a field line, and λ is the linear eigenfrequency squared. Because $A > 0$, after a standard Liouville transformation, this Sturm-Liouville problem can be transformed into Schrödinger form:

$$\left[\frac{d^2}{d\theta^2} + E - V(\theta) \right] y = 0, \quad (2)$$

where $y(\theta) = A^{1/2}(\theta)\xi(\theta)$. The effective potential $V(\theta)$ depends on equilibrium properties sampled along the field line, and E is the “energy” eigenvalue. Our notation is identical to that of [5].

Axisymmetric versus 3D geometry—In both cases, the potential contains a secular magnetic shear term growing as $|\theta| \rightarrow \infty$ that is not periodic. The crucial distinction lies in the behavior of the geometric coefficients. For axisymmetry, the curvature, metric coefficients, magnetic field strength, and pressure-gradient drive are periodic functions of θ : A field line samples the same geometric sequence repeatedly. The resulting eigenfunctions are spatially extended Bloch solutions [5,9].

In 3D geometry, by contrast, different field lines on the same flux surface sample *different* sequences of curvature and metric coefficients. Along a single field line, these coefficients vary aperiodically as the line winds around the

^{*}Contact author: amitava@princeton.edu

torus. This aperiodic geometric sampling—not the shear term—is what produces Anderson localization.

Transfer-matrix analysis and the Lyapunov exponent—Discretizing Eq. (2) on a uniform grid $\theta_n = n\Delta\theta$ gives the transfer-matrix form

$$\begin{pmatrix} y_{n+1} \\ y_n \end{pmatrix} = T_n \begin{pmatrix} y_n \\ y_{n-1} \end{pmatrix}, \quad (3)$$

where

$$T_n = \begin{pmatrix} 2 + (\Delta\theta)^2[V(\theta_n) - E] & -1 \\ 1 & 0 \end{pmatrix}. \quad (4)$$

The Lyapunov exponent of the transfer-matrix product is

$$\gamma = \lim_{N \rightarrow \infty} \frac{1}{N} \ln \|T_N T_{N-1} \cdots T_1\|, \quad (5)$$

where $\|\cdot\|$ denotes any convenient matrix norm (the limit is norm independent). For 1D Schrödinger operators with aperiodic coefficients, products of transfer matrices generically satisfy $\gamma > 0$ —this is guaranteed under mild assumptions by the Furstenberg theorem [10]. A positive Lyapunov exponent implies exponential localization of eigenfunctions:

$$|y_n| \lesssim e^{-|n-n_0|/\ell}, \quad (6)$$

where n_0 is the localization center and $\ell \equiv \gamma^{-1}$ is the localization length. The inequality in Eq. (6) should be read as a smooth upper-bound envelope; the eigenfunctions computed numerically by Cuthbert and Dewar exhibit significant fine structure within the envelope [5].

Furstenberg-type arguments apply under specific assumptions: A given stellarator field line may be quasiperiodic or nearly integrable, in which case γ should be computed directly from the transfer matrix. In particular, quasiperiodic potentials—which may arise in quasisymmetric stellarators where $|B|$ has an approximate continuous symmetry—can support extended or critical states rather than Anderson-localized ones.

The Ginzburg-Landau network—We derive a time-dependent weakly nonlinear form of the ideal-magnetohydrodynamics (MHD) equations by expanding the fully nonlinear ideal-MHD displacement equation in Lagrangian coordinates [11–13]. The plasma position is $\mathbf{r}(\mathbf{r}_0, t) = \mathbf{r}_0 + \boldsymbol{\xi}(\mathbf{r}_0, t)$, where $\boldsymbol{\xi}$ is the Lagrangian displacement. Expanding in powers of $\boldsymbol{\xi}$ yields

$$\partial_t \boldsymbol{\xi} = \mathcal{L}\boldsymbol{\xi} + \mathcal{N}_2(\boldsymbol{\xi}, \boldsymbol{\xi}) + \mathcal{N}_3(\boldsymbol{\xi}, \boldsymbol{\xi}, \boldsymbol{\xi}), \quad (7)$$

where $\mathcal{L}$ is the linearized ideal-MHD operator and $\mathcal{N}_2$ and $\mathcal{N}_3$ are quadratic and cubic nonlinearities, respectively. The reduction from the second-order ideal-MHD system to the

first-order form (7) follows from the fact that near marginal ballooning stability the unstable eigenvalues are purely real and growing, so the center manifold is one-dimensional and the dynamics project onto a nonoscillatory equation for each mode amplitude [14]. Near marginal stability, the perturbation on this reduced manifold is

$$\xi(\theta, t) = \sum_j a_j(t) \phi_j(\theta), \quad (8)$$

where $\phi_j(\theta)$ are the weakly unstable and Anderson-localized eigenfunctions. As derived in the Appendix and End Matter, the amplitude equations take the Ginzburg-Landau form

$$\dot{a}_j = \gamma_j a_j - \mu_j a_j^3 + \sum_{k \neq j} J_{jk} a_k, \quad (9)$$

where the linear growth rate γ_j (not to be confused with the Lyapunov exponent) and the cubic coefficient $\mu_j > 0$ are mode-specific. The intermode coupling decays exponentially with separation:

$$J_{jk} \sim J_0 e^{-d_{jk}/\ell}. \quad (10)$$

Here, each localized packet is labeled by a field-line label α_j and a ballooning-space localization center θ_j ; its projection onto the flux surface defines a point x_j , and d_{jk} is the metric distance on the surface. The suppression of the quadratic term $\mathcal{N}_2$ from Eq. (9) is a central result whose derivation is given in End Matter.

The Ginzburg-Landau network (9) naturally contains the intermediate nonlinear regime first identified by Zhu, Bhattacharjee, and Germaschewski [15] and subsequently developed for toroidal ballooning modes [13,16]. Prior to cubic saturation, when $\mu_j a_j^2 \ll \gamma_j$, each mode amplitude continues to grow exponentially at the linear rate γ_j with its spatial structure unchanged. In 3D geometry, the intermediate regime is followed by cubic saturation. In axisymmetric geometry, sufficiently close to marginal stability, the quadratic drive [12] can, in principle, follow the intermediate regime and produce explosive growth; whether it does so in practice depends on proximity to marginality and the absence of competing saturation mechanisms.

Two localized structures interact effectively only when the coupling J_{jk} exceeds a characteristic dynamical scale $\sim \gamma_j$, the linear growth rate of the mode. This defines an effective interaction radius

$$R_* = \ell \ln \left(\frac{J_0}{\gamma_j} \right). \quad (11)$$

Here, R_* is determined entirely by equilibrium and linear stability properties with no free parameters. On a flux surface, the centers θ_j of unstable localized packets are




FIG. 1. Percolation realizations of the three topological connectivity regimes on a toroidal flux surface. Each disk represents a localized ballooning-mode packet; disk radii equal the interaction radius R_* , and two packets are linked when their disks overlap (centers within $2R_*$). The surface-spanning cluster is shown in red. (a) Subcritical ($\eta \ll \eta_c$): high magnetic aperiodicity produces short localization lengths and sparse, nonpercolating clusters; global synchronization is prevented and instability saturates benignly. (b) Near-critical ($\eta \approx \eta_c = 1.128$): a surface-spanning cluster path (highlighted) intermittently forms, enabling bursty, intermittent MHD activity. (c) Supercritical ($\eta \gg \eta_c$): the spanning cluster persists, phase locking modes globally and enabling the explosive growth characteristic of a major profile crash.

distributed with areal density $\rho(\beta)$, which increases as the pressure gradient exceeds the marginal value. The set of packets and their interaction radii defines a random geometric graph on the flux surface. A standard result of continuum percolation theory [8] is that such a graph develops a surface-spanning connected cluster when the dimensionless parameter

$$\eta(\beta) = \rho(\beta)\pi R_*^2(\beta) \quad (12)$$

exceeds the critical value $\eta_c = 1.128$. Recent M3D-C¹ simulations of W7-X plasmas show regimes of benign saturation ($\beta < 5\%$) and crashlike behavior in configurations with near-zero magnetic shear [17,18]. In the present framework, these behaviors correspond naturally to subcritical and supercritical connectivity of localized ballooning structures.

We note that this percolation threshold provides a necessary but not sufficient condition for global instability. The present framework treats local growth rates as approximately uniform and coupling phases as random. In practice, heterogeneity in growth rates could produce mode clustering that locally restores quadratic coupling even in a globally subcritical network. Conversely, phase interference among coupling terms could raise the effective threshold above η_c . A fully quantitative stability criterion would require accounting for these effects, which we leave for future work. The percolation threshold on the toroidal flux surface (topologically a 2-torus) differs from the flat-plane value by finite-size corrections that are small for large N . Furthermore, if the interaction region is anisotropic (elongated along field lines), the relevant percolation threshold for elliptical objects differs from 1.128, though the qualitative phase-transition structure is preserved.

To connect η with experiment and simulation, Fig. 1 shows the topological connectivity for three characteristic magnetic geometries, and Fig. 2 provides both a schematic regime map and the theoretical scaling.




FIG. 2. (a) Schematic estimates of the connectivity parameter $\eta = \rho\pi R_*^2$ for representative operating regimes. Bands reflect order-of-magnitude uncertainty; no published study directly reports ρ or R_* in the percolation sense. Device positions are based on the qualitative reasoning described in the text: W7-X (M3D-C¹ simulations show benign saturation to $\beta \approx 5\%$ [17]); LHD (bursty MHD activity and density crashes near the ballooning threshold [2]); DIII-D inter-ELM (radial correlation lengths 0.5–1 cm from reflectometry and BES [19–21], filamentary scales from [22,23]). Dashed line: continuum percolation threshold $\eta_c = 1.128$ [8]. (b) Theoretical scaling $\eta \sim (\ell/a)^2 [\ln(J_0/\gamma)]^2$ for three values of the dimensionless coupling ratio. Device positions are qualitative estimates of ℓ/a ; axisymmetric geometry ($\ell \rightarrow \infty$, $a/\ell \rightarrow 0$) gives $\eta \rightarrow \infty$.

For the schematic regime map (Fig. 2, left panel), device positions are based on qualitative reasoning from published results rather than direct extraction of ρ and R_* . W7-X simulations show benign nonlinear saturation up to $\beta \approx 5\%$ [17], consistent with $\eta \ll \eta_c$; LHD exhibits bursty MHD activity and core density collapse events near its ballooning threshold [2], consistent with $\eta \lesssim \eta_c$; and DIII-D inter-ELM measurements of radial correlation lengths of density fluctuations (0.5–1 cm) from reflectometry and beam emission spectroscopy (BES) diagnostics [19–21], combined with filamentary scales in W7-X and LHD [22,23], provide proxy estimates placing DIII-D in a moderately supercritical regime. We repeat for emphasis that no published study directly reports ρ or R_* in the percolation sense; the values in Fig. 2 are order-of-magnitude estimates with uncertainty bands spanning roughly one order of magnitude. The right panel in Fig. 2 shows the theoretical scaling $\eta \sim (\ell/a)^2 [\ln(J_0/\gamma)]^2$, derived in End Matter, as a function of the ratio a/ℓ of machine size to localization length, with device positions placed at qualitative estimates of this ratio. This scaling makes explicit that axisymmetric geometry ($\ell \rightarrow \infty$, $a/\ell \rightarrow 0$) formally gives $\eta \rightarrow \infty$ for any finite β above marginal stability.

A quantitative determination of η would require (i) computing γ (and, hence, ℓ) from the transfer-matrix Eq. (3) evaluated along field lines of each equilibrium reconstruction; (ii) measuring $\rho = N/A$ from the linear stability boundary as a function of field-line label α , i.e., the fraction of field lines on the flux surface that are ballooning unstable at a given β ; and (iii) determining R_* via Eq. (11) with γ_j from linear stability calculations. Steps (i) and (ii) are computable from equilibrium reconstructions and would convert Fig. 2 from a schematic regime map into a parameter-free theoretical prediction testable against simulation. Step (iii) could be validated experimentally from spatiotemporal correlation functions of coherent fluctuation structures in imaging or probe data.

Conclusions—We have demonstrated that 3D geometry converts ballooning dynamics from a single global mode into a Ginzburg-Landau system on a network of Anderson-localized states. Nonlinear stability is, therefore, a connectivity property of this network. Localization is a *consequence* of the Furstenberg theorem applied to the aperiodic ballooning potential in 3D geometry. The Anderson mechanism is absent in axisymmetric geometry, where extended Bloch eigenfunctions support the explosive quadratic nonlinearity of the detonation theory [12].

In the presence of symmetry breaking imposed by resonant magnetic perturbations in a tokamak, the framework predicts that sufficient magnetic aperiodicity could push η below η_c , providing a topological interpretation of ELM suppression. This is a prediction of the theory amenable to experimental test.

Recent work suggests that in quasisymmetric equilibria the magnetic field strength along a flux surface may possess an underlying structure associated with integrable systems such

as the Korteweg–de Vries equation [24]. The spectrum of the associated Schrödinger-like operator for Korteweg–de Vries-type (reflectionless) potentials consists of extended, transmitting states rather than Anderson-localized ones—precisely the case where the Furstenberg theorem does not apply. This connection strengthens the conjecture that stellarators with near-perfect quasisymmetry may prove as vulnerable to nonlinear ballooning modes as axisymmetric tokamaks. Recent M3D-C¹ simulations of Landreman-Buller-Drevlak-type quasisymmetric equilibria showing sustained ballooning instability [18] seem qualitatively consistent with this prediction and motivate dedicated linear calculations of the Lyapunov exponent for these configurations.

Density crashes have been observed in LHD by Ohdachi *et al.* [2], showing a ballooning mode in the gradient region driving a beta-limiting collapse. This provides possible evidence for the Anderson-localization mechanism: It is spatially localized, is destabilized by 3D geometry, and appears in a region of globally negative magnetic shear where axisymmetric ballooning theory predicts stability. Within the present framework, this mode is a localized packet that survives stabilizing global shear because the aperiodic 3D potential creates an isolated potential well along a specific field line—a phenomenon that cannot occur in axisymmetric geometry. The intermittent, bursty character of the density collapse is consistent with the near-critical regime $\eta \approx \eta_c$, in which the spanning cluster forms and dissolves repeatedly as the pressure profile oscillates about the percolation threshold.

The percolation parameter η could serve as an additional optimization target in large-scale stellarator design codes alongside existing neoclassical and linear-MHD metrics—providing a nonlinear stability criterion that is, in principle, computable from equilibrium and linear stability data. By strategically retaining a degree of geometric aperiodicity, it may be possible to design reactors that are topologically immune to global ballooning crashes, trading near-perfect quasisymmetry for robust nonlinear stability.

Acknowledgments—We thank N. Bohlsen, A. Brown, S. Buller, I. Dodin, P. Helander, N. Nikulsin, H. Qin, and W. Sengupta for helpful discussions. We acknowledge the use of large language models [Anthropic (Claude), ChatGPT, and Gemini] for assistance with drafting, organizing ideas, and navigating the literature. The author is responsible for all scientific content and conclusions. This research was supported by a grant from the Simons Foundation/SFARI (Grant No. 560651, A. B.) and DOE Award No. DE-SC0024548. We are grateful to Princeton University for supporting our research.

Data availability—There are no publicly available research data or software supporting this manuscript. Requests for further information or data should be sent to the author.

- [1] E. Strait, *Phys. Plasmas* **1**, 1415 (1994).
- [2] S. Ohdachi *et al.*, *Nucl. Fusion* **57**, 066042 (2017).
- [3] H. Yamada *et al.*, *Fusion Sci. Technol.* **51**, 12 (2010).
- [4] O. Grulke *et al.*, *Nucl. Fusion* **64**, 112002 (2024).
- [5] P. Cuthbert and R. L. Dewar, *Phys. Plasmas* **7**, 2302 (2000).
- [6] P. W. Anderson, *Phys. Rev.* **109**, 1492 (1958).
- [7] M. H. Redi, J. L. Johnson, S. Klasky, J. Canik, R. L. Dewar, and W. A. Cooper, *Phys. Plasmas* **9**, 1990 (2002).
- [8] J. Quintanilla, S. Torquato, and R. M. Ziff, *J. Phys. A* **33**, L399 (2000).
- [9] R. L. Dewar and A. H. Glasser, *Phys. Fluids* **26**, 3038 (1983).
- [10] A. Crisanti, G. Paladin, and A. Vulpiani, *Products of Random Matrices in Statistical Physics* (Springer-Verlag, Berlin, 1993).
- [11] D. Pfirsch and R. N. Sudan, *Phys. Fluids B* **5**, 2052 (1993).
- [12] S. C. Cowley and M. Artun, *Phys. Rep.* **283**, 185 (1997).
- [13] P. Zhu, C. C. Hegna, C. R. Sovinec, A. Bhattacharjee, and K. Germaschewski, *Phys. Plasmas* **14**, 055903 (2007).
- [14] J. D. Crawford, *Rev. Mod. Phys.* **63**, 991 (1991).
- [15] P. Zhu, A. Bhattacharjee, and K. Germaschewski, *Phys. Rev. Lett.* **96**, 065001 (2006).
- [16] P. Zhu, C. C. Hegna, and C. R. Sovinec, *Phys. Rev. Lett.* **102**, 235003 (2009).
- [17] Y. Zhou, K. Aleynikova, C. Liu, and N. M. Ferraro, *Phys. Rev. Lett.* **133**, 135102 (2024).
- [18] A. M. Wright and N. M. Ferraro, *Phys. Plasmas* **31**, 082509 (2024).
- [19] Z. Yan, G. R. McKee, R. J. Groebner, P. B. Snyder, T. H. Osborne, M. N. Beurskens, and K. H. Burrell, *Phys. Plasmas* **18**, 056117 (2011).
- [20] T. L. Rhodes, R. J. Taylor, and W. A. Peebles, *Rev. Sci. Instrum.* **66**, 824 (1995).
- [21] K. Barada *et al.*, *Nucl. Fusion* **61**, 126037 (2021).
- [22] A. Buzas *et al.*, *Nucl. Fusion* **64**, 066012 (2020).
- [23] K. Tanaka, N. Ohno, Y. Tsuji, S. Kajita, S. Masuzaki, M. Kobayashi, T. Morisaki, A. Komori, and the LHD Experimental Group, *Plasma Fusion Res.* **7**, 1402152 (2012).
- [24] W. Sengupta, R. Madan, S. Buller, N. Nikulsin, E. J. Paul, R. Nies, A. A. Kaptanoglu, S. R. Hudson, and A. Bhattacharjee, *Phys. Plasmas* **32**, 102509 (2025).

End Matter

Appendix A: Derivation of the Ginzburg-Landau equation—We describe how the weakly nonlinear Ginzburg-Landau evolution follows from the fully nonlinear Lagrangian ideal-MHD displacement equation. In Lagrangian variables, ideal MHD admits an exact equation of motion [11–13]

$$\rho_0(\mathbf{r}_0)\partial_t^2\xi = \mathbf{F}[\xi], \quad (\text{A1})$$

where $\mathbf{F}[\xi]$ is a nonlinear force functional. Expanding about equilibrium gives

$$\mathbf{F}[\xi] = \mathcal{L}\xi + \mathcal{Q}(\xi, \xi) + \mathcal{C}(\xi, \xi, \xi) + \dots, \quad (\text{A2})$$

where $\mathcal{L}$ is the linear ideal-MHD force operator and $\mathcal{Q}$ and $\mathcal{C}$ are quadratic and cubic nonlinearities, respectively. Near marginal ballooning stability, the spectrum of $\mathcal{L}$ contains a finite set of weakly unstable purely real eigenvalues. The corresponding unstable center manifold is 1D, and standard center-manifold analysis [14] reduces Eq. (A1) to the first-order system (7). Projecting onto the adjoint eigenfunctions $\phi_j^\dagger$ yields the amplitude equations

$$\dot{a}_j = \gamma_j a_j + \sum_{kl} c_{j;kl}^{(2)} a_k a_l + \sum_{klm} c_{j;klm}^{(3)} a_k a_l a_m + \dots, \quad (\text{A3})$$

where $c_{j;kl}^{(2)} = \langle \phi_j^\dagger, \mathcal{Q}(\phi_k, \phi_l) \rangle$ and analogously for $c^{(3)}$. The suppression of the quadratic coupling in 3D geometry is demonstrated below. Retaining the leading contributions gives Eq. (9) with $J_{jk} \sim e^{-d_{jk}/\ell}$.

Appendix B: Suppression of quadratic couplings—Localization does not make $c_{j;kl}^{(2)}$ vanish identically, but it does make most couplings exponentially small. If packet envelopes satisfy $|\phi_j| \lesssim e^{-d(\theta, \theta_j)/\ell}$, then

$$|c_{j;kl}^{(2)}| \lesssim C_2 e^{-D_{jkl}/\ell}, \quad (\text{B1})$$

where D_{jkl} measures the mutual separation of the three packets. Off-diagonal quadratic couplings are, therefore, exponentially suppressed in the strongly localized regime.

The self-coupling $k = l = j$ is not suppressed by overlap. In stellarator-symmetric equilibria, where $V(\theta)$ is even about the localization center θ_j , the ground-state eigenfunction is even: $\phi_j = f_j(\theta - \theta_j)$, $f_j(-u) = f_j(u)$, $f_j \rightarrow 0$ exponentially. Writing $A f_j^2 f_j' = (\frac{1}{3}) A (f_j^3)'$ and integrating by parts (boundary term vanishing as $f_j \rightarrow 0$) $c_{j;jj}^{(2)} = -\frac{1}{3} \int A' f_j^3 d\theta \sim O(\ell/L_A)$. Here, L_A is the variation scale of A , and $b_j \equiv c_{j;jj}^{(2)} \sim O(\ell/L_A)$ is the self-quadratic coefficient. Neglecting it relative to the cubic requires $|b_j| \ll \mu_j \sqrt{\gamma_j/\mu_j}$ at the saturation amplitude $a_j \sim \sqrt{\gamma_j/\mu_j}$.

The key distinction from axisymmetric geometry is *phase coherence*. In axisymmetric geometry, the Bloch eigenfunction $\phi \sim P(\theta) e^{-i\frac{1}{2}\theta^2/2\ell_s^2}$ carries a periodic phase structure that supports a two-step harmonic cascade: The primary mode at m drives a second harmonic at $2m$, which feeds back to drive the detonation mechanism [12]. Anderson localization destroys this extended phase coherence, converting the quadratic interaction into short-ranged incoherent couplings.

For quasirandomly distributed centers, the N terms in Q_j contribute with quasirandom signs. By the Brownian-walk argument, $|Q_j| \sim \sqrt{N}$ while the collective cubic scales as N :

$$\frac{|Q_j|}{\sum_j \mu_j a_j^3} \sim \frac{1}{\sqrt{N}} \xrightarrow{N \rightarrow \infty} 0. \quad (\text{B2})$$

The cubic Ginzburg-Landau network (9) is, therefore, the correct leading dynamics in the large- N , strongly localized regime.

Appendix C: Percolation parameter scaling—With $\rho = f(\beta)/a^2$ (number of packets $\sim f$ over machine area a^2)

and $R_* = \ell \ln(J_0/\gamma)$,

$$\eta = \rho \pi R_*^2 \sim \pi f \left(\frac{\ell}{a} \right)^2 \ln^2 \left(\frac{J_0}{\gamma} \right). \quad (\text{C1})$$

As $\ell \rightarrow \infty$ (axisymmetric limit), $\eta \rightarrow \infty$ for any $\beta > \beta_c$, consistent with the explosive character of tokamak ballooning instabilities and with Fig. 2(b). Near marginal β_c , $f \sim \epsilon \equiv (\beta - \beta_c)/\beta_c$, giving critical overshoot $\epsilon_c \sim a^2 \eta_c / [\pi \ell^2 \ln^2(J_0/\gamma)]$; for $\ell/a \sim 0.1$ – 0.3 and $\ln(J_0/\gamma) \sim 2$ – 3 , $\epsilon_c \sim 0.03$ – 0.10 , consistent with observed ballooning-limit windows.