

First Measurement of Charged-Current Muon-Neutrino-Induced K^+ Production on Argon Using the MicroBooNE Detector

P. Abratenko,³⁹ D. Andrade Aldana,¹⁴ L. Arellano,²² J. Asaadi,³⁸ A. Ashkenazi,³⁷ S. Balasubramanian,¹² B. Baller,¹² A. Barnard,²⁹ G. Barr,²⁹ D. Barrow,²⁹ J. Barrow,²⁶ V. Basque,¹² J. Bateman,^{15,22} O. Benevides Rodrigues,¹⁴ S. Berkman,²⁵ A. Bhat,⁷ M. Bhattacharya,¹² M. Bishai,³ A. Blake,¹⁹ B. Bogart,²⁴ T. Bolton,¹⁸ M. B. Brunetti,^{17,41} L. Camilleri,¹⁰ D. Caratelli,⁴ F. Cavanna,¹² G. Cerati,¹² A. Chappell,⁴¹ Y. Chen,³³ J. M. Conrad,²³ M. Convery,³³ L. Cooper-Troendle,³⁰ J. I. Crespo-Anadón,⁶ R. Cross,⁴¹ M. Del Tutto,¹² S. R. Dennis,⁵ P. Detje,⁵ R. Diurba,² Z. Djurcic,¹ K. Duffy,²⁹ S. Dytman,³⁰ B. Eberly,³⁵ P. Englezos,³² A. Ereditato,^{7,12} J. J. Evans,²² C. Fang,⁴ G. A. Fiorentini Aguirre,³⁴ W. Foreman,^{14,20} B. T. Fleming,⁷ D. Franco,⁷ A. P. Furmanski,²⁶ F. Gao,⁴ D. Garcia-Gamez,¹³ S. Gardiner,¹² G. Ge,¹⁰ S. Gollapinni,²⁰ E. Gramellini,²² P. Green,²⁹ H. Greenlee,¹² L. Gu,¹⁹ W. Gu,³ R. Guenette,²² P. Guzowski,²² L. Hagaman,⁷ M. D. Handley,⁵ O. Hen,²³ C. Hilgenberg,²⁶ G. A. Horton-Smith,¹⁸ A. Hussain,¹⁸ B. Irwin,²⁶ M. S. Ismail,³⁰ C. James,¹² X. Ji,²⁷ J. H. Jo,³ R. A. Johnson,⁸ D. Kalra,¹⁰ G. Karagiorgi,¹⁰ W. Ketchum,¹² M. Kirby,³ T. Kobilarcik,¹² N. Lane,^{15,22} J.-Y. Li,¹¹ Y. Li,³ K. Lin,³² B. R. Littlejohn,¹⁴ L. Liu,¹² W. C. Louis,²⁰ X. Luo,⁴ T. Mahmud,¹⁹ C. Mariani,⁴⁰ D. Marsden,²² J. Marshall,⁴¹ N. Martinez,¹⁸ D. A. Martinez Caicedo,³⁴ S. Martynenko,³ A. Mastbaum,³² I. Mawby,¹⁹ N. McConkey,³¹ V. Meddage,¹⁸ L. Mellet,²⁵ J. Mendez,²¹ J. Micallef,^{23,39} A. Mogan,⁹ T. Mohayai,¹⁶ M. Mooney,⁹ A. F. Moor,⁵ C. D. Moore,¹² L. Mora Lepin,²² M. M. Moudgalya,²² S. Mullerlababu,² D. Naples,³⁰ A. Navrer-Agasson,¹⁵ N. Nayak,³ M. Nebot-Guinot,¹¹ C. Nguyen,³² J. Nowak,¹⁹ N. Oza,¹⁰ O. Palamara,¹² N. Pallat,²⁶ V. Paolone,³⁰ A. Papadopoulou,¹ V. Papavassiliou,²⁸ H. B. Parkinson,¹¹ S. F. Pate,²⁸ N. Patel,¹⁹ Z. Pavlovic,¹² E. Piasetzky,³⁷ K. Pletcher,²⁵ I. Pophale,¹⁹ X. Qian,³ J. L. Raaf,¹² V. Radeka,³ A. Rafique,¹ M. Reggiani-Guzzo,¹¹ J. Rodriguez Rondon,^{16,34} M. Rosenberg,³⁹ M. Ross-Lonergan,²⁰ I. Safa,¹⁰ D. W. Schmitz,⁷ A. Schukraft,¹² W. Seligman,¹⁰ M. H. Shaevitz,¹⁰ R. Sharankova,¹² J. Shi,⁵ E. L. Snider,¹² M. Soderberg,³⁶ S. Söldner-Rembold,¹⁵ J. Spitz,²⁴ M. Stancari,¹² J. St. John,¹² T. Strauss,¹² A. M. Szelc,¹¹ N. Taniuchi,⁵ K. Terao,³³ C. Thorpe,²² D. Torbunov,³ D. Totani,⁴ M. Troups,¹² A. Trettin,²² Y.-T. Tsai,³³ J. Tyler,¹⁸ M. A. Uchida,⁵ T. Usher,³³ B. Viren,³ J. Wang,²⁷ M. Weber,² H. Wei,²¹ A. J. White,⁷ S. Wolbers,¹² T. Wongjirad,³⁹ K. Wresilo,⁵ W. Wu,³⁰ E. Yandel,^{4,20} T. Yang,¹² L. E. Yates,¹² H. W. Yu,³ G. P. Zeller,¹² J. Zennamo,¹² and C. Zhang³

(MicroBooNE Collaboration)*

¹Argonne National Laboratory (ANL), Lemont, Illinois 60439, USA

²Universität Bern, Bern CH-3012, Switzerland

³Brookhaven National Laboratory (BNL), Upton, New York 11973, USA

⁴University of California, Santa Barbara, California 93106, USA

⁵University of Cambridge, Cambridge CB3 0HE, United Kingdom

⁶Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), Madrid E-28040, Spain

⁷University of Chicago, Chicago, Illinois 60637, USA

⁸University of Cincinnati, Cincinnati, Ohio 45221, USA

⁹Colorado State University, Fort Collins, Colorado 80523, USA

¹⁰Columbia University, New York, New York 10027, USA

¹¹University of Edinburgh, Edinburgh EH9 3FD, United Kingdom

¹²Fermi National Accelerator Laboratory (FNAL), Batavia, Illinois 60510, USA

¹³Universidad de Granada, Granada E-18071, Spain

¹⁴Illinois Institute of Technology (IIT), Chicago, Illinois 60616, USA

¹⁵Imperial College London, London SW7 2AZ, United Kingdom

¹⁶Indiana University, Bloomington, Indiana 47405, USA

¹⁷The University of Kansas, Lawrence, Kansas 66045, USA

¹⁸Kansas State University (KSU), Manhattan, Kansas 66506, USA

¹⁹Lancaster University, Lancaster LA1 4YW, United Kingdom

²⁰Los Alamos National Laboratory (LANL), Los Alamos, New Mexico 87545, USA

²¹Louisiana State University, Baton Rouge, Louisiana 70803, USA

²²The University of Manchester, Manchester M13 9PL, United Kingdom

²³Massachusetts Institute of Technology (MIT), Cambridge, Massachusetts 02139, USA

²⁴University of Michigan, Ann Arbor, Michigan 48109, USA

²⁵Michigan State University, East Lansing, Michigan 48824, USA

- ²⁶University of Minnesota, Minneapolis, Minnesota 55455, USA
²⁷Nankai University, Nankai District, Tianjin 300071, China
²⁸New Mexico State University (NMSU), Las Cruces, New Mexico 88003, USA
²⁹University of Oxford, Oxford OX1 3RH, United Kingdom
³⁰University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
³¹Queen Mary University of London, London E1 4NS, United Kingdom
³²Rutgers University, Piscataway, New Jersey 08854, USA
³³SLAC National Accelerator Laboratory, Menlo Park, California 94025, USA
³⁴South Dakota School of Mines and Technology (SDSMT), Rapid City, South Dakota 57701, USA
³⁵University of Southern Maine, Portland, Maine 04104, USA
³⁶Syracuse University, Syracuse, New York 13244, USA
³⁷Tel Aviv University, Tel Aviv, Israel, 69978
³⁸University of Texas, Arlington, Texas 76019, USA
³⁹Tufts University, Medford, Massachusetts 02155, USA
⁴⁰Center for Neutrino Physics, Virginia Tech, Blacksburg, Virginia 24061, USA
⁴¹University of Warwick, Coventry CV4 7AL, United Kingdom



(Received 6 March 2025; revised 13 November 2025; accepted 19 November 2025; published 19 December 2025)

The MicroBooNE experiment is an 85 tonne active mass liquid argon time projection chamber neutrino detector exposed to the on-axis Booster Neutrino Beam at Fermilab. One of MicroBooNE's physics goals is the precise measurement of neutrino interactions on argon in the 1 GeV energy regime. Building on the capabilities of the MicroBooNE detector, this analysis identifies K^+ mesons, a key signature for the study of strange particle production in neutrino interactions. This measurement is furthermore valuable for background estimation for future nucleon decay searches and for improved reconstruction and particle identification capabilities in experiments such as the Deep Underground Neutrino Experiment. In this Letter, we present the first-ever measurement of a flux-integrated cross section for charged-current muon neutrino induced K^+ production on argon nuclei, determined to be $7.93 \pm 3.22(\text{stat}) \pm 2.83(\text{syst}) \times 10^{-42} \text{ cm}^2/\text{nucleon}$ based on an analysis of 6.88×10^{20} protons on target. This result was found to be consistent with model predictions from different neutrino event generators within the reported uncertainties.

DOI: [10.1103/96vx-gq2g](https://doi.org/10.1103/96vx-gq2g)

Within the standard model, the stability of protons is a consequence of the global symmetry that conserves baryon number. However, various grand unified theories predicting baryon number violation through nucleon decay suggest different decay channels, with a preference for $p \rightarrow e^+\pi^0$, with lifetimes ranging from 10^{34} to 10^{39} yr [1]. Similarly, the Supersymmetric Grand Unified Theory (SUSY GUT) favors protons decaying via $p \rightarrow \nu K^+$, with lifetimes of 10^{32} to 10^{35} yr [2].

Upcoming large long-baseline neutrino oscillation experiments, such as the Deep Underground Neutrino Experiment (DUNE) [3] and Hyper-Kamiokande [4], have the sensitivity to search for proton decay in the range of 10^{32} to 10^{39} yr. Cherenkov detector experiments like Super-Kamiokande [5–7] and Hyper-Kamiokande are

particularly suited for $p \rightarrow e^+\pi^0$ decay searches. However, for water Cherenkov detectors, the $p \rightarrow \nu K^+$ decay channel needs to be indirectly detected since the momentum of the produced K^+ (339 MeV/c) is below the Cherenkov detection threshold in water (560 MeV/c). Furthermore, backgrounds from atmospheric neutrino interactions, such as the $\nu_\mu n \rightarrow \mu^- K^+ n$ and $\nu p \rightarrow \nu K^+ \Lambda$ channels, are challenging to distinguish in water Cherenkov detectors.

In contrast, liquid argon time projection chambers (LArTPCs) can directly measure kaons that range out and decay by analyzing their unique energy loss (dE/dx) profile. These profiles, along with topological signatures such as distinctive kinks from kaon decay and the high ionization Bragg peak, allow an enhanced kaon identification. All of these topological and calorimetric features are easily measured using the millimeter-precision imaging capabilities of an LArTPC.

The first measurements of charged-current (CC) and neutral-current (NC) kaon production, reported with the Argonne National Laboratory [8] and GARGAMELLE bubble chambers [9] in 1974 and 1975, observed $\mathcal{O}(10)$ events. Later, in the 1980's, the Brookhaven National Laboratory bubble chamber reported $\mathcal{O}(10)$ CC and NC

*Contact author: microboone_info@fnal.gov

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

events with strange particle production in neutrino-hydrogen and neutrino-deuterium interactions [10,11], and the Fermilab bubble chamber reported $\mathcal{O}(10)$ events and a measured cross section for CC neutrino interactions $\nu_\mu + n \rightarrow \mu^- + \Lambda + K^+$ on a deuterium target for energy ranges between 10 and 250 GeV [12]. Most recently, the MINERvA experiment measured K^+ production in NC and CC modes on a hydrocarbon target, reporting $\mathcal{O}(100)$ and $\mathcal{O}(1000)$ events respectively [13–15]. In addition, MINERvA reported the first evidence of coherent kaon production [15,16].

This Letter presents the first measurement of the cross section for K^+ production in neutrino interactions on argon nuclei. The signature of the signal events is a CC ν_μ interaction in the presence of a K^+ in the final state with momentum $230 \text{ MeV}/c < p_{K^+} < 2900 \text{ MeV}/c$. The K^+ should decay either to $\mu^+\nu_\mu$ or $\pi^+\pi^0$, which have branching ratios of 63.56% and 20.67%, respectively [17]. This measurement uses MicroBooNE datasets collected between 2016 and 2018 from the on-axis Booster Neutrino Beam (BNB), equivalent to 6.88×10^{20} protons on target (POT).

As the first quantitative measurement of K^+ production on argon nuclei, this analysis will impact nuclear, neutrino, and proton decay research on two fronts. First, it provides background constraints in future $p \rightarrow \nu K^+$ nucleon decay experiments. Second, this measurement provides input to improve neutrino generator models. In the case of bound proton decay, the K^+ is generated within the nucleus and may interact with other nucleons as it escapes. These interactions can decrease the detected momentum of the K^+ in the detector and may also eject nucleons from the nucleus.

Charged-current neutrino-induced K^+ production occurs through three interaction modes: (1) *single kaon production*, where a CC interaction produces a single kaon with a production energy threshold of $E_{\nu_\mu} \geq 0.79 \text{ GeV}$, and strangeness is not conserved ($|\Delta S| = 1$); (2) *associated kaon production*, where both a kaon and a hyperon are produced, with a threshold of $E_{\nu_\mu} \geq 1.2 \text{ GeV}$, and strangeness is conserved; and (3) *coherent production*, where only one kaon is produced in the final state, and the target nucleus remains intact after the neutrino interaction. Figure 1 shows an event candidate of CC ν_μ kaon production recorded in MicroBooNE data. The kaon decays at rest into a 388 MeV energy muon, which subsequently decays into a Michel electron.

The MicroBooNE detector [18,19] consists of an 85 tonne LArTPC and a photon detection system composed of 32 photomultiplier tubes. The detector identifies neutrino interactions by detecting ionization and scintillation light produced by charged particles moving through the detector. Ionization charge is recorded across three wire planes (two induction, one collection) oriented at different angles (with induction planes at $\pm 60^\circ$ relative to the vertical




FIG. 1. A $\nu_\mu + \text{Ar} \rightarrow \mu^- + K^+$ interaction candidate observed in MicroBooNE data recorded by the collection plane. The color scale represents the intensity of the ionization charge collected on the time projection chamber (TPC) wires (green, low intensity; red, high intensity). The x and y axes represent the wire position in the beam direction and the distance in the drift direction, respectively. The gap shown on each track is due to a region in the detector with no active wires.

collection plane), and 3 mm wire spacing. This configuration allows the generation of mm-scale resolution, three-dimensional images of neutrino interactions. Scintillation light collected by the photomultiplier tubes provides ns-scale timing information, essential for identifying neutrino interactions that are synchronized with the BNB, and for rejecting cosmic-ray backgrounds.

In MicroBooNE, the LArSoft framework [20] provides tools for simulating the neutrino flux, neutrino interactions, particle propagation, and detector response, including ionization and scintillation processes as well as reconstruction of neutrino events. The BNB neutrino flux through the MicroBooNE detector is simulated using the MiniBooNE flux model [21] adapted for MicroBooNE's location along the beamline. GENIE v3.0.6 (G18_10a_02_11a) [22,23] is the event generator used to simulate neutrino-argon interactions, including both single and associated kaon production. In GENIE, associated kaons are produced via individual resonances simulated by the Rein-Sehgal model [24]. Kaons produced via deep inelastic scattering are simulated by the Koba-Nielsen-Olesen scaling parametrization [25]. For higher energies (hadronic invariant mass $W > 2.3 \text{ GeV}$), the AGKY model [26] and Pythia6 [27] are used. The hadronization models are tuned to match strange particle production (Λ and K_S^0) observed in experiments such as the Big European Bubble Chamber [28–31] and the Fermilab bubble chamber [32,33]. The single kaon production model [34] is implemented in GENIE, generating events in the channels $\nu_l + n \rightarrow l^- + n + K^+$, $\nu_l + p \rightarrow l^- + p + K^+$, and $\nu_l + n \rightarrow l^- + p + K^0$. The region of validity assumed for the model is for $E_\nu \leq 2 \text{ GeV}$ [34,35], while the GENIE authors note that the model can be applied to higher energies as well [36].




FIG. 2. BDT classifier score distribution. The black dots represent the MicroBooNE data. The stacked histogram represents the expected BDT distribution from the simulation with its statistical uncertainty (gray band). The orange arrow represents the BDT event selection cut. The red triangles are the K^+ candidates selected in MicroBooNE data by the BDT.

Particle trajectories and interactions within the detector are simulated using GEANT4 v4_10_3_p03c using the QGSP_BERT hadron physics list [37–39], which models particle propagation in the detector. Simulated neutrino interactions are overlayed with data events collected using an unbiased trigger that operates in anticoincidence with the beam, enabling data-driven modeling of cosmic-ray and detector noise. Both simulation and data reconstruction use the Pandora framework [40], which uses a multialgorithm approach to pattern recognition and event reconstruction in LArTPCs. Pandora creates clusters from wire hits, identifies the neutrino vertex, matches clusters across wire planes to form 3D objects, classifies reconstructed particles as tracks or showers, and establishes particle hierarchies by identifying parent-daughter relationships among particles.

Before performing a dedicated selection for K^+ production signal events, a preselection is applied as follows: (1) events must pass a CC inclusive filter that isolates CC neutrino interactions by identifying the presence of an outgoing muon from the vertex of a neutrino interaction [41], (2) all reconstructed tracks in an event must originate within a $2.4 \times 2.1 \times 9.8$ m fiducial volume centered inside the MicroBooNE TPC, and (3) the daughter track (defined as a track connected to the endpoint of one primary signal track candidate) should stop at least 5 cm away from any of the physical edges of the TPC to ensure containment.

For the K^+ event selection, we implemented a boosted decision tree (BDT) classifier [42]. The BDT classifier

inputs include reconstructed variables such as χ^2 values, which compare the expected dE/dx for kaon (χ_k^2), proton (χ_p^2), pion (χ_π^2), and muon (χ_μ^2) hypotheses with the observed data for each wire plane, the combined three-wire-plane χ^2 , the daughter track length, and the log-likelihood ratio particle identification [43]. We trained the BDT classifier to identify true K^+ that decay to $\mu^+\nu_\mu$ or $\pi^+\pi^0$, using Monte Carlo (MC) simulation samples of reconstructed neutrino interactions that meet the preselection criteria described above. To train the BDT classifier, a background sample corresponding to 3.67×10^{20} POT (equivalent to 10% of the MicroBooNE BNB simulation) was used. For the simulation of the signal, a dedicated high-statistics sample of single and associated K^+ production was used. We optimized the BDT classifier score selection criteria by maximizing the product of efficiency and purity. Figure 2 shows the BDT classifier score distribution, with the orange arrow at 0.41 representing the selected BDT score cut. Events with a score higher than the BDT selection are kept as K^+ candidates, and 10 such K^+ candidates are identified in the data. The total number of background events predicted from neutrino-induced interactions, cosmic rays recorded during neutrino-triggered beam spills, and off-beam data is 2.21 events. The simulation predicts a selection efficiency of 3.95%, and an associated selection purity of 79%. The primary background category for K^+ candidates consists of protons generated at the neutrino vertex that interact with an argon nucleus, producing secondary particles at the proton endpoint that could mimic a K^+ decay product. This background category comprises 2.08 out of 2.21 predicted background events.

Figure 3 shows the reconstructed daughter track length distribution of the K^+ candidates selected in MicroBooNE data, including the event shown in Fig. 1. The stacked histogram shows the expected daughter track length distribution from the MicroBooNE BNB simulation. Because of the two-body decay nature of the K^+ branching ratios analyzed in this Letter, the resulting monoenergetic μ^+ (π^+) provides a powerful discriminant, as it leads to a fixed track length that can be measured with high accuracy and is not present in background events. The expected track length for μ^+ (π^+) accumulates around 53 cm (30 cm). The K^+ candidates selected by the BDT selection in Fig. 3 (red triangles) exhibit a pileup in the region compatible with K^+ decay.

To evaluate the accuracy of the background modeling, two sideband regions were defined using a two-dimensional distribution of the BDT classifier score versus daughter track length. A far sideband region is defined to include tracks with low BDT classifier scores ($\text{BDT} < 0$) and track lengths less than 40 cm. A near sideband region selects tracks with BDT classifier scores close to the signal region ($\text{BDT} < 0.41$) and across all track length ranges, but excludes tracks in the far sideband region. The data and MC




FIG. 3. Reconstructed daughter track length distribution. The stacked histogram shows the MC simulation with its statistical uncertainty (gray band). The black dots represent the MicroBooNE data. The red triangles are the K^+ candidates selected in MicroBooNE data by the BDT, where three data points lie close to 60 cm, nearly overlapping.

comparisons of the near and far sidebands show agreement for reconstructed kinematic variables such as K^+ candidate track length, ϕ (azimuthal angle around the beam direction), and θ (angle of the track with respect to the beam direction). For the K^+ candidate track length data and MC comparison, χ^2/ndf (number of degrees of freedom) values (including statistical uncertainties) of 16.87/24 and 17.93/24 were obtained for the near and far sidebands. The agreement with the data suggests that the simulation is able to adequately model the background in the cross-section extraction. More information about the sideband comparisons is provided in Supplemental Material [44].

This analysis presents the measurement of the single-bin flux-integrated cross section for $\nu_\mu + \text{Ar} \rightarrow \mu^- + K^+ + X$ interactions, where muon neutrinos interact with Ar and produce one μ^- , one K^+ , and X (any number of other hadrons in the final state) on argon nuclei. The cross section (σ) was found to be $7.93 \pm 3.22(\text{stat}) \pm 2.83(\text{syst}) \times 10^{-42} \text{ cm}^2/\text{nucleon}$, and it was calculated using the following expression:

$$\sigma = \frac{N - B}{\epsilon \times N_{\text{target}} \times \Phi_{\nu_\mu}}, \quad (1)$$

where N represents the number of selected signal events (10), B is the number of expected background events (2.21), ϵ is the selection efficiency (3.95%), N_{target} is the number of target nucleons (4.13×10^{31}), and Φ_{ν_μ} stands for the integrated BNB ν_μ flux ($6.02 \times 10^{11} \text{ } \nu/\text{cm}^2$). The

selection efficiency accounts for all K^+ decays within the fiducial volume.

The systematic uncertainties in the measured cross section are evaluated by reweighting and generating simulated events to account for each source of uncertainty [46]. For the detector-related uncertainties, a dedicated simulation is used to model the variation in detector responses [47,48]. In contrast, uncertainties from neutrino interactions and BNB flux modeling are treated with a reweighting technique. To evaluate the uncertainty associated with kaon reinteractions within the detector, we implement a methodology similar to that presented by MINERvA [13]. For each selected K^+ candidate track, a weight W_{inel} is calculated using the following expression to estimate the kaon inelastic reinteraction uncertainty:

$$W_{\text{inel}} = \frac{1 - e^{-\rho x \sigma_{\text{var}}^{\text{tot}}}}{1 - e^{-\rho x \sigma_{\text{geant}}^{\text{tot}}}} \times \frac{\sigma_{\text{var}}^{\text{inel}}}{\sigma_{\text{var}}^{\text{tot}}} \times \frac{\sigma_{\text{geant}}^{\text{tot}}}{\sigma_{\text{geant}}^{\text{inel}}}, \quad (2)$$

where ρ is the density of the material, x is the track length, $\sigma_{\text{geant}}^{\text{tot}}$ and $\sigma_{\text{geant}}^{\text{inel}}$ are the K^+ total and inelastic GEANT4 cross sections, and $\sigma_{\text{var}}^{\text{tot}}$ and $\sigma_{\text{var}}^{\text{inel}}$ are the K^+ total and inelastic cross sections estimated by applying a variation of $\pm 40\%$ to the $\sigma_{\text{geant}}^{\text{inel}}$. The 40% variation is motivated by observed disagreement between experimental measurements of $K^+ - \text{Ca}$ interactions [45] and the predictions from the GEANT4 interaction model shown in Supplemental Material [44].

The leading source of systematic uncertainty comes from the detector response (30.3%), with a major contribution from the modeling of ion recombination of electrons with argon. This analysis relies on identifying stopping K^+ and μ^+ (π^+) through their Bragg peaks, which are characterized by a sharp rise in energy deposition near the end of the particle track. These effects occur in a highly ionizing regime, where ion recombination effects are most pronounced. As the charge profile is impacted by recombination, any mismodeling can distort the Bragg peak shape and affect particle identification. The modeling of neutrino interactions, the BNB neutrino flux, and particle reinteractions in the detector, including kaon inelastic interactions, lead to systematic uncertainties of 3.7%, 11.7%, and 10.5%, respectively. The neutrino interaction modeling uncertainty on the cross-section measurement enters through the modeling of the subtracted background and the signal efficiency prediction. The neutrino interaction modeling uncertainty was evaluated by varying the GENIE parameters within their nominal 1σ ranges [46]. These variations were propagated through the full simulation to assess their impact on the measured kaon production cross-section uncertainty budget. The main contributions to the neutrino interaction modeling arise from CC resonance production (axial and vector masses). The resulting uncertainty shows limited impact on the selection efficiency, as the selection depends mainly on topological and

TABLE I. The $\nu_\mu + \text{Ar} \rightarrow \mu^- + K^+ + X$ cross section extracted from different neutrino event generators compared with the cross section extracted from data.

Generator	Cross section (10^{-42} cm ² /nucleon)
GENIE v2.12.10	8.67
GENIE v3.00.06	8.42
GENIE v3.4.0 (AR23)	9.85
GiBUU 2025, patch 1	6.52
NEUT 5.4.0.1 [51]	9.71
NuWro 19.02.1	10.87
MicroBooNE data	$7.93 \pm 3.22(\text{stat}) \pm 2.83(\text{syst})$

calorimetric features rather than interaction kinematics that are more tied to detailed cross-section modeling. The effect of statistics in the estimation of the selection efficiency (6.1%), the number of target nuclei N_{target} (1%), MC statistical uncertainties on the background estimation (7.4%), and POT exposure (2%) are the remaining sources of uncertainty. The overall magnitude of the uncertainty (statistical and systematic) on our measurement is 54.1%, where the major contribution is due to the statistical uncertainty, with a reported value of 40.6%. Table I presents the measured cross section of the $\nu_\mu + \text{Ar} \rightarrow \mu^- + K^+ + X$ interactions and the predictions obtained from various neutrino event generators with the NUISANCE framework [49].

The flux-integrated cross-section measurement reported in this Letter indicates consistency with the predictions from different neutrino event generators, although the large uncertainties prevent a stringent test of these models. The cross sections predicted by NuWro [50] and GiBUU 2025, patch 1 [52] show larger deviations from the data, but remain within the 1σ uncertainty of the measurement.

In conclusion, this Letter presents the first measurement of the $\nu_\mu + \text{Ar} \rightarrow \mu^- + K^+ + X$ cross section on argon, using the MicroBooNE dataset collected from the BNB flux. The results reported here represent a significant step forward for future studies aiming to refine background estimates for proton decay searches, particularly within the framework of grand unified theories and supersymmetry. To advance this analysis in future LArTPC measurements, it is necessary to increase statistics, reduce large detector systematics (especially those related to the recombination of electrons with argon nuclei), refine reconstruction algorithms for short tracks, and improve the identification of kaon tracks that undergo interactions. Experiments such as SBND [53], ICARUS [54], and the DUNE near detector will allow precision measurements of neutrino-induced K^+ production on argon, including differential cross-section measurements for the first time ever. Furthermore, the uncertainty related to kaon reinteractions on argon nuclei could be refined by using data from LArIAT [55], and the DUNE prototypes at CERN [56]. More statistics will come from analyzing MicroBooNE's full dataset with a total of

1.2×10^{21} POT. The future DUNE near detector data could also provide more accurate measurements of CC neutrino-induced K^+ production on argon nuclei, enabling improved comparisons among the neutrino generator models. These high-precision measurements will enhance our understanding of rare neutrino interactions and final state interaction modeling, and will improve the background predictions for future nucleon decay searches at DUNE, which currently rely on model predictions.

Acknowledgments—This document was prepared by the MicroBooNE collaboration using the resources of the Fermi National Accelerator Laboratory (Fermilab), a U.S. Department of Energy, Office of Science, Office of High Energy Physics HEP User Facility. Fermilab is managed by Fermi Forward Discovery Group, LLC, acting under Contract No. 89243024CSC000002. MicroBooNE is supported by the following: the U.S. Department of Energy, Office of Science, Offices of High Energy Physics and Nuclear Physics; the U.S. National Science Foundation; the Swiss National Science Foundation; the Science and Technology Facilities Council (STFC), part of United Kingdom Research and Innovation (UKRI); the Royal Society (United Kingdom); the UKRI Future Leaders Fellowship; the NSF AI Institute for Artificial Intelligence and Fundamental Interactions; and the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement No. 101003460 (PROBES). Additional support for the laser calibration system and cosmic-ray tagger was provided by the Albert Einstein Center for Fundamental Physics, Bern, Switzerland. We also acknowledge the contributions of technical and scientific staff to the design, construction, and operation of the MicroBooNE detector as well as the contributions of past collaborators to the development of MicroBooNE analyses, without whom this work would not have been possible.

Data availability—The data that support the findings of this article are not publicly available. The data are available from the authors upon reasonable request.

-
- [1] H. Georgi and S.L. Glashow, Unity of all elementary-particle forces, *Phys. Rev. Lett.* **32**, 438 (1974).
 - [2] J. Hisano, H. Murayama, and T. Yanagida, Nucleon decay in the minimal supersymmetric SU(5) grand unification, *Nucl. Phys. B* **402**, 46 (1993).
 - [3] B. Abi *et al.*, Deep underground neutrino experiment (DUNE), far detector technical design report, volume I. Introduction to DUNE, *J. Instrum.* **15**, T08008 (2020).
 - [4] K. Abe *et al.* (Super-Kamiokande Collaboration), Hyper-Kamiokande design report, [arXiv:1805.04163](https://arxiv.org/abs/1805.04163).
 - [5] K. Abe *et al.* (Super-Kamiokande Collaboration), Search for proton decay via $p \rightarrow \nu K^+$ using 260 kiloton Δ year data of Super-Kamiokande, *Phys. Rev. D* **90**, 072005 (2014).

- [6] K. Abe *et al.* (Super-Kamiokande Collaboration), Search for proton decay via $p \rightarrow e^+\pi^0$ and $p \rightarrow \mu^+\pi^0$ in 0.31 megaton $\cdot$ years exposure of the Super-Kamiokande water Cherenkov detector, *Phys. Rev. D* **95**, 012004 (2017).
- [7] K. Abe *et al.* (Super-Kamiokande Collaboration), Search for proton decay via $p \rightarrow e^+\pi^0$ and $p \rightarrow \mu^+\pi^0$ with an enlarged fiducial volume in Super-Kamiokande I-IV, *Phys. Rev. D* **102**, 112011 (2020).
- [8] S. J. Barish *et al.*, Strange-particle production in neutrino interactions, *Phys. Rev. Lett.* **33**, 1446 (1974).
- [9] H. Deden *et al.*, Strange particle production and charmed particle search in the Gargamelle neutrino experiment, *Phys. Lett. B* **58**, 361 (1975).
- [10] J. E. Jensen, The Brookhaven National Laboratory 80-inch liquid hydrogen bubble chamber, in *Proceedings of the 1964 Cryogenic Engineering Conference* (Brookhaven National Laboratory, Upton, New York, 1964).
- [11] N. J. Baker, P. L. Connolly, S. A. Kahn, H. G. Kirk, M. J. Murtagh, R. B. Palmer, N. P. Samios, and M. Tanaka, Strange-particle production from neutrino interactions in the BNL 7-foot bubble chamber, *Phys. Rev. D* **24**, 2779 (1981).
- [12] D. Son *et al.*, Quasielastic charmed-baryon production and exclusive strange-particle production by high-energy neutrinos, *Phys. Rev. D* **28**, 1219 (1983).
- [13] C. M. Marshall *et al.* (MINERvA Collaboration), Measurement of K^+ production in charged-current ν_μ interactions, *Phys. Rev. D* **94**, 012002 (2016).
- [14] C. M. Marshall *et al.* (MINERvA Collaboration), Measurement of neutral-current K^+ production by neutrinos using MINERvA, *Phys. Rev. Lett.* **119**, 011802 (2017).
- [15] C. M. Marshall, Measurement of charged kaon production by neutrinos at MINERvA, Ph.D. thesis, University of Rochester, 2016.
- [16] Z. Wang, C. M. Marshall *et al.* (MINERvA Collaboration), Evidence of coherent K^+ meson production in neutrino-nucleus scattering, *Phys. Rev. Lett.* **117**, 061802 (2016).
- [17] S. Navas *et al.* (Particle Data Group), Review of particle physics, *Phys. Rev. D* **110**, 030001 (2024).
- [18] B. Fleming (MicroBooNE Collaboration), The MicroBooNE technical design report, 10.2172/1333130 (2012).
- [19] R. Acciarri *et al.* (MicroBooNE Collaboration), Design and construction of the MicroBooNE detector, *J. Instrum.* **12**, P02017 (2017).
- [20] E. L. Snider and G. Petrillo, LArSoft: Toolkit for simulation, reconstruction and analysis of liquid argon TPC neutrino detectors, *J. Phys. Conf. Ser.* **898**, 042057 (2017).
- [21] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), Neutrino flux prediction at MiniBooNE, *Phys. Rev. D* **79**, 072002 (2009).
- [22] C. Andreopoulos *et al.*, The GENIE neutrino Monte Carlo generator, *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 87 (2010).
- [23] C. Andreopoulos *et al.* (GENIE Collaboration), Neutrino-nucleon cross-section model tuning in GENIE v3, *Phys. Rev. D* **104**, 072009 (2021).
- [24] D. Rein and L. M. Sehgal, Neutrino-excitation of baryon resonances and single pion production, *Ann. Phys. (N.Y.)* **133**, 79 (1981).
- [25] Z. Koba, H. B. Nielsen, and P. Olesen, Scaling of multiplicity distributions in high energy hadron collisions, *Nucl. Phys. B* **40**, 317 (1972).
- [26] T. Yang, C. Andreopoulos, H. Gallagher, K. Hofmann, and P. Kehayias, A hadronization model for few-GeV neutrino interactions, *Eur. Phys. J. C* **63**, 1 (2009).
- [27] T. Sjöstrand, S. Mrenna, and P. Skands, Pythia 6.4 physics and manual, *J. High Energy Phys.* **05** (2006) 026.
- [28] G. T. Jones *et al.*, Neutral strange particle production in neutrino and anti-neutrino charged current interactions on protons, *Z. Phys. C* **57**, 197 (1993).
- [29] D. Allasia *et al.*, Production of neutral strange particles in $\bar{\nu}_\mu d_2$ and $\nu_\mu d_2$ charged current interactions, *Nucl. Phys. B* **224**, 1 (1983).
- [30] S. Willocq *et al.* (WA59 Collaboration), Neutral strange particle production in anti-neutrino—neon charged current interactions, *Z. Phys. C* **53**, 207 (1992).
- [31] P. Bosetti *et al.*, Strange particle production in ν and $\bar{\nu}$ neon interactions, *Nucl. Phys. B* **209**, 29 (1982).
- [32] N. J. Baker *et al.*, Strange-particle production in neutrino-neon charged-current interactions, *Phys. Rev. D* **34**, 1251 (1986).
- [33] D. DeProspo *et al.*, Neutral strange particle production in neutrino and antineutrino charged-current interactions on neon, *Phys. Rev. D* **50**, 6691 (1994).
- [34] M. Rafi Alam, I. Ruiz Simo, M. Sajjad Athar, and M. J. Vicente Vacas, Weak kaon production off the nucleon, *Phys. Rev. D* **82**, 033001 (2010).
- [35] L. Alvarez-Ruso, J. Nieves, I. Ruiz Simo, M. Valverde, and M. J. Vicente Vacas, Charged kaon production by coherent scattering of neutrinos and antineutrinos on nuclei, *Phys. Rev. C* **87**, 015503 (2013).
- [36] M. Alam, C. Andreopoulos *et al.*, GENIE production release 2.10.0, [arXiv:1512.06882](https://arxiv.org/abs/1512.06882).
- [37] S. Agostinelli *et al.*, GEANT4—a simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [38] J. Allison *et al.*, GEANT4 developments and applications, *IEEE Trans. Nucl. Sci.* **53**, 270 (2006).
- [39] J. Allison *et al.*, Recent developments in GEANT4, *Nucl. Instrum. Methods Phys. Res., Sect. A* **835**, 186 (2016).
- [40] R. Acciarri *et al.*, The Pandora multi-algorithm approach to automated pattern recognition of cosmic-ray muon and neutrino events in the MicroBooNE detector, *Eur. Phys. J. C* **78**, 82 (2018).
- [41] P. Abratenko *et al.* (MicroBooNE Collaboration), First measurement of inclusive muon neutrino charged current differential cross sections on argon at $E_\nu \sim 0.8$ GeV with the MicroBooNE detector, *Phys. Rev. Lett.* **123**, 131801 (2019).
- [42] A. Hoecker *et al.*, TMVA—toolkit for multivariate data analysis, [arXiv:physics/0703039](https://arxiv.org/abs/physics/0703039).
- [43] P. Abratenko *et al.* (MicroBooNE Collaboration), Calorimetric classification of track-like signatures in liquid argon TPCs using MicroBooNE data, *J. High Energy Phys.* **12** (2021) 153.
- [44] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/96vx-gq2g> for further details on side-band studies, event selection displays, and K^+ -Ca data-GEANT4 cross-section comparison, which includes Ref. [45].
- [45] E. Friedman *et al.*, K^+ nucleus reaction and total cross sections: New analysis of transmission experiments, *Phys. Rev. C* **55**, 1304 (1997).

- [46] P. Abratenko *et al.* (MicroBooNE Collaboration), New $CC0\pi$ GENIE model tune for MicroBooNE, *Phys. Rev. D* **105**, 072001 (2022).
- [47] P. Abratenko *et al.* (MicroBooNE Collaboration), Novel approach for evaluating detector-related uncertainties in a LArTPC using MicroBooNE data, *Eur. Phys. J. C* **82**, 454 (2022).
- [48] J. Calcutt, C. Thorpe, K. Mahn, and L. Fields, GEANT4Re-weight: A framework for evaluating and propagating hadronic interaction uncertainties in GEANT4, *J. Instrum.* **16**, P08042 (2021).
- [49] P. Stowell *et al.*, NUISANCE: A neutrino cross-section generator tuning and comparison framework, *J. Instrum.* **12**, P01016 (2017).
- [50] T. Golan, J. T. Sobczyk, and J. Żmuda, NuWro: The Wrocław Monte Carlo generator of neutrino interactions, *Nucl. Phys. B, Proc. Suppl.* **229–232**, 499 (2012).
- [51] Y. Hayato and L. Pickering, The NEUT neutrino interaction simulation program library, *Eur. Phys. J.: Spec. Top.* **230**, 24 (2021).
- [52] GiBUU Collaboration, The Giessen Boltzmann-Uehling-Uhlenbeck (GiBUU) project, <https://gibuu.hepforge.org> (2025).
- [53] R. Acciarri *et al.* (SBND Collaboration), The short-baseline near detector at Fermilab, [arXiv:2504.00245](https://arxiv.org/abs/2504.00245).
- [54] S. Amerio *et al.*, Design, construction and tests of the ICARUS T600 detector, *Nucl. Instrum. Methods Phys. Res., Sect. A* **527**, 329 (2004).
- [55] E. Gramellini, Measurement of the negative pion and positive kaon total hadronic cross sections on argon at the LArIAT experiment, Ph.D. thesis, Yale University, 2018.
- [56] A. Abed Abud *et al.* (DUNE Collaboration), First measurement of the total inelastic cross section of positively charged kaons on argon at energies between 5.0 and 7.5 GeV, *Phys. Rev. D* **110**, 092011 (2024).