

Observation of the Radiative Decay $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$

M. Abumusabh¹, I. Adachi², L. Aggarwal³, H. Ahmed⁴, Y. Ahn⁵, H. Aihara⁶, N. Akopov⁷, S. Alghamdi⁸, M. Alhakami⁹, A. Aloisio¹⁰, N. Althubiti¹¹, K. Amos¹², N. Anh Ky¹³, C. Antonioli¹⁴, D. M. Asner¹⁵, H. Atmacan¹⁶, T. Aushev¹⁷, R. Ayad¹⁸, V. Babu¹⁹, N. K. Baghel²⁰, S. Bahinipati²¹, P. Bambade²², Sw. Banerjee²³, M. Barrett²⁴, M. Bartl²⁵, J. Baudot²⁶, A. Beaubien²⁷, J. Becker²⁸, J. V. Bennett²⁹, F. U. Bernlochner³⁰, V. Bertacchi³¹, M. Bertemes³², E. Bertholet³³, M. Bessner³⁴, S. Bettarini³⁵, F. Bianchi³⁶, T. Bilka³⁷, D. Biswas³⁸, A. Bobrov³⁹, D. Bodrov⁴⁰, A. Bondar⁴¹, G. Bonvicini⁴², J. Borah⁴³, A. Boschetti⁴⁴, A. Bozek⁴⁵, M. Bračko⁴⁶, P. Branchini⁴⁷, R. A. Briere⁴⁸, T. E. Browder⁴⁹, A. Budano⁵⁰, S. Bussino⁵¹, Q. Campagna⁵², M. Campajola⁵³, G. Casarosa⁵⁴, C. Cecchi⁵⁵, P. Chang⁵⁶, P. Cheema⁵⁷, L. Chen⁵⁸, B. G. Cheon⁵⁹, C. Cheshta⁶⁰, H. Chetri⁶¹, K. Chilikin⁶², K. Chirapatpimol⁶³, H.-E. Cho⁶⁴, K. Cho⁶⁵, S.-J. Cho⁶⁶, S.-K. Choi⁶⁷, S. Choudhury⁶⁸, S. Chutia⁶⁹, J. A. Colorado-Cacedo⁷⁰, I. Consigny⁷¹, L. Corona⁷², J. X. Cui⁷³, E. De La Cruz-Burelo⁷⁴, S. A. De La Motte⁷⁵, G. De Nardo⁷⁶, G. De Pietro⁷⁷, R. de Sangro⁷⁸, M. Destefanis⁷⁹, S. Dey⁸⁰, A. Di Canto⁸¹, Z. Doležal⁸², I. Domínguez Jiménez⁸³, T. V. Dong⁸⁴, X. Dong⁸⁵, M. Dorigo⁸⁶, G. Dujany⁸⁷, P. Ecker⁸⁸, J. Eppelt⁸⁹, R. Farkas⁹⁰, P. Feichtinger⁹¹, T. Ferber⁹², T. Fillinger⁹³, C. Finck⁹⁴, G. Finocchiaro⁹⁵, F. Forti⁹⁶, B. G. Fulsom⁹⁷, A. Gabrielli⁹⁸, E. Ganiev⁹⁹, M. Garcia-Hernandez¹⁰⁰, R. Garg¹⁰¹, G. Gaudino¹⁰², V. Gaur¹⁰³, V. Gautam¹⁰⁴, A. Gaz¹⁰⁵, A. Gellrich¹⁰⁶, G. Ghevondyan¹⁰⁷, D. Ghosh¹⁰⁸, H. Ghumaryan¹⁰⁹, R. Giordano¹¹⁰, A. Giri¹¹¹, P. Gironella Gironell¹¹², A. Glazov¹¹³, B. Gobbo¹¹⁴, R. Godang¹¹⁵, O. Gogota¹¹⁶, P. Goldenzweig¹¹⁷, W. Gradl¹¹⁸, E. Graziani¹¹⁹, D. Greenwald¹²⁰, K. Gudkova¹²¹, I. Haide¹²², Y. Han¹²³, H. Hayashii¹²⁴, S. Hazra¹²⁵, C. Hearty¹²⁶, M. T. Hedges¹²⁷, G. Heine¹²⁸, I. Heredia de la Cruz¹²⁹, T. Higuchi¹³⁰, M. Hoek¹³¹, M. Hohmann¹³², R. Hoppe¹³³, P. Horak¹³⁴, X. T. Hou¹³⁵, C.-L. Hsu¹³⁶, A. Huang¹³⁷, T. Humair¹³⁸, T. Iijima¹³⁹, N. Ipsita¹⁴⁰, A. Ishikawa¹⁴¹, R. Itoh¹⁴², M. Iwasaki¹⁴³, D. Jacobi¹⁴⁴, W. W. Jacobs¹⁴⁵, E.-J. Jang¹⁴⁶, Q. P. Ji¹⁴⁷, S. Jia¹⁴⁸, Y. Jin¹⁴⁹, A. Johnson¹⁵⁰, J. Kandra¹⁵¹, K. H. Kang¹⁵², S. Kang¹⁵³, G. Karyan¹⁵⁴, F. Keil¹⁵⁵, C. Kiesling¹⁵⁶, D. Y. Kim¹⁵⁷, J.-Y. Kim¹⁵⁸, K.-H. Kim¹⁵⁹, H. Kindo¹⁶⁰, K. Kinoshita¹⁶¹, P. Kodyš¹⁶², T. Koga¹⁶³, S. Kohani¹⁶⁴, A. Korobov¹⁶⁵, S. Korpar¹⁶⁶, E. Kovalenko¹⁶⁷, R. Kowalewski¹⁶⁸, P. Križan¹⁶⁹, P. Krokovny¹⁷⁰, T. Kuhr¹⁷¹, D. Kumar¹⁷², K. Kumara¹⁷³, T. Kunigo¹⁷⁴, A. Kuzmin¹⁷⁵, Y.-J. Kwon¹⁷⁶, S. Lacaprara¹⁷⁷, T. Lam¹⁷⁸, J. S. Lange¹⁷⁹, T. S. Lau¹⁸⁰, M. Laurenza¹⁸¹, R. Leboucher¹⁸², F. R. Le Diberder¹⁸³, H. Lee¹⁸⁴, M. J. Lee¹⁸⁵, C. Lemettais¹⁸⁶, P. Leo¹⁸⁷, P. M. Lewis¹⁸⁸, C. Li¹⁸⁹, H.-J. Li¹⁹⁰, L. K. Li¹⁹¹, Q. M. Li¹⁹², S. X. Li¹⁹³, W. Z. Li¹⁹⁴, Y. Li¹⁹⁵, Y. B. Li¹⁹⁶, Y. P. Liao¹⁹⁷, J. Libby¹⁹⁸, J. Lin¹⁹⁹, V. Lisovskyi²⁰⁰, M. H. Liu²⁰¹, Q. Y. Liu²⁰², Z. Q. Liu²⁰³, D. Liventsev²⁰⁴, S. Longo²⁰⁵, A. Lozar²⁰⁶, T. Lueck²⁰⁷, C. Lyu²⁰⁸, J. L. Ma²⁰⁹, Y. Ma²¹⁰, M. Maggiora²¹¹, S. P. Maharana²¹², R. Maiti²¹³, G. Mancinelli²¹⁴, R. Manfredi²¹⁵, M. Mantovano²¹⁶, D. Marcantonio²¹⁷, M. Marfoli²¹⁸, C. Marinas²¹⁹, C. Martellini²²⁰, A. Martens²²¹, T. Martinov²²², L. Massaccesi²²³, M. Masuda²²⁴, D. Matvienko²²⁵, M. Maushart²²⁶, J. A. McKenna²²⁷, Z. Mediankin Gruberová²²⁸, R. Mehta²²⁹, F. Meier²³⁰, D. Meleshko²³¹, M. Merola²³², C. Miller²³³, M. Mirra²³⁴, H. Miyake²³⁵, R. Mizuk²³⁶, G. B. Mohanty²³⁷, S. Moneta²³⁸, H.-G. Moser²³⁹, I. Nakamura²⁴⁰, M. Nakao²⁴¹, M. Naruki²⁴², Z. Natkaniec²⁴³, A. Natchii²⁴⁴, M. Nayak²⁴⁵, S. Nishida²⁴⁶, R. Nomaru²⁴⁷, S. Ogawa²⁴⁸, H. Ono²⁴⁹, F. Otani²⁵⁰, G. Pakhlova²⁵¹, A. Panta²⁵², S. Pardi²⁵³, K. Parham²⁵⁴, J. Park²⁵⁵, S.-H. Park²⁵⁶, A. Passeri²⁵⁷, S. Patra²⁵⁸, S. Paul²⁵⁹, T. K. Pedlar²⁶⁰, R. Pestotnik²⁶¹, M. Piccolo²⁶², L. E. Piilonen²⁶³, P. L. M. Podesta-Lerma²⁶⁴, T. Podobnik²⁶⁵, C. Praz²⁶⁶, S. Prell²⁶⁷, M. T. Prim²⁶⁸, S. Privalov²⁶⁹, H. Purwar²⁷⁰, P. Rados²⁷¹, S. Raiz²⁷², K. Ravindran²⁷³, J. U. Rehman²⁷⁴, M. Reif²⁷⁵, S. Reiter²⁷⁶, L. Reuter²⁷⁷, D. Ricalde Herrmann²⁷⁸, I. Ripp-Baudot²⁷⁹, G. Rizzo²⁸⁰, S. H. Robertson²⁸¹, J. M. Roney²⁸², A. Rostomyan²⁸³, S. Saha²⁸⁴, L. Salutati²⁸⁵, D. A. Sanders²⁸⁶, L. Santelj²⁸⁷, C. Santos²⁸⁸, V. Savinov²⁸⁹, B. Scavino²⁹⁰, S. Schneider²⁹¹, K. Schoenning²⁹², C. Schwanda²⁹³, Y. Seino²⁹⁴, K. Senyo²⁹⁵, J. Serrano²⁹⁶, C. Sfienti²⁹⁷, W. Shan²⁹⁸, G. Sharma²⁹⁹, C. P. Shen³⁰⁰, X. D. Shi³⁰¹, T. Shillington³⁰², J.-G. Shiu³⁰³, D. Shtol³⁰⁴, B. Shwartz³⁰⁵, A. Sibidanov³⁰⁶, F. Simon³⁰⁷, J. Skorupa³⁰⁸, R. J. Sobie³⁰⁹, M. Sobotzik³¹⁰, A. Soffer³¹¹, A. Sokolov³¹², E. Solovieva³¹³, S. Spataro³¹⁴, K. Špenko³¹⁵, B. Spruck³¹⁶, M. Starič³¹⁷, P. Stavroulakis³¹⁸, R. Stroili³¹⁹, M. Sumihama³²⁰, S. S. Tang³²¹, K. Tanida³²², F. Tenchini³²³, F. Testa³²⁴, A. Thaller³²⁵, T. Tien Manh³²⁶, O. Tittel³²⁷, R. Tiwary³²⁸, E. Torassa³²⁹, K. Trabelsi³³⁰, F. F. Trantou³³¹, I. Ueda³³², K. Unger³³³, Y. Unno³³⁴, K. Uno³³⁵, S. Uno³³⁶, P. Urquijo³³⁷, Y. Ushiroda³³⁸, S. E. Vahsen³³⁹, R. van Tonder³⁴⁰, K. E. Varvell³⁴¹, M. Veronesi³⁴², V. S. Vismaya³⁴³, L. Vitale³⁴⁴, V. Vobbilisetti³⁴⁵, R. Volpe³⁴⁶, M. Wakai³⁴⁷, S. Wallner³⁴⁸, M.-Z. Wang³⁴⁹, A. Warburton³⁵⁰, M. Watanabe³⁵¹, S. Watanuki³⁵², C. Wessel³⁵³, E. Won³⁵⁴, X. P. Xu³⁵⁵, B. D. Yabsley³⁵⁶, W. Yan³⁵⁷, J. Yelton³⁵⁸, K. Yi³⁵⁹, J. H. Yin³⁶⁰, K. Yoshihara³⁶¹, J. Yuan³⁶², Y. Yusa³⁶³, L. Zani³⁶⁴, M. Zeyrek³⁶⁵, J. S. Zhou³⁶⁶, Q. D. Zhou³⁶⁷, L. Zhu³⁶⁸, and R. Žlebčík³⁶⁹

(Belle and Belle II Collaboration)

 (Received 3 November 2025; revised 25 March 2026; accepted 4 May 2026; published 16 June 2026)

We report the first observation of the radiative decay $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ with a statistical significance exceeding 10 standard deviations. The signal is observed in the continuum $e^+e^- \rightarrow c\bar{c}$ process, using combined data samples of 980.4 fb^{-1} from Belle and 427.9 fb^{-1} from Belle II, collected at the KEKB and SuperKEKB asymmetric-energy e^+e^- colliders, respectively. The branching fraction ratio $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0]$ is measured to be $[7.13 \pm 0.70(\text{stat}) \pm 0.26(\text{syst})]\%$. This result provides crucial discrimination between theoretical models of the $D_{s0}^*(2317)^+$ structure.

DOI: [10.1103/vcld-225s](https://doi.org/10.1103/vcld-225s)

The study of exotic hadrons represents a pivotal frontier in particle physics, providing critical insights into the nonperturbative regime of quantum chromodynamics [1–3]. The scalar charm-strange meson $D_{s0}^*(2317)^+$ and the axial-vector meson $D_{s1}(2460)^+$ have garnered significant attention because their masses are significantly below those predicted by the quark model [4–6] for $c\bar{s}$ mesons with their respective J^P quantum numbers. Several theoretical frameworks have been proposed to explain the nature of $D_{s0}^*(2317)^+$ and $D_{s1}(2460)^+$, including molecular states [7–11], quark-antiquark configurations [12–26], tetraquark structures [27–32], and mixed states [33–38]. Despite these efforts, the precise nature of $D_{s0}^*(2317)^+$ and $D_{s1}(2460)^+$ remains unresolved, underscoring the need for new and improved experimental data.

The $D_{s0}^*(2317)^+$ has been a subject of intense experimental and theoretical interest since its discovery by the BABAR Collaboration in the decay mode $D_s^+\pi^0$ [39], later confirmed by CLEO [40] and Belle [41]. Its mass, $2317.8 \pm 0.5 \text{ MeV}/c^2$, lies below the DK threshold, restricting its decay to the isospin-violating strong decay channel $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$. This channel has been measured with a branching fraction of $1.00^{+0.00}_{-0.20}$ by BESIII [42]. Radiative transitions are particularly sensitive probes of the internal structure of such hadrons, as they involve electromagnetic interactions that are well understood [43]. CLEO [40], Belle [41], and BABAR [44] searched for $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ using 13.5 fb^{-1} , 86.9 fb^{-1} , and 232 fb^{-1} data samples, respectively, at center-of-mass (c.m.) energies near 10.6 GeV, but did not find any evidence for this channel. The most restrictive upper limit on the ratio $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0]$ is set to 5.9% at 90% confidence level by CLEO with a $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ signal yield of -6.5 ± 5.2 [40]. Assuming its spin parity is 0^+ , the $D_{s0}^*(2317)^+ \rightarrow D_s^+\gamma$ decay is forbidden. Though the decay

width of $D_{s0}^*(2317)^+$ is unknown, a determination of the branching fraction ratio $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0]$ would provide a direct experimental constraint on various theoretical models used to explain the nature of $D_{s0}^*(2317)^+$. For instance, a ratio in the range of 0.5% to 4.25% would strongly favor molecular interpretations [45–47], while a larger value ($> 8.1\%$) would align more closely with conventional $c\bar{s}$ configurations [21,48]. Therefore, a precise measurement of this ratio is essential to resolve the nature of the $D_{s0}^*(2317)^+$ meson.

In this Letter, we report the first observation of the radiative decay $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$, with $D_s^{*+} \rightarrow D_s^+\gamma$. The rate for this decay is measured relative to the hadronic decay $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$, using 980.4 fb^{-1} of Belle data collected at c.m. energies near the $\Upsilon(nS)$ ($n = 1, \dots, 5$) resonances and 427.9 fb^{-1} of Belle II data collected at or near the c.m. energies of $\Upsilon(4S)$ and 10.75 GeV. Inclusion of charge conjugate states is implicit. The D_s^+ candidates are reconstructed via the $\phi\pi^+$ and $K^+\bar{K}^{*0}$ decay modes, both of which result in the $K^+K^-\pi^+$ final state. The hadronic decay serves as a reference channel, enabling cancellation of the systematic uncertainties associated with D_s^+ and γ selection in the branching fraction ratio measurement.

The Belle detector [49,50] was a large-solid-angle spectrometer that operated at the KEKB asymmetric-energy e^+e^- collider [51,52]. The Belle II detector [53] is a significant upgrade of Belle and operates at the SuperKEKB e^+e^- collider [54]. The z -axis is defined parallel to the e^+ beam at Belle and to the principal axis of the solenoid at Belle II with the interaction point as the origin of the coordinate system.

Data and simulated Monte Carlo (MC) samples for both Belle and Belle II are processed with the Belle II analysis software framework [55–57]. MC simulations are used to optimize selection criteria, investigate background sources, calculate reconstruction efficiencies, and determine the probability density functions (PDFs) employed in fitting the data. The MC events for the continuum $e^+e^- \rightarrow c\bar{c}$ process are generated with KKMC [58] and PYTHIA [59,60], where at least one of the charm quarks hadronizes into a $D_{s0}^*(2317)^+$ meson for the signal events. The $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ and $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$ decays

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

are simulated with the phase space model, while the decay $D_s^{*+} \rightarrow D_s^+ \gamma$ is simulated as a P -wave decay. The decay $D_s^+ \rightarrow K^+ K^- \pi^+$ is modeled based on previous measurements [61,62]. Simulated events undergo detector simulation with Geant3 [63] for Belle and Geant4 [64] for Belle II. The signal MC samples are corrected with a reweighting method based on the measured x_p distribution from the reference channel, where $x_p \equiv p_{D_{s0}^*(2317)^+}^*/p_{\max}^*$ is the reduced momentum of the selected $D_{s0}^*(2317)^+$ candidate. Here, $p_{D_{s0}^*(2317)^+}^*$ is its momentum in the c.m. frame, $p_{\max}^* \equiv \sqrt{E_{\text{beam}}^2/c^2 - M^2[D_{s0}^*(2317)^+]c^2}$ is the maximum kinematically allowed momentum, E_{beam} is the beam energy, and $M[D_{s0}^*(2317)^+]$ is the invariant mass of the $D_{s0}^*(2317)^+$ candidates.

To study backgrounds, we use MC samples generated with the Belle and Belle II configurations, which correspond to four times the sizes of the corresponding datasets. Belle's MC samples include $\Upsilon(1S, 2S, 3S)$ decays, $\Upsilon(4S) \rightarrow B\bar{B}$, $\Upsilon(5S) \rightarrow B_{(s)}^{(*)}\bar{B}_{(s)}^{(*)}, B\bar{B}^{(*)}\pi$, and $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) at c.m. energies of $\sqrt{s} = 10.52, 10.58$, and 10.867 GeV. Belle II's MC samples consist of $e^+e^- \rightarrow q\bar{q}$ and $\Upsilon(4S) \rightarrow B\bar{B}$.

For the signal event selection, tracks are required to satisfy $dr < 0.5$ cm and $|dz| < 3.0$ cm, where dr and dz are transverse and longitudinal impact parameters, respectively. For charged particle identification, information from different subdetectors is combined to form the likelihood $\mathcal{L}_i$ for species i , where $i = \pi$ or K [65,66]. A track with a likelihood ratio $\mathcal{L}_K/(\mathcal{L}_K + \mathcal{L}_\pi) > 0.6 (< 0.4)$ is identified as a kaon (pion). With this selection, for Belle, the kaon (pion) identification efficiency is about 88% (90%), while 8% (8%) of the pions (kaons) are misidentified as kaons (pions); for Belle II, the identification efficiency is about 97% (98%) for kaon (pion), with 1.3% (1.7%) of the pions (kaons) are misidentified as kaons (pions).

The electromagnetic calorimeter clusters not associated with tracks with energy greater than 0.10 GeV in the c.m. frame are regarded as photons. For the signal channel $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}(\rightarrow D_s^+ \gamma_2) \gamma_1$, the energy of γ_1 is required to be greater than 0.22 GeV in the c.m. frame. Additionally, the invariant mass $M_{\gamma_1 \gamma_2}$ must lie outside the region $[0.10, 0.16]$ GeV/ c^2 to exclude $D_{s0}^*(2317)^+ \rightarrow D_s^+ \pi^0$ candidates. For the reference channel $D_{s0}^*(2317)^+ \rightarrow D_s^+ \pi^0$, one of the signal photons should have energy greater than 0.22 GeV in the c.m. frame, and $M(\gamma_1 \gamma_2)$ must be within 15 MeV/ c^2 of the known π^0 mass to form π^0 candidates ($\sim 2.5\sigma$) [61]. For both decay channels, the invariant mass of any combination of a signal photon and any other photon in the event must not fall within 15 MeV/ c^2 of the known π^0 mass [61].

The K^+ , K^- , and π^+ candidates are combined to form D_s^+ candidates. For the decay channel $D_s^+ \rightarrow \phi \pi^+$, we require the invariant mass of the $K^+ K^-$ pair to satisfy

$|M(K^+ K^-) - m(\phi)| < 0.01$ GeV/ c^2 ($\sim 2.5\sigma$), where $m(\phi)$ is the known ϕ mass [61]. For the decay channel $D_s^+ \rightarrow K^+ \bar{K}^{*0}$, the invariant mass of the $K^- \pi^+$ pair must satisfy $|M(K^- \pi^+) - m(K^{*0})| < 0.05$ GeV/ c^2 , where $m(K^{*0})$ is the known K^{*0} mass [61]. The invariant mass of the $K^+ K^- \pi^+$ system must satisfy $|M(K^+ K^- \pi^+) - m(D_s^+)| < 0.01$ GeV/ c^2 , where $m(D_s^+)$ is the known D_s^+ mass [61].

The D_s^+ candidates are combined with a photon to form D_s^{*+} candidates. The mass window of D_s^{*+} is $|M(D_s^+ \gamma_2) - m(D_s^{*+})| < 0.015$ GeV/ c^2 , where $m(D_s^{*+})$ is the known mass of D_s^{*+} [61], and $M(D_s^+ \gamma_2) = M^{\text{rec}}(D_s^+ \gamma_2) - M^{\text{rec}}(K^+ K^- \pi^+) + m(D_s^+)$ is used to cancel the contribution to the mass resolution from the measurement of D_s^+ . Here and below, we use $M^{\text{rec}}(X)$ to indicate the reconstructed invariant mass of the X system. After applying these requirements, there are no candidates for which $M(D_s^+ \gamma_1)$ falls within the D_s^{*+} mass window. Then, the combinations of $D_s^{*+} \gamma_1$ or $D_s^+ \pi^0$ are considered as $D_{s0}^*(2317)^+$ candidates.

To suppress the combinatorial background, we require x_p to be larger than 0.7, which also removes all $D_{s0}^*(2317)^+$ from B decays. The D_s^+ and D_s^{*+} signal purities are 81% (86%) and 68% (63%) for Belle (Belle II) for the radiative decay channel, respectively. All the possible candidates in an event are retained for further analysis, with the multiplicity of 1.03 (1.02) for radiative (hadronic) decay channel in the studied regions shown in Figs. 1 and 2.

We optimize the selection criteria by maximizing the Punzi figure of merit $\varepsilon/(5/2 + \sqrt{N_B})$ [67] in the signal region $[2.29 < M(D_s^{*+} \gamma) < 2.34]$ GeV/ c^2 of the $D_{s0}^*(2317)^+ \rightarrow D_s^{*+} \gamma$ channel, where ε is the detection efficiency. The background yield N_B is estimated in a data-driven way by linearly extrapolating the yield from the upper sideband $[2.35 < M(D_s^{*+} \gamma) < 2.40]$ GeV/ c^2 , as the lower sideband contains a peaking background due to a random photon combining with a real D_s^+ candidate to form the fake D_s^{*+} candidate, while the true γ_2 is mistakenly selected as γ_1 . Similar peaking backgrounds have been reported previously by the CLEO experiment [40]. Since modeling the shapes of these backgrounds is challenging, and they do not affect the extracted signal yields of the $D_{s0}^*(2317)^+$, we exclude this lower mass region in this analysis.

Here, we use $M(D_s^{*+} \gamma) = M^{\text{rec}}(D_s^+ \gamma_1) - M^{\text{rec}}(D_s^+ \gamma_2) + m(D_s^{*+})$, as this cancels the contribution to the mass resolution from the measurement of the D_s^+ . Following a blind analysis strategy, we do not examine the $M(D_s^{*+} \gamma)$ distributions in the signal region until the analysis procedure is finalized.

After applying the aforementioned selections, the $M(D_s^+ \pi^0)$ distributions are presented in Fig. 1, revealing distinct peaks corresponding to the $D_{s0}^*(2317)^+$ state in both datasets. Here, we use $M(D_s^+ \pi^0) = M^{\text{rec}}(D_s^+ \pi^0) - M^{\text{rec}}(K^+ K^- \pi^+) + m(D_s^+) - M^{\text{rec}}(\gamma_1 \gamma_2) + m(\pi^0)$, as this




FIG. 1. Invariant mass distributions of $D_s^+ \pi^0$ from (a) Belle and (b) Belle II data. The data samples are represented by the dots with error bars. The blue curves, green dotted curves, and green filled areas are the fitted total PDFs, total backgrounds, and combinatorial backgrounds, respectively. The areas between total and combinatorial backgrounds are from the fitted cross-feed contributions. The distributions from the normalized D_s^+ and π^0 mass sidebands are shown with yellow blank and red slash filled histograms, respectively.

cancels the contributions to the mass resolutions from the measurements of the D_s^+ and π^0 . The TopoAna package [68] is used for the background study. Based on the study of the MC samples, apart from the combinatorial background, the $D_{s1}(2460)^+ \rightarrow D_s^{*+} \pi^0$ decay with the photon from D_s^{*+} missed can introduce an excess under the $D_{s0}^*(2317)^+$ peak (denoted as the cross feed). The distributions from the normalized D_s^+ and π^0 mass sidebands $[|M(K^+ K^- \pi^+) - m(D_s^+) \pm 0.04| < 0.01 \text{ GeV}/c^2$ and $|M(\gamma\gamma) - m(\pi^0) \pm 0.05| < 0.0075 \text{ GeV}/c^2]$ are shown with yellow blank and red slash filled histograms in Fig. 1, which exhibit no peaking structures; i.e., the background from the $D_{s0}^*(2317)^+ \rightarrow D_s^{*+} \gamma$ channel is negligible.




FIG. 2. Simultaneous fits to the $M(D_s^{*+} \gamma)$ distributions from (a) Belle and (b) Belle II. The blue and violet curves are the best fit results and the fitted total background PDFs, respectively. The filled green areas are the fitted combinatorial backgrounds. The histograms in red slashes represent the normalized D_s^{*+} sidebands.

The signal yields of $D_{s0}^*(2317)^+ \rightarrow D_s^+ \pi^0$ are extracted from the unbinned extended maximum-likelihood fits to the $M(D_s^+ \pi^0)$ spectra. In each fit, the signal PDF is represented by a crystal ball (CB) function [69] convolved with a triple-Gaussian function, whose parameters, except the mean values of the CB functions, are fixed according to signal MC simulations. The cross-feed PDF is constructed from smoothed histograms of MC events. The combinatorial backgrounds are described by the second-order Chebyshev polynomials. The yields of these components are floated in the fits, and the fit results are shown in Fig. 1. The fit method is validated by the MC samples. The similar fits are performed to $M(D_s^+ \pi^0)$ spectra from different x_p bins to measure the x_p distribution of $D_{s0}^*(2317)^+$. The obtained efficiency-corrected x_p distribution is used to correct the MC simulation. The fitted yields of the hadronic

decay channel $N_{\text{exp}}^{\text{fit}}(D_s^+\pi^0)$ are 10820 ± 230 for Belle and 6108 ± 163 for Belle II. For events with $x_p > 0.7$, the detection efficiencies $\epsilon_{\text{exp}}(D_s^+\pi^0)$ are 4.6% and 5.2% for Belle and Belle II, respectively. Here and after, the subscript exp indicates an experiment (Belle or Belle II).

For the $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ channel, the $M(D_s^{*+}\gamma)$ spectra from Belle and Belle II data are presented in Fig. 2, where the $D_{s0}^*(2317)^+$ signal peak is clearly visible in both plots. According to the studies done on MC simulations [68], we do not anticipate any peaking contribution from $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$ or from $D_{s1}(2460)^+ \rightarrow D_s^{*+}\pi^0, D_s^{*+}\gamma$, and $D_{s0}^*(2317)^+\gamma$ decays. Furthermore, no peaking in the $D_{s0}^*(2317)^+$ signal region is found in the normalized D_s^{*+} mass sidebands $[|M(D_s^+\gamma) - m(D_s^{*+}) \pm 0.05| < 0.015 \text{ GeV}/c^2]$. A small peaking background, which we label as “broken signal,” may arise when a correctly reconstructed $D_s^+\gamma_1$ candidate is combined with a wrongly selected γ_2 .

We extract the branching fraction ratio $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0]$, denoted $\mathcal{R}$, through a simultaneous unbinned extended maximum-likelihood fit to the $M(D_s^{*+}\gamma)$ spectra from Belle and Belle II, as shown in Fig. 2. Each $D_{s0}^*(2317)^+$ signal PDF is modeled by a CB function convolved with a triple-Gaussian function, while the corresponding broken signal contribution is described by an asymmetric Gaussian. All parameters of the broken signal and signal PDF, as well as the ratios of their yields, are fixed from MC simulations, except for the mean values of the CB functions. The yield ratio of the broken signal to signal component is 7.5% (9.3%) for Belle (Belle II). The combinatorial backgrounds are described by first-order polynomials. The mass resolution is approximately 7 MeV for both Belle and Belle II. The value of $\mathcal{R}$ is shared as a common free parameter in the simultaneous fit, while the $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ signal yields are set according to $N_{\text{exp}}(D_s^{*+}\gamma) = \mathcal{R}N_{\text{exp}}^{\text{fit}}(D_s^+\pi^0)\epsilon_{\text{exp}}(D_s^{*+}\gamma)/\epsilon_{\text{exp}}(D_s^+\pi^0)$ separately for Belle and Belle II. Here, $\epsilon_{\text{exp}}(D_s^{*+}\gamma)$ is the detection efficiency of $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ decay, which is 4.2% for Belle and 4.6% for Belle II. The fit results are shown in Fig. 2. The fitted masses of the $D_{s0}^*(2317)^+$ in the Belle and Belle II datasets differ by $3.6 \pm 1.5 \text{ MeV}/c^2$. This mass difference is consistent with known discrepancies in low-energy photon calibration between the two detectors and does not affect the $\mathcal{R}$ measurement, as the simultaneous fit accounts for this offset through independent mean values for each dataset. The fitted $\mathcal{R}$ value is $[7.13 \pm 0.70(\text{stat})]\%$. The corresponding $N_{\text{exp}}(D_s^{*+}\gamma)$ are 712 ± 69 and 387 ± 38 for Belle and Belle II, respectively.

The significance of $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ is 10.1σ , estimated from the negative log-likelihood ratio $-2\ln(\mathcal{L}_0/\mathcal{L}_{\text{max}}) = 111.9$ [70] with the difference in degrees of freedom ($\Delta\text{d.o.f.} = 3$) and the systematic uncertainty discussed below considered. Here, $\mathcal{L}_0$ and

$\mathcal{L}_{\text{max}}$ represent the maximized likelihoods of the simultaneous fits without and with the $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ signal components, respectively. Systematic uncertainties are incorporated by convolving the likelihood ratio distribution with a Gaussian function whose width corresponds to the total systematic uncertainty. We also perform separate fits to the Belle and Belle II data using the same fit components as those in the simultaneous fit. The fitted signal yields $N_{\text{exp}}^{\text{fit}}(D_s^{*+}\gamma)$ are 742 ± 82 and 348 ± 69 for Belle and Belle II, respectively. The corresponding $\mathcal{R}$ values are $[7.43 \pm 0.83(\text{stat})]\%$ and $[6.43 \pm 1.27(\text{stat})]\%$, demonstrating good consistency between the results of the simultaneous fit and the fits to each individual dataset.

The systematic uncertainties due to D_s^+ and γ selection cancel in the $\mathcal{R}$ measurement. The dominant systematic uncertainties are from the fit model and x_p weighting. All systematic sources are described below, and the resulting percent variations relative to the nominal fit are listed in Table I.

To characterize possible systematic effects in the $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$ reference channel, (a) the order of the background polynomials and the fit range are varied, (b) the widths of the triple-Gaussian functions are increased by 1σ , and (c) the resolution PDF variations are also propagated into the modeling of the $D_{s1}(2460)^+ \rightarrow D_s^{*+}\pi^0$ cross-feed components. Finally, the differences in the fitted $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$ yields are taken as systematic uncertainties of $N_{\text{exp}}^{\text{fit}}(D_s^+\pi^0)$, which are 1.3% (1.0%), 0.7% (1.5%), and 0.8% (0.3%) from the fit region and background PDF, resolution, and cross-feed PDF for Belle (Belle II), respectively.

A series of pseudoexperiments is conducted to estimate the systematic uncertainty contribution to $\mathcal{R}$ from the $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$ channel. In each trial, we randomly fluctuate the $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$ yields for both Belle and Belle II by sampling from Gaussian distributions. Each Gaussian distribution is constructed with its mean value set to the corresponding nominal $D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0$ yield and its standard deviation equal to the systematic uncertainty of $N_{\text{exp}}^{\text{fit}}(D_s^+\pi^0)$. A simultaneous fit similar to the

TABLE I. The summary of the systematic uncertainties of the measurement of the branching fraction ratio $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^+\pi^0]$ (in %).

Source	$D_s^+\pi^0$	$D_s^{*+}\gamma$
Fit region and background PDF	0.8	1.3
Fixed PDF parameters	0.7	2.5
Cross feed or broken signal	0.6	0.7
x_p reweighting		0.5
$\cos\theta_{\gamma_1\gamma_2}$ reweighting		1.7
MC sample size		0.5
Sum		3.6

nominal fit to the data described above is then performed to the $M(D_s^{*+}\gamma)$ distributions from data for each set of the pseudoyields of $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\pi^0$. From these results, an ensemble of Gaussian-distributed varied $\mathcal{R}$ values is obtained whose width is taken as the systematic uncertainty on $\mathcal{R}$ from $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\pi^0$ decay.

We characterize systematic effects in the signal channel $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ fits by examining the changes in fitted $\mathcal{R}$ values in the simultaneous fit to $M(D_s^{*+}\gamma)$ distributions from data after (a) varying the order of the background polynomials and the fit range, (b) increasing the widths of the triple-Gaussian functions by 1σ , and (c) adjusting the ratios and widths of broken signal to signal yields by 2σ to conservatively estimate the systematic uncertainty. The differences of the fitted $\mathcal{R}$ values from the nominal result are taken as systematic uncertainties.

To estimate the uncertainty due to x_p reweighting, we vary the polynomial order when fitting the efficiency-corrected x_p distribution and reweight the signal MC samples accordingly. Then, the new detection efficiencies derived from the reweighted signal MC samples are used in the simultaneous fit to $M(D_s^{*+}\gamma)$ distributions from data. The change in the fitted $\mathcal{R}$ value from the nominal result is taken as the systematic uncertainty.

By reweighting the $\cos\theta_{\gamma_1\gamma_2}$ distribution in the $D_{s0}^*(2317)^+$ signal MC samples to match a $1 + \cos^2\theta_{\gamma_1\gamma_2}$ distribution, we obtain a new set of detection efficiencies. Here, $\theta_{\gamma_1\gamma_2}$ is the angle between the two photons in the D_s^* rest frame. The corresponding systematic uncertainty is estimated using a method similar to that for the x_p reweighting.

The systematic uncertainty of detection efficiencies due to the limited size of the signal MC sample is estimated by $\sqrt{(1-\varepsilon)\varepsilon/N}$, where ε and N are the detection efficiency and number of simulated signal events, respectively. By varying the detection efficiencies by 1σ in the simultaneous fit to $M(D_s^{*+}\gamma)$ from data, the change in the fitted $\mathcal{R}$ from the nominal result is taken as the systematic uncertainty.

Assuming that all the systematic uncertainties detailed above are independent, they are added in quadrature to obtain the total systematic uncertainty of 3.2%, as listed in Table I.

In summary, using combined data samples of 980 fb⁻¹ from Belle and 428 fb⁻¹ from Belle II collected at the KEKB and SuperKEKB asymmetric-energy e^+e^- colliders, respectively, we have made the first observation of the radiative decay $D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma$ in the continuum $e^+e^- \rightarrow c\bar{c}$ process with a significance exceeding 10 standard deviations. A comparison between theoretical predictions and the measured $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\pi^0]$ value is presented in Fig. 3. The branching fraction ratio $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\pi^0]$ is measured to be $[7.13 \pm 0.70(\text{stat}) \pm 0.26(\text{syst})]\%$, which is generally




FIG. 3. Comparison of the measured $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\pi^0]$ with theoretical predictions. The theoretical approaches of the references are effective field theory for Liu [26] and Cleven [11], traditional quark model for Godfrey [48], heavy quark symmetries for Colangelo [24], phenomenological for Azimov [25], light front quark model for Ke [71], effective Lagrangian with chiral symmetry for Bardeen [21], effective Lagrangian for Faessler [45], heavy quark flavor symmetry for Fu [46], and chiral Lagrangian with coupled-channel dynamics for Lutz [47]. The prediction of Ke [71] is a lower limit indicated by an arrow. The uncertainty of the experimental measurement is the quadrature sum of statistical and systematic uncertainties.

larger than theoretical predictions suggesting $D_{s0}^*(2317)^+$ as a molecular state [45–47], while smaller than the $c\bar{s}$ state assumption under the quark model [48] and under effective field theory for both assumptions [11,26]. However, predictions based on the light front quark model [71] and chiral quark model [21] agree with our measurement under the pure $c\bar{s}$ state expectation. While most theoretical models address an anomalously low mass of the $D_{s0}^*(2317)^+$ and provide estimates of its width, the intrinsic width cannot currently be measured due to limited experimental resolution. Our result provides a complementary input that will help to further discriminate among competing explanations for the nature of the $D_{s0}^*(2317)^+$ and motivates dedicated theoretical calculations of the ratio $\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\gamma]/\mathcal{B}[D_{s0}^*(2317)^+ \rightarrow D_s^{*+}\pi^0]$ under various hypotheses.

Acknowledgments—We thank the SuperKEKB team for delivering high-luminosity collisions; the KEK cryogenics group for the efficient operation of the detector solenoid magnet and IBBelle on site; the KEK Computer Research Center for on-site computing support; the NII for SINET6 network support; and the raw-data centers hosted by BNL, DESY, GridKa, IN2P3, INFN, PNNL/EMSL, and the University of Victoria. This work, based on data collected using the Belle II detector, which was built and commissioned prior to March 2019, and data collected using the Belle detector, which was operated until June 2010, was

supported by Higher Education and Science Committee of the Republic of Armenia Grant No. 23LCG-1C011; Australian Research Council and Research Grants No. DP200101792, No. DP210101900, No. DP210102831, No. DE220100462, No. LE210100098, and No. LE230100085; Austrian Federal Ministry of Education, Science and Research, Austrian Science Fund (FWF) [Grants DOI: 10.55776/P34529, DOI: 10.55776/J4731, DOI: 10.55776/J4625, DOI: 10.55776/M3153, and DOI: 10.55776/PAT1836324] and Horizon 2020 ERC Starting Grant No. 947006 “InterLeptons”; Natural Sciences and Engineering Research Council of Canada, Digital Research Alliance of Canada, and Canada Foundation for Innovation; National Key R&D Program of China under Contracts No. 2024YFA1610503 and No. 2024YFA1610504, National Natural Science Foundation of China and Research Grants No. 11575017, No. 11761141009, No. 11705209, No. 11975076, No. 12135005, No. 12150004, No. 12161141008, No. 12405099, No. 12475093, No. 12175041, and No. 12405102 and Shandong Provincial Natural Science Foundation Project ZR2022JQ02; the Czech Science Foundation Grant No. 22-18469S; Regional funds of EU/MEYS: OPJAK FORTE CZ.02.01.01/00/22_008/0004632 and Charles University Grant Agency Project No. 246122; European Research Council, Seventh Framework PIF-GA-2013-622527, Horizon 2020 ERC-Advanced Grants No. 267104 and No. 884719, Horizon 2020 ERC-Consolidator Grant No. 819127, Horizon 2020 Marie Skłodowska-Curie Grant Agreements No. 700525 “NIOBE” and No. 101026516, and Horizon 2020 Marie Skłodowska-Curie RISE project JENNIFER2 Grant Agreement No. 822070 (European grants); L’Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) du CNRS and L’Agence Nationale de la Recherche (ANR) under Grant No. ANR-23-CE31-0018 (France); BMFT, DFG, HGF, MPG, and AvH Foundation (Germany); Department of Atomic Energy under Project Identification No. RTI 4002, Department of Science and Technology, and UPES SEED funding Programs No. UPES/R&D-SEED-INFRA/17052023/01 and No. UPES/R&D-SOE/20062022/06 (India); Israel Science Foundation Grant No. 2476/17, U.S.-Israel Binational Science Foundation Grant No. 2016113, and Israel Ministry of Science Grant No. 3-16543; Istituto Nazionale di Fisica Nucleare and the Research Grants BELLE2, and the ICSC—Centro Nazionale di Ricerca in High Performance Computing, Big Data and Quantum Computing, funded by European Union—NextGenerationEU; Japan Society for the Promotion of Science, Grant-in-Aid for Scientific Research Grants No. 16H03968, No. 16H03993, No. 16H06492, No. 16K05323, No. 17H01133, No. 17H05405,

No. 18K03621, No. 18H03710, No. 18H05226, No. 19H00682, No. 20H05850, No. 20H05858, No. 22H00144, No. 22K14056, No. 22K21347, No. 23H05433, No. 26220706, and No. 26400255 and the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan; National Research Foundation (NRF) of Korea Grants No. 2021R1-F1A-1064008, No. 2022R1-A2C-1003993, No. 2022R1-A2C-1092335, No. RS-2016-NR017151, No. RS-2018-NR031074, No. RS-2021-NR060129, No. RS-2023-00208693, No. RS-2024-00354342, and No. RS-2025-02219521, Radiation Science Research Institute, Foreign Large-Size Research Facility Application Supporting project, the Global Science Experimental Data Hub Center, the Korea Institute of Science and Technology Information (K25L2M2C3), and KREONET/GLORIAD; Universiti Malaya RU grant, Akademi Sains Malaysia, and Ministry of Education Malaysia; Frontiers of Science Program Contracts No. FOINS-296, No. CB-221329, No. CB-236394, No. CB-254409, and No. CB-180023 and SEP-CINVESTAV Research Grant No. 237 (Mexico); the Polish Ministry of Science and Higher Education and the National Science Center; the Ministry of Science and Higher Education of the Russian Federation and the HSE University Basic Research Program, Moscow; University of Tabuk Research Grants No. S-0256-1438 and No. S-0280-1439 (Saudi Arabia) and Researchers Supporting Project Number (RSPD2025R873), King Saud University, Riyadh, Saudi Arabia; Slovenian Research Agency and Research Grants No. J1-50010 and No. P1-0135; Ikerbasque, Basque Foundation for Science, State Agency for Research of the Spanish Ministry of Science and Innovation through Grant No. PID2022-136510NB-C33, Spain, Agencia Estatal de Investigación, Spain Grant No. RYC2020-029875-I and Generalitat Valenciana, Spain Grant No. CIDEGENT/2018/020; The Knut and Alice Wallenberg Foundation (Sweden), Contracts No. 2021.0174, No. 2021.0299, and No. 2023.0315; National Science and Technology Council and Ministry of Education (Taiwan); Thailand Center of Excellence in Physics; TUBITAK ULAKBIM (Turkey); National Research Foundation of Ukraine, Project No. 2020.02/0257 and Ministry of Education and Science of Ukraine; the U.S. National Science Foundation and Research Grants No. PHY-1913789 and No. PHY-2111604 and the U.S. Department of Energy and Research Awards No. DE-AC06-76RLO1830, No. DE-SC0007983, No. DE-SC0009824, No. DE-SC0009973, No. DE-SC0010007, No. DE-SC0010073, No. DE-SC0010118, No. DE-SC0010504, No. DE-SC0011784, No. DE-SC0012704, No. DE-SC0019230, No. DE-SC0021274, No. DE-SC0021616, No. DE-SC0022350, No. DE-SC0023470; and the Vietnam Academy of Science and Technology (VAST) under Grants No. NVCC.05.02/25-25 and No. DL0000.05/26-27.

These acknowledgements are not to be interpreted as an endorsement of any statement made by any of our institutes, funding agencies, governments, or their representatives.

Data availability—The data are not publicly available. The data are available from the authors upon reasonable request.

-
- [1] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C. P. Shen, C. E. Thomas, A. Vairo, and C. Z. Yuan, *Phys. Rep.* **873**, 1 (2020).
 - [2] H. X. Chen, W. Chen, X. Liu, Y. R. Liu, and S. L. Zhu, *Rep. Prog. Phys.* **80**, 076201 (2017).
 - [3] F. K. Guo, C. Hanhart, U. G. Meißner, Q. Wang, Q. Zhao, and B. S. Zou, *Rev. Mod. Phys.* **90**, 015004 (2018); **94**, 029901(E) (2022).
 - [4] S. Godfrey and N. Isgur, *Phys. Rev. D* **32**, 189 (1985).
 - [5] S. Godfrey and R. Kokoski, *Phys. Rev. D* **43**, 1679 (1991).
 - [6] M. Di Pierro and E. Eichten, *Phys. Rev. D* **64**, 114004 (2001).
 - [7] Y. Q. Chen and X. Q. Li, *Phys. Rev. Lett.* **93**, 232001 (2004).
 - [8] F. K. Guo, P. N. Shen, H. C. Chiang, R. G. Ping, and B. S. Zou, *Phys. Lett. B* **641**, 278 (2006).
 - [9] D. Gamermann, E. Oset, D. Strottman, and M. J. Vicente Vacas, *Phys. Rev. D* **76**, 074016 (2007).
 - [10] D. Gamermann, L. R. Dai, and E. Oset, *Phys. Rev. C* **76**, 055205 (2007).
 - [11] M. Cleven, H. W. Griebhammer, F. K. Guo, C. Hanhart, and U. G. Meißner, *Eur. Phys. J. A* **50**, 149 (2014).
 - [12] E. van Beveren and G. Rupp, *Phys. Rev. Lett.* **91**, 012003 (2003).
 - [13] E. van Beveren and G. Rupp, *Eur. Phys. J. C* **32**, 493 (2004).
 - [14] S. Coito, G. Rupp, and E. van Beveren, *Phys. Rev. D* **84**, 094020 (2011).
 - [15] D. S. Hwang and D. W. Kim, *Phys. Lett. B* **601**, 137 (2004).
 - [16] D. S. Hwang and D. W. Kim, *J. Phys. Conf. Ser.* **9**, 63 (2005).
 - [17] Y. A. Simonov and J. A. Tjon, *Phys. Rev. D* **70**, 114013 (2004).
 - [18] I. W. Lee, T. Lee, D. P. Min, and B. Y. Park, *Eur. Phys. J. C* **49**, 737 (2007).
 - [19] Z. Y. Zhou and Z. Xiao, *Phys. Rev. D* **84**, 034023 (2011).
 - [20] A. M. Badalian, Y. A. Simonov, and M. A. Trusov, *Phys. Rev. D* **77**, 074017 (2008).
 - [21] W. A. Bardeen, E. J. Eichten, and C. T. Hill, *Phys. Rev. D* **68**, 054024 (2003).
 - [22] M. A. Nowak, M. Rho, and I. Zahed, *Acta Phys. Pol. B* **35**, 2377 (2004), <https://inspirehep.net/literature/623008>.
 - [23] E. E. Kolomeitsev and M. F. M. Lutz, *Phys. Lett. B* **582**, 39 (2004).
 - [24] P. Colangelo and F. De Fazio, *Phys. Lett. B* **570**, 180 (2003).
 - [25] Y. I. Azimov and K. Goeke, *Eur. Phys. J. A* **21**, 501 (2004).
 - [26] X. Liu, Y. M. Yu, S. M. Zhao, and X. Q. Li, *Eur. Phys. J. C* **47**, 445 (2006).
 - [27] H. Y. Cheng and W. S. Hou, *Phys. Lett. B* **566**, 193 (2003).
 - [28] V. Dmitrasinovic, *Phys. Rev. D* **70**, 096011 (2004).
 - [29] V. Dmitrasinovic, *Phys. Rev. D* **86**, 016006 (2012).
 - [30] A. Hayashigaki and K. Terasaki, *Prog. Theor. Phys.* **114**, 1191 (2006).
 - [31] L. Maiani, F. Piccinini, A. D. Polosa, and V. Riquer, *Phys. Rev. D* **71**, 014028 (2005).
 - [32] M. E. Bracco, A. Lozea, R. D. Matheus, F. S. Navarra, and M. Nielsen, *Phys. Lett. B* **624**, 217 (2005).
 - [33] T. E. Browder, S. Pakvasa, and A. A. Petrov, *Phys. Lett. B* **578**, 365 (2004).
 - [34] J. Lu, X. L. Chen, W. Z. Deng, and S. L. Zhu, *Phys. Rev. D* **73**, 054012 (2006).
 - [35] P. Bicudo, *Phys. Rev. D* **74**, 036008 (2006).
 - [36] A. Martinez Torres, L. R. Dai, C. Koren, D. Jido, and E. Oset, *Phys. Rev. D* **85**, 014027 (2012).
 - [37] D. Mohler, C. B. Lang, L. Leskovec, S. Prelovsek, and R. M. Woloshyn, *Phys. Rev. Lett.* **111**, 222001 (2013).
 - [38] M. N. Tang, Y. H. Lin, F. K. Guo, C. Hanhart, and U. G. Meißner, *Commun. Theor. Phys.* **75**, 055203 (2023).
 - [39] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **90**, 242001 (2003).
 - [40] D. Besson *et al.* (CLEO Collaboration), *Phys. Rev. D* **68**, 032002 (2003); **75**, 119908(E) (2007).
 - [41] Y. Mikami *et al.* (Belle Collaboration), *Phys. Rev. Lett.* **92**, 012002 (2004).
 - [42] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **97**, 051103 (2018).
 - [43] H. X. Chen, W. Chen, X. Liu, Y. R. Liu, and S. L. Zhu, *Rep. Prog. Phys.* **86**, 026201 (2023).
 - [44] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. D* **74**, 032007 (2006).
 - [45] A. Faessler, T. Gutsche, V. E. Lyubovitskij, and Y. L. Ma, *Phys. Rev. D* **76**, 014005 (2007).
 - [46] H. L. Fu, H. W. Griebhammer, F. K. Guo, C. Hanhart, and U. G. Meißner, *Eur. Phys. J. A* **58**, 70 (2022).
 - [47] M. F. M. Lutz and M. Soyeur, *Nucl. Phys. A* **813**, 14 (2008).
 - [48] S. Godfrey, *Phys. Lett. B* **568**, 254 (2003).
 - [49] A. Abashian *et al.* (Belle Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 117 (2002).
 - [50] J. Brodzicka *et al.*, *Prog. Theor. Exp. Phys.* **2012**, 04D001 (2012).
 - [51] S. Kurokawa and E. Kikutani, *Nucl. Instrum. Methods Phys. Res., Sect. A* **499**, 1 (2003), <https://inspirehep.net/literature/572361>.
 - [52] T. Abe *et al.*, *Prog. Theor. Exp. Phys.* **2013**, 03A001 (2013).
 - [53] T. Abe *et al.* (Belle II Collaboration), [arXiv:1011.0352](https://arxiv.org/abs/1011.0352).
 - [54] K. Akai, K. Furukawa, and H. Koiso, *Nucl. Instrum. Methods Phys. Res., Sect. A* **907**, 188 (2018).
 - [55] T. Kuhr, C. Pulvermacher, M. Ritter, T. Hauth, and N. Braun (Belle II Software Framework Group), *Comput. Software Big Sci.* **3**, 1 (2019).
 - [56] Belle II Collaboration, Belle II Analysis Software Framework (basf2), [10.5281/zenodo.5574115](https://zenodo.org/record/5574115).
 - [57] M. Gelb *et al.*, *Comput. Software Big Sci.* **2**, 9 (2018).
 - [58] S. Jadach, B. F. L. Ward, and Z. Was, *Comput. Phys. Commun.* **130**, 260 (2000).
 - [59] T. Sjöstrand, P. Edén, C. Friberg, L. Lönnblad, G. Miu, S. Mrenna, and E. Norrbin, *Comput. Phys. Commun.* **135**, 238 (2001).
 - [60] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, *Comput. Phys. Commun.* **191**, 159 (2015).

- [61] S. Navas *et al.* (Particle Data Group), *Phys. Rev. D* **110**, 030001 (2024).
- [62] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **104**, 012016 (2021).
- [63] R. Brun *et al.*, Geant 3: User's guide Geant 3.10, Geant 3.11, Report No. DD/EE/84-1, CERN, 1984.
- [64] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [65] E. Nakano, *Nucl. Instrum. Methods Phys. Res., Sect. A* **494**, 402 (2002).
- [66] I. Adachi *et al.* (Belle II Collaboration), *Eur. Phys. J. C* **85**, 1237 (2025).
- [67] G. Punzi, eConf **C030908**, MODT002 (2003), <https://inspirehep.net/literature/634798>.
- [68] X. Zhou, S. Du, G. Li, and C. Shen, *Comput. Phys. Commun.* **258**, 107540 (2021).
- [69] M. Oreglia, Report No. SLAC-0236, SLAC, 1980.
- [70] S. S. Wilks, *Ann. Math. Stat.* **9**, 60 (1938).
- [71] H. W. Ke, X. Q. Li, and Y. L. Shi, *Phys. Rev. D* **87**, 054022 (2013).