

Low Energy Limit of Banks-Fischler-Shenker-Susskind Quantum Mechanics

Óscar J. C. Dias^{*}

STAG research centre and Mathematical Sciences, *University of Southampton*,
University Road Southampton SO17 1BJ, United Kingdom

Jorge E. Santos[†]

DAMTP, Centre for Mathematical Sciences, *University of Cambridge*,
Wilberforce Road, Cambridge CB3 0WA, United Kingdom

 (Received 3 October 2024; revised 3 April 2025; accepted 28 May 2025; published 12 June 2025)

We investigate the low energy regime of BFSS quantum mechanics using its holographic dual. We identify three distinct thermodynamic phases (black holes) and analyze their thermodynamic properties extensively, including phase transitions amongst the several phases. While the properties of the canonical ensemble aligns with existing conjectures on BFSS thermodynamics, we uncover intriguing and unexpected behavior in the microcanonical ensemble. Specifically, for sufficiently low energies, we observe the dominance of the localized phase. Surprisingly, we also identify an energy range where the non-uniform phase becomes dominant. The transition between these phases is mediated by a Kol-type topology-changing phenomenon.

DOI: [10.1103/wws8-m34w](https://doi.org/10.1103/wws8-m34w)

Introduction—The holographic principle, also known as gauge-gravity duality, connects certain quantum field theories (QFT) to higher-dimensional quantum gravity, and it is foundational in modern high-energy physics. In this context, finite temperature phases in the QFT correspond one-to-one with black hole phases in the gravitational theory, including their thermodynamic properties. Maldacena’s original duality, proposed in [1], relates ten-dimensional type IIB supergravity on $\text{AdS}_5 \times S^5$ to four-dimensional $\mathcal{N} = 4$ supersymmetric Yang-Mills (SYM) with large N and strong ’t Hooft coupling. Despite substantial supporting evidence, no formal proof of this duality exists, as the gravitational descriptions apply only when the gauge theory duals are strongly coupled [1–3]. Additionally, $\mathcal{N} = 4$ SYM, though highly symmetric, is impractical to simulate with standard lattice Monte Carlo methods. Its four-dimensional nature introduces computational complexity, and its supersymmetric ultraviolet (UV) regime includes many fermionic degrees of freedom, complicating standard path integral formulations due to the sign problem [4] (recently, there has been however significant progress dealing with these obstacles [5,6]).

Shortly after [1], Ref. [7] proposed a network of additional holographic dualities pairing maximally supersymmetric theories with their gravitational duals. These were identified by studying D-branes in string theory and decoupling the brane world volume theory from gravity. This Letter focuses on the simplest duality of [7], namely, between the low energy limit of D0-branes and maximally supersymmetric quantum mechanics (BFSS) [8]. Recent years have seen the development of novel techniques for analyzing the gauge theory side of this duality at strong coupling using computer lattice simulations [5,9–19]. Our study will offer key insights into the gravitational aspect of the duality, crucial for understanding BFSS theory in its strongly coupled regime and enabling precise holography matchings.

BFSS is a gauge theory that includes finite $N \times N$ traceless Hermitian bosonic degrees of freedom X^j and fermionic degrees of freedom θ^α , transforming as spinors of $SO(9)$. The action is

$$S_{\text{BFSS}} = \frac{N}{2\lambda} \int dt \text{Tr} \left\{ (D_t X^j)^2 + \theta^\alpha D_t \theta^\alpha + \frac{1}{2} [X^j, X^k]^2 + i \theta^\alpha \gamma_{\alpha\beta}^j [\theta^\beta, X^j] \right\}, \quad (1)$$

where $\lambda \equiv g_{\text{YM}}^2 N$ is the ’t Hooft coupling, g_{YM} is the gauge coupling, and D_t is the gauge covariant derivative with respect to time t . Summation over spatial indices $i, j = 1, \dots, 9$ and spinor indices $\alpha, \beta = 1, \dots, 16$ is implicit. λ has units of energy cubed so, at finite temperature $\mathcal{T}_H$, the

^{*}Contact author: ojcd1r13@soton.ac.uk

[†]Contact author: jss55@cam.ac.uk

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article’s title, journal citation, and DOI. Funded by SCOAP³.

system's thermodynamics depends on the dimensionless parameters $\tau \equiv \mathcal{T}_H/\lambda^{1/3}$ and N .

According to [7], a dual gravitational description involving 11-dimensional supergravity is expected for

$$N^{-5/6} \ll \tau \ll 1, \quad (2)$$

where the lower bound ensures black holes are larger than the 11-dimensional Planck length and the upper bound ensures small curvatures in string units. When a classical description is adequate, the action for this supergravity is

$$S_{11D} = \frac{1}{16\pi G_{11}} \int \left(\eta R + \frac{1}{2} dC \wedge \star dC - \frac{1}{6} C \wedge dC \wedge dC \right), \quad (3)$$

where η is the spacetime volume form, R is the Ricci scalar, C is a three-form gauge potential, $\star$ is the standard 11-dimensional Hodge dual operation, and G_{11} is the 11-dimensional Newton constant. In this Letter, we will focus on geometries with $C = 0$.

Following [7], we search for solutions to the equations of motion derived from (3) that asymptotically approach, at large r , the metric

$$ds^2 = -\frac{r^7}{\mathcal{R}^7} dt^2 + dr^2 + \frac{\mathcal{R}^7}{r^7} \left(dz + \frac{r^7}{\mathcal{R}^7} dt \right)^2 + r^2 d\Omega_8^2, \quad (4)$$

where $d\Omega_8^2$ is the line element of a unit radius sphere S^8 and $z \sim z + L_{11}$ with $L_{11} = 2\pi g_s \ell_s$ parametrizing the M-theory circle (ℓ_s , g_s are the string length and string coupling constant, respectively). The map between BFSS and 11-dimensional supergravity is

$$g_s = \frac{4\pi^2 \ell_s^3}{N} \lambda, \quad G_{11} = 16\pi^7 g_s^3 \ell_s^9, \quad \left(\frac{\mathcal{R}}{\ell_s} \right)^7 = 60\pi^3 g_s N. \quad (5)$$

For sufficiently high temperatures, namely, when $N^{-5/6} \lesssim \tau \ll 1$, [7] conjectured that the dominant phase retains translational invariance in z , has horizon topology $S^1 \times S^8$, and is simply given by

$$ds^2 = -\frac{r^7}{\mathcal{R}^7} f dt^2 + \frac{dr^2}{f} + \frac{\mathcal{R}^7}{r^7} \left[dz + \frac{r^7}{\mathcal{R}^7} f(r) dt \right]^2 + r^2 d\Omega_8^2, \quad (6a)$$

with

$$f(r) = 1 - \frac{r_0^7}{r^7} \quad \text{where} \quad \left(\frac{r_0}{\ell_s} \right)^5 = \frac{120\pi^2}{49} (2\pi g_s N)^{5/3} \tau^2. \quad (6b)$$

This solution describes a stack of uniform D0-branes at a finite temperature τ extended along the M-theory circle.

However, for sufficiently low temperatures, specifically $N^{-25/21} \ll \tau \lesssim N^{-5/6}$, [7] conjectured that the relevant phase should be a localized black hole on the M-theory circle. In this Letter, we will show that while this expectation is met in the canonical ensemble, it surprisingly fails in the microcanonical ensemble. Indeed, we will present compelling numerical evidence for the existence of a dominant phase, within a specific range of energies, that breaks translational invariance in z while maintaining a horizon topology of $S^1 \times S^8$. Moreover, in the canonical ensemble, we will determine the precise temperature at which the first-order transition occurs between localised black holes and the translationally preserving phase.

We will describe the construction of black hole phases in supergravity, map them to BFSS thermodynamic phases, and present BFSS phase diagrams in both canonical and microcanonical ensembles.

Constructing the black hole phases—The black holes/strings we aim to construct can be derived from Ricci-flat, *static* configurations with Kaluza-Klein asymptotics $\mathbb{R}^{1,9} \times S^1_L$. We will detail the construction of these configurations and then present the mapping that transforms such solutions to those approaching (4) at large r .

We use the DeTurck trick, introduced in [20] and reviewed in [21,22]. The method starts by considering the so-called Einstein-DeTurck equation,

$$R_{ab} - \nabla_{(a} \xi_{b)} = 0, \quad (7)$$

with ξ being the DeTurck vector, given by

$$\xi^a = g^{cd} [\Gamma_{cd}^a(g) - \Gamma_{cd}^a(\bar{g})], \quad (8)$$

where $\Gamma_{cd}^a(\mathbf{g})$ is the Christoffel connection associated to a metric $\mathbf{g}$, and $\bar{g}$ is a reference metric which we are free to choose. Solutions to the Einstein-DeTurck equation are generally not always Ricci flat. However, if we focus on static configurations that admit a regular Euclidean section, and have Kaluza-Klein asymptotics, it has been shown in [23] that solutions with $\xi \neq 0$ do not exist. As such, for such spacetimes, we expect solutions of the Einstein-DeTurck equation to coincide with Ricci-flat solutions. Solving (7) instead of $R_{ab} = 0$ offers significant advantages. Once a reference metric $\bar{g}$ is chosen, the Einstein-DeTurck equation forms a well-posed elliptic system [23]. Solutions must have $\xi^a = 0$, implying $\square x^a = g^{cd} \Gamma_{cd}^a(\bar{g})$. Thus, these solutions are naturally expressed in generalized harmonic coordinates [24], with the gauge source term fully controlled by $\bar{g}$. A good gauge choice involves choosing $\bar{g}$ judiciously. The choices of $\bar{g}$ across the solutions described in this Letter are presented in Supplemental Material [25].

Since the solutions we aim to construct first are static, they possess a hypersurface orthogonal Killing vector

field k . By introducing a time coordinate T that parametrizes the integral curves of k , we have $k = \partial/\partial T$. Additionally, these solutions are spherically symmetric with an $SO(9)$ isometry group. The most general line element compatible with such symmetries reads,

$$ds^2 = -Q_1(X, Z)dT^2 + \frac{4Q_2(X, Z)dX^2}{(1-X^2)^4} + L^2Q_4(X, Z)[dZ + Q_3(X, Z)dX]^2 + \frac{Q_5(X, Z)}{(1-X^2)^2}d\Omega_8^2, \quad (9)$$

where $Z \sim Z + 1$. The factors of $(1-X^2)$ were chosen so that near the asymptotic boundary, $r \sim [1/(1-X^2)]$, implying $X \rightarrow 1^-$ as $r \rightarrow +\infty$. Near $X = 1$, we demand $Q_1 = Q_4 = 1$, $Q_2 = Q_5 = K^2$, with $K > 0$ constant, and $Q_3 = 0$ so that the spacetime approaches

$$ds^2 = -dT^2 + dr^2 + r^2d\Omega_8^2 + L^2dZ^2 \quad (10)$$

at large r , showing that it exhibits the desired asymptotic structure. All the functions Q_i are periodic with respect Z , but as X becomes small these functions behave differently across the different black hole phases. In Fig. 1 we sketch the expected behavior for the different black hole topologies.

There are three phases of interest. In the uniform phase (rightmost panel of Fig. 1) translational invariant along the Z direction is preserved. In this phase the functions Q_i can be found analytically,

$$Q_1(X, Z) = X^2 \sum_{i=0}^6 (1-X^2)^i, \quad Q_2(X, Z) = \frac{X^2 r_0^2}{Q_1(X, Z)}, \\ Q_3(X, Z) = 0, \quad Q_4(X, Z) = 1 \quad \text{and} \quad Q_5(X, Z) = r_0^2, \quad (11)$$

with r_0/L parametrizing the thickness of the black string in terms of the length L of the Kaluza-Klein circle and with the black hole event horizon being located at $X = 0$. This solution is simply $\text{Schw}_{10D} \times S^1$, with Schw_{10D} being a ten-dimensional Schwarzschild black hole. Spatial cross

sections of the event horizon have topology $S^1 \times S^8$. Hereafter, phases with this topology will be referred to as black strings.

It has long been known [26] that for $r_0/L < 0.39858(3)$, the uniform phase becomes unstable to the so-called Gregory-Laflamme instability [27]. Furthermore, from the onset of the instability, a new family of *nonuniform* strings (middle panel of Fig. 1) emerges [28–34]. This phase still has an horizon whose spatial cross section has topology $S^1 \times S^8$, but no longer exhibits translations invariance along the Z direction. We have constructed this phase using the DeTurck trick, confirming, and extending, the results reported in [32].

The last phase of interest to this letter (leftmost panel in Fig. 1) corresponds to a solution where spatial cross sections of the event horizon no longer wrap the periodic direction, but instead have S^9 topology—these are often referred to as *localized* or *caged black holes*. Although this phase has been constructed in $d = 5, 6$ and $d = 10$ (see [20,35–42]; see also [43,44]), this has not been the case in $d = 11$ (the dimension of interest in this Letter). This phase is entirely novel and the details of its construction are given in Supplemental Material [25].

All of the static phases satisfy a first law of black hole mechanics as well as a Smarr type relation, namely,

$$d\mathcal{E}^{(0)} = \mathcal{T}_H^{(0)} dS_H^{(0)} \quad \text{and} \quad 8\mathcal{E}^{(0)} = 9\mathcal{T}_H^{(0)} S_H^{(0)} + \mathcal{T}_Z^{(0)}, \quad (12)$$

where $\mathcal{E}^{(0)}/G_{11}$, $S_H^{(0)}/G_{11}$, $\mathcal{T}_Z^{(0)}/G_{11}$ are the energy, entropy and tension densities, respectively, while $\mathcal{T}_H^{(0)}/L$ is the Hawking temperature.

BFSS thermodynamics—Having provided some details on the solutions that can exist with standard Kaluza-Klein asymptotics, we now describe how to map these to the D0-brane asymptotics of (4). This is achieved by applying a Carrollian transformation of the form [45]

$$T = \frac{L^{7/2}}{\mathcal{R}^{7/2}} \gamma t - \frac{1}{\gamma} \frac{\mathcal{R}^{7/2}}{L^{7/2}} z \quad \text{and} \quad Z = \frac{1}{\gamma} \frac{\mathcal{R}^{7/2}}{L^{7/2}} \frac{z}{L}, \quad (13a)$$

where γ is a constant to be adjusted in what follows so that the resulting metric has the correct asymptotic structure given in (4). It is a simple exercise to show that this transformation preserves regularity at the horizon, while modifying the asymptotic structure. This is crucial for the gravitational thermodynamics to directly correspond one to one with the thermodynamics of the BFSS dual theory. Using the general decaying properties of the several Q_i 's near spatial infinity, one finds,

$$\gamma = \frac{1}{\pi^{3/2}} \sqrt{\frac{15}{2}} \sqrt{\mathcal{E}^{(0)} - \mathcal{T}_Z^{(0)}}. \quad (13b)$$

As an example, the Carrollian transformation (13), takes the uniform black string (11) to (6), which we recall




FIG. 1. A schematic drawing of the transverse horizon radius for different black hole phases with standard Kaluza-Klein asymptotics: (a) the localized black hole phase, (b) the nonuniform string phase, and (c) the uniform string phase.

describes a finite temperature stack of D0-branes. Furthermore, since the static Kaluza-Klein circle has size L at spatial infinity, and we require the M-theory circle—parametrized by z —to have size $L_{11} = 2\pi g_s \ell_s$ there, the transformation (13) uniquely determines L to be

$$L = N^{\frac{1}{3}} (32\pi^8)^{1/9} g_s^{1/3} \ell_s \left(\mathcal{E}^{(0)} - T_Z^{(0)} \right)^{-1/9}. \quad (14)$$

Using [46,47], one can read off the energy ε , entropy σ , and free energy $\mathfrak{f}$ densities, as well as the temperature τ of all the BFSS field theory phases through the map (13),

$$\begin{aligned} \varepsilon &= N^{\frac{4}{3}} (32\pi^8)^{1/9} \frac{\mathcal{E}^{(0)} + T_Z^{(0)}}{\left(\mathcal{E}^{(0)} - T_Z^{(0)} \right)^{7/9}}, \\ \sigma &= N \frac{2\pi \mathcal{S}_H^{(0)}}{\mathcal{E}^{(0)} - T_Z^{(0)}}, \\ \mathfrak{f} &= N^{\frac{4}{3}} (32\pi^8)^{1/9} \frac{\mathcal{E}^{(0)} + T_Z^{(0)} - 2T_H^{(0)} \mathcal{S}_H^{(0)}}{\left(\mathcal{E}^{(0)} - T_Z^{(0)} \right)^{7/9}}, \\ \tau &= N^{-\frac{5}{3}} \left(\frac{32}{\pi} \right)^{1/9} T_H^{(0)} \left(\mathcal{E}^{(0)} - T_Z^{(0)} \right)^{2/9}, \end{aligned} \quad (15)$$

respectively. These expressions, which will be derived in [48], are central to the discussion in this Letter. In particular, given a static (non)uniform or localized gravitational solution, one can readily compute its thermodynamic properties $\{\mathcal{E}^{(0)}, T_Z^{(0)}, T_H^{(0)}, \mathcal{S}_H^{(0)}\}$ and, via (15), obtain the corresponding field theory thermodynamics $\{\varepsilon, \sigma, \mathfrak{f}, \tau\}$ of the (non)uniform or localized BFSS phases.

Results—We begin with the canonical ensemble, fixing the temperature $\tau N^{5/9}$ and identifying the phase with the lower free energy density, $\mathfrak{f} N^{-4/9}$. Figure 2 shows $\Delta \mathfrak{f}$, the free energy difference between a given BFSS thermal phase and the uniform phase at the same τ . The black disks show exact numerical data for the localized BFSS phase [49], and the red squares correspond to the nonuniform BFSS phase. We observe a first order phase transition between the localized phase and the uniform phase at $\tau N^{5/9} \approx 0.6118$. This is one of the main results of this Letter. This aligns with the phase diagram conjectured by [7], but here we provide a precise prediction for the critical temperature where the first-order phase transition occurs. In principle, it should be possible to derive the latter from a field theory calculation. We have also computed the specific heat associated with all the dominant phases in the canonical ensemble and found it to be always positive. At first glance, this might seem at odds with expectations for small localized black holes. However, the map given in (15) ensures that the specific heat remains positive even for the localized black hole phase.

On the other hand, the microcanonical ensemble reveals a much more interesting and unexpected structure.




FIG. 2. $\Delta \mathfrak{f}$, the free energy difference between a given thermal phase and the uniform phase at the same τ , as a function of τ . The black disks represent exact numerical data for the localized BFSS phase, while the red squares denote the nonuniform BFSS phase.

In Fig. 3, using the same color coding as Fig. 2, we plot $\Delta \sigma/N$, the entropy density difference between a given BFSS phase and the uniform phase at fixed energy density $\varepsilon N^{-4/9}$. For $0 < \varepsilon < \varepsilon_m$, the localized (black) phase dominates, while for $\varepsilon_m < \varepsilon < \varepsilon_M$, the nonuniform (red) phase dominates the microcanonical ensemble. A topology-changing transition occurs at $\varepsilon = \varepsilon_m$, as described by Kol in [51]. One finds that $1.505(4) \lesssim \varepsilon_m \lesssim 1.512(6)$, where the lower bound is found computing the maximum value of ε for which localized solutions exist, and the upper bound follows from finding the minimum value of ε for which nonuniform solutions exist. Improved numerics would pin down ε_m even further. On the other hand, we find the value of ε_M to be $\varepsilon_M \approx 1.713(2)$. This simply




FIG. 3. $\Delta \sigma$, the entropy density difference between a given thermodynamic phase and the uniform phase at the same energy ε , as a function of ε . The black disks represent exact numerical data for the localized BFSS phase, while the red squares denote the nonuniform BFSS phase.

follows from applying the map (15) to the gravitational Gregory-Laflamme instability onset of the uniform phase. It remains a formidable challenge to predict ϵ_m and ϵ_M from a field theory calculation in the microcanonical ensemble and we believe our precise identification of these values should motivate and guide such a search. The existence of an energy window where the non-uniform phase dominates the microcanonical ensemble is surprising and constitutes the main result of this Letter.

Discussion—We have studied the low energy limit of BFSS using its holographic dual. Our numerical results obtained in the canonical ensemble corroborate, and make precise, a longstanding conjecture set in [7]. Namely, that the dominant phase at small enough temperature is given by a localized thermal phase, dual to a black hole whose spatial horizon cross section has S^9 topology. The transition between the uniform BFSS phase and the localized BFSS phase occurs at $\tau N^{5/9} \approx 0.6118$. However, in the microcanonical ensemble we find a surprising result. For small enough energies, we find that the localized BFSS phase dominates, but we also find an energy *window* where a nonuniform phase with $S^8 \times S^1$ topology dominates, while breaking translational invariance along the M-theory circle.

One might wonder how our results can be made consistent with the dynamics of the original Gregory-Laflamme instability [27,52,53], which show an instability of the uniform black string at sufficiently low energies. To address this question it is, however, important to note that Carrollian transformations (13) do not preserve dynamics, despite mapping stationary solutions. Specifically, the linear analysis in [27] expands all metric functions in Fourier modes of the form $e^{-i\omega T + 2\pi i n Z}$. It should be noted that (13) maps these Fourier modes into modes in t and z that do *not* respect the required periodicity in z . Recall that for the original Gregory-Laflamme unstable modes, $\omega = i\Omega$ with $\Omega > 0$.

While there has been extensive research in high energy physics aimed at understanding the canonical ensemble of various field theories—using approaches such as localization; see for instance [54] and references therein—it is only recently that the microcanonical ensemble has garnered significant attention [55–57]. The increased complexity of studying the microcanonical ensemble stems from the necessity of understanding the detailed nature of field theory microstates. Addressing this challenge is notoriously difficult and is a key reason why a complete solution to the black hole information paradox remains elusive. Indeed, recent calculations show how to compute the entropy of Hawking radiation (see for instance [58–62]), but not how to compute its precise quantum state [63]. Our work reveals an intricate microcanonical ensemble structure for BFSS quantum mechanics at low energies. It would be desirable to identify all the relevant phases using BFSS degrees of freedom directly, though this task will

invariably require an understanding of the underlying microstates.

One possible interpretation for the absence of a first-order phase transition in the microcanonical ensemble, typically observed in phase transitions between localized black holes and uniform phases in $d \leq 13$ [20,35–42] with standard Kaluza-Klein asymptotics, is as follows. In standard thermodynamic systems, first-order phase transitions in the microcanonical ensemble can lead to mixed (also known as coexistent) phases. However, the boundary system in this context is quantum mechanical, which prevents the spatial separation of phases. Instead, we observe the presence of an intermediate phase (the non-uniform black string) that effectively interpolates between the localized and uniform phases [64].

The Carrollian map (13) together with the associated thermodynamical map (15) allows us to associate *any* black hole solution with standard Kaluza-Klein asymptotics into one with D0-brane asymptotics (and thus with any BFSS state). In particular, given the multitude of black hole solutions in higher dimensions—see, for instance, [65–69]—and the nontrivial nature of the map (15), one wonders how many more unexpected phases can possibly dominate a generalized ensemble that includes rotations and/or supersymmetric mass deformations *à la* BMN [70,71]. We leave these exploratory investigations to future work.

Acknowledgments—We thank Gary Horowitz, Juan Maldacena, and Yoav Zigdon for reading an earlier version of this Letter and for providing critical comments. O. D. acknowledges financial support from the STFC “Particle Physics Grants Panel (PPGP) 2018” Grant No. ST/T000775/1 and PPGP 2020 Grant No. ST/X000583/1. O. D.’s research was also supported in part by the YITP-ExU long-term workshop Quantum Information, Quantum Matter and Quantum Gravity (QIMG2023), Yukawa Institute for Theoretical Physics, Kyoto University, where part of this work was completed. J. E. S. has been partially supported by STFC consolidated Grants No. ST/T000694/1 and No. ST/X000664/1.

-
- [1] J. Maldacena, The large N limit of superconformal field theories and supergravity, *Adv. Theor. Math. Phys.* **2**, 231 (1998).
 - [2] E. Witten, Anti-de Sitter space and holography, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
 - [3] O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri, and Y. Oz, Large N field theories, string theory and gravity, *Phys. Rep.* **323**, 183 (2000).
 - [4] E. Y. Loh, J. E. Gubernatis, R. T. Scalettar, S. R. White, D. J. Scalapino, and R. L. Sugar, Sign problem in the numerical simulation of many-electron systems, *Phys. Rev. B* **41**, 9301 (1990).
 - [5] G. Bergner, N. Bodendorfer, M. Hanada, S. Pateloudis, E. Rinaldi, A. Schäfer, P. Vranas, and H. Watanabe (Monte Carlo String/M-theory Collaboration (MCSMC)),

- Confinement/deconfinement transition in the D0-brane matrix model: A signature of M-theory?, *J. High Energy Phys.* **05** (2022) 096.
- [6] S. Catterall, J. Giedt, and G.C. Toga, Holography from lattice $\mathcal{N} = 4$ super Yang-Mills, *J. High Energy Phys.* **08** (2023) 084.
- [7] N. Itzhaki, J.M. Maldacena, J. Sonnenschein, and S. Yankielowicz, Supergravity and the large N limit of theories with sixteen supercharges, *Phys. Rev. D* **58**, 046004 (1998).
- [8] T. Banks, W. Fischler, S.H. Shenker, and L. Susskind, M theory as a matrix model: A conjecture, *Phys. Rev. D* **55**, 5112 (1997).
- [9] M. Hanada, J. Nishimura, and S. Takeuchi, Non-lattice simulation for supersymmetric gauge theories in one dimension, *Phys. Rev. Lett.* **99**, 161602 (2007).
- [10] S. Catterall and T. Wiseman, Towards lattice simulation of the gauge theory duals to black holes and hot strings, *J. High Energy Phys.* **12** (2007) 104.
- [11] K.N. Anagnostopoulos, M. Hanada, J. Nishimura, and S. Takeuchi, Monte Carlo studies of supersymmetric matrix quantum mechanics with sixteen supercharges at finite temperature, *Phys. Rev. Lett.* **100**, 021601 (2008).
- [12] S. Catterall and T. Wiseman, Black hole thermodynamics from simulations of lattice Yang-Mills theory, *Phys. Rev. D* **78**, 041502(R) (2008).
- [13] M. Hanada, A. Miwa, J. Nishimura, and S. Takeuchi, Schwarzschild radius from Monte Carlo calculation of the Wilson loop in supersymmetric matrix quantum mechanics, *Phys. Rev. Lett.* **102**, 181602 (2009).
- [14] M. Hanada, Y. Hyakutake, J. Nishimura, and S. Takeuchi, Higher derivative corrections to black hole thermodynamics from supersymmetric matrix quantum mechanics, *Phys. Rev. Lett.* **102**, 191602 (2009).
- [15] S. Catterall and T. Wiseman, Extracting black hole physics from the lattice, *J. High Energy Phys.* **04** (2010) 077.
- [16] M. Hanada, J. Nishimura, Y. Sekino, and T. Yoneya, Monte Carlo studies of Matrix theory correlation functions, *Phys. Rev. Lett.* **104**, 151601 (2010).
- [17] M. Hanada, Y. Hyakutake, G. Ishiki, and J. Nishimura, Holographic description of quantum black hole on a computer, *Science* **344**, 882 (2014).
- [18] M. Hanada and J. Maltz, A proposal of the gauge theory description of the small Schwarzschild black hole in $\text{AdS}_5 \times S^5$, *J. High Energy Phys.* **02** (2017) 012.
- [19] M. Hanada and H. Watanabe, On thermal transition in QCD, *Prog. Theor. Exp. Phys.* **2024**, 043B02 (2024).
- [20] M. Headrick, S. Kitchen, and T. Wiseman, A new approach to static numerical relativity, and its application to Kaluza-Klein black holes, *Classical Quantum Gravity* **27**, 035002 (2010).
- [21] O. J. C. Dias, J. E. Santos, and B. Way, Numerical methods for finding stationary gravitational solutions, *Classical Quantum Gravity* **33**, 133001 (2016).
- [22] T. Wiseman, Numerical construction of static and stationary black holes, in *Black Holes in Higher Dimensions*, edited by G. T. Horowitz (Cambridge University Press, Cambridge, England, 2012), pp. 233–270.
- [23] P. Figueras, J. Lucietti, and T. Wiseman, Ricci solitons, Ricci flow, and strongly coupled CFT in the Schwarzschild Unruh or Boulware vacua, *Classical Quantum Gravity* **28**, 215018 (2011).
- [24] H. Friedrich, On the hyperbolicity of Einstein's and other gauge field equations, *Commun. Math. Phys.* **100**, 525 (1985).
- [25] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/wws8-m34w> for a self-contained detailed description of the method employed to find the two families of solutions of the main text and how to compute the thermodynamical observables of the systems.
- [26] E. Sorkin, A critical dimension in the black string phase transition, *Phys. Rev. Lett.* **93**, 031601 (2004).
- [27] R. Gregory and R. Laflamme, Black strings and p-branes are unstable, *Phys. Rev. Lett.* **70**, 2837 (1993).
- [28] G.T. Horowitz and K. Maeda, Fate of the black string instability, *Phys. Rev. Lett.* **87**, 131301 (2001).
- [29] S. S. Gubser, On nonuniform black branes, *Classical Quantum Gravity* **19**, 4825 (2002).
- [30] T. Wiseman, Static axisymmetric vacuum solutions and nonuniform black strings, *Classical Quantum Gravity* **20**, 1137 (2003).
- [31] E. Sorkin, Non-uniform black strings in various dimensions, *Phys. Rev. D* **74**, 104027 (2006).
- [32] P. Figueras, K. Murata, and H. S. Reall, Stable non-uniform black strings below the critical dimension, *J. High Energy Phys.* **11** (2012) 071.
- [33] M. Kalisch and M. Ansorg, Highly deformed non-uniform black strings in six dimensions, in *Proceedings of the 14th Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Astrophysics, and Relativistic Field Theories* (2017), Vol. 2, pp. 1799–1804, [10.1142/9789813226609_0185](https://arxiv.org/abs/10.1142/9789813226609_0185).
- [34] M. Kalisch and M. Ansorg, Pseudo-spectral construction of non-uniform black string solutions in five and six spacetime dimensions, *Classical Quantum Gravity* **33**, 215005 (2016).
- [35] T. Wiseman, From black strings to black holes, *Classical Quantum Gravity* **20**, 1177 (2003).
- [36] E. Sorkin, B. Kol, and T. Piran, Caged black holes: Black holes in compactified space-times. 2. 5-d numerical implementation, *Phys. Rev. D* **69**, 064032 (2004).
- [37] H. Kudoh and T. Wiseman, Properties of Kaluza-Klein black holes, *Prog. Theor. Phys.* **111**, 475 (2004).
- [38] H. Kudoh and T. Wiseman, Connecting black holes and black strings, *Phys. Rev. Lett.* **94**, 161102 (2005).
- [39] M. Kalisch, S. Möckel, and M. Ammon, Critical behavior of the black hole/black string transition, *J. High Energy Phys.* **08** (2017) 049.
- [40] O. J. C. Dias, J. E. Santos, and B. Way, Localised and nonuniform thermal states of super-Yang-Mills on a circle, *J. High Energy Phys.* **06** (2017) 029.
- [41] M. Ammon, M. Kalisch, and S. Moeckel, Notes on ten-dimensional localized black holes and deconfined states in two-dimensional SYM, *J. High Energy Phys.* **11** (2018) 090.
- [42] B. Cardona and P. Figueras, Critical Kaluza-Klein black holes and black strings in $D = 10$, *J. High Energy Phys.* **11** (2018) 120.
- [43] O. J. C. Dias, J. E. Santos, and B. Way, Lumpy $\text{AdS}_5 \times S^5$ black holes and black belts, *J. High Energy Phys.* **04** (2015) 060.

- [44] O. J. C. Dias, J. E. Santos, and B. Way, Localised $\text{AdS}_5 \times S^5$ Black holes, *Phys. Rev. Lett.* **117**, 151101 (2016).
- [45] This transformation can also be interpreted as a lightlike compactification of the form $(T, Z) \rightarrow (T, Z) + (-R, +R)$, followed by a coordinate transformation and a particular choice of R . We are grateful to Juan Maldacena for bringing this terminology to our attention.
- [46] R. M. Wald and A. Zoupas, A general definition of ‘conserved quantities’ in general relativity and other theories of gravity, *Phys. Rev. D* **61**, 084027 (2000).
- [47] O. J. C. Dias, G. S. Hartnett, and J. E. Santos, Covariant Noether charges for type IIB and 11-dimensional supergravities, *Classical Quantum Gravity* **38**, 015003 (2021).
- [48] O. J. C. Dias and J. E. Santos, Localized states of super quantum mechanics (to be published).
- [49] In a forthcoming publication, we compare our numerical localized solutions with the second-order perturbative results from [50] and find excellent agreement in regions of moduli space where localized black holes do exist.
- [50] T. Harmark, Small black holes on cylinders, *Phys. Rev. D* **69**, 104015 (2004).
- [51] B. Kol, Topology change in general relativity, and the black hole black string transition, *J. High Energy Phys.* **10** (2005) 049.
- [52] L. Lehner and F. Pretorius, Black strings, low viscosity fluids, and violation of cosmic censorship, *Phys. Rev. Lett.* **105**, 101102 (2010).
- [53] P. Figueras, T. França, C. Gu, and T. Andrade, Endpoint of the Gregory-Laflamme instability of black strings revisited, *Phys. Rev. D* **107**, 044028 (2023).
- [54] N. Bobev, J. Hong, and V. Reys, Holographic thermal observables and M2-branes, *J. High Energy Phys.* **12** (2023) 054.
- [55] O. J. C. Dias, P. Mitra, and J. E. Santos, New phases of $\mathcal{N} = 4$ SYM at finite chemical potential, *J. High Energy Phys.* **05** (2023) 053.
- [56] D. Marolf and J. E. Santos, Stability of the microcanonical ensemble in Euclidean quantum gravity, *J. High Energy Phys.* **11** (2022) 046.
- [57] S. Kim, S. Kundu, E. Lee, J. Lee, S. Minwalla, and C. Patel, Grey Galaxies’ as an endpoint of the Kerr-AdS superradiant instability, *J. High Energy Phys.* **11** (2023) 024.
- [58] G. Penington, Entanglement wedge reconstruction and the information paradox, *J. High Energy Phys.* **09** (2020) 002.
- [59] A. Almheiri, N. Engelhardt, D. Marolf, and H. Maxfield, The entropy of bulk quantum fields and the entanglement wedge of an evaporating black hole, *J. High Energy Phys.* **12** (2019) 063.
- [60] A. Almheiri, R. Mahajan, and J. Maldacena, Islands outside the horizon, [arXiv:1910.11077](https://arxiv.org/abs/1910.11077).
- [61] A. Almheiri, R. Mahajan, J. Maldacena, and Y. Zhao, The page curve of Hawking radiation from semiclassical geometry, *J. High Energy Phys.* **03** (2020) 149.
- [62] A. Almheiri, R. Mahajan, and J. E. Santos, Entanglement islands in higher dimensions, *SciPost Phys.* **9**, 001 (2020).
- [63] A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian, and A. Tajdini, The entropy of Hawking radiation, *Rev. Mod. Phys.* **93**, 035002 (2021).
- [64] We would like to thank Juan Maldacena for suggesting this interpretation.
- [65] R. Emparan and H. S. Reall, Black holes in higher dimensions, *Living Rev. Relativity* **11**, 6 (2008).
- [66] O. J. C. Dias, P. Figueras, R. Monteiro, H. S. Reall, and J. E. Santos, An instability of higher-dimensional rotating black holes, *J. High Energy Phys.* **05** (2010) 076.
- [67] O. J. C. Dias, T. Ishii, K. Murata, J. E. Santos, and B. Way, Gregory-Laflamme encounters superradiance, *J. High Energy Phys.* **01** (2023) 147.
- [68] O. J. C. Dias, T. Ishii, K. Murata, J. E. Santos, and B. Way, Gregory-Laflamme and superradiance encounter black resonator strings, *J. High Energy Phys.* **02** (2023) 069.
- [69] O. J. C. Dias, T. Ishii, K. Murata, J. E. Santos, and B. Way, Superradiance and black resonator strings encounter helical black strings, *J. High Energy Phys.* **05** (2023) 041.
- [70] D. E. Berenstein, J. M. Maldacena, and H. S. Nastase, Strings in flat space and pp waves from $N = 4$ super-Yang-Mills, *J. High Energy Phys.* **04** (2002) 013.
- [71] M. S. Costa, L. Greenspan, J. Penedones, and J. Santos, Thermodynamics of the BMN matrix model at strong coupling, *J. High Energy Phys.* **03** (2015) 069.