

Erratum: Modeling the high-pressure solid and liquid phases of tin from deep potentials with *ab initio* accuracy [Phys. Rev. Materials 7, 053603 (2023)]

Tao Chen , Fengbo Yuan, Jianchuan Liu, Huayun Geng, Linfeng Zhang, Han Wang, and Mohan Chen 



(Received 12 October 2025; published 28 October 2025)

DOI: [10.1103/cnm2-fcqv](https://doi.org/10.1103/cnm2-fcqv)

In Fig. 7(a) of our original paper, the region that should have been labelled as β phase was mistakenly marked as bct phase, and vice versa. Here is the revised figure.

The discussion and conclusions presented in our original paper remain unaffected.




FIG. 7. Phase diagram of Sn with temperatures ranging from 0 to 2000 K and pressures ranging from 0 to 20 GPa from (a) experiments [12] and (b) the DP-SCAN model. Four phases of Sn are presented including the α -Sn, β -Sn, bct-Sn, and liquid phases. The coexistence points on the phase diagram of (b) are labeled as A (5 GPa, 741 K), B (20 GPa, 531 K), C (20 GPa, 1387 K), and D (10 GPa, 853.5 K). In addition, (c) shows the phase diagram with temperatures ranging from 0 to 1000 K and pressures ranging from 0 to 3 GPa from the DP-PBE model.