

Direct observation of saturated heat-flux inhibition by magnetic fields in laser-produced plasmas

A. Triantafyllidis^{1,*}, J.-R. Marquès¹, P. Loiseau^{2,3}, C. Vlachos⁴, J. J. Santos⁴, J. Béard⁵, J.-M. Lagarrigue⁵,
L. Lancia¹, N. Ozaki^{6,7}, M. Koenig¹ and B. Albertazzi^{1,‡}

¹LULI—CNRS, CEA, Sorbonne Universités, École Polytechnique, Institut Polytechnique de Paris, F-91120 Palaiseau cedex, France

²CEA, DAM, DIF, F-91297 Arpaçon, France

³Université Paris-Saclay, CEA, Laboratoire Matière en Conditions Extrêmes, F-91680 Bruyères-le-Châtel, France

⁴Université de Bordeaux, CNRS, CEA, Centre Lasers Intenses et Applications (CELIA), F-33405 Talence, France

⁵Laboratoire National des Champs Magnétiques Intenses (LNCMI), UPR 3228, CNRS-Université Joseph Fourier (UJF)-Université Paul Sabatier (UPS)-Institut National des Sciences Appliquées (INSA), F-31400 Toulouse, France

⁶Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

⁷Institute of Laser Engineering, Osaka University, Suita, Osaka 565-0871, Japan



(Received 16 March 2026; accepted 2 September 2026; published 5 October 2026)

We report direct measurements of magnetically inhibited heat transport in an underdense laser-produced plasma subjected to externally applied magnetic fields of up to 20 T. Spatially and temporally resolved Thomson-scattering data show that the peak electron temperature increases by a factor ~ 1.4 relative to the unmagnetized case, as cross-field thermal conduction is maximally suppressed. This state is observed for $B \geq 10$ T and is sustained over ~ 0.5 ns. The subsequent temperature decrease is primarily driven by a reduction in inverse-bremsstrahlung heating as the electron density cavitates. Our experimental results are consistent with simple analytical estimates and are corroborated by Vlasov-Fokker-Planck simulations, suggesting that the plasma reaches the maximum temperature gain for an electron Hall parameter $\chi_{\text{Hall}} \gtrsim 50$ ($T_e \sim 800$ eV, $n_e \sim 3 \times 10^{19}$ cm $^{-3}$). These findings demonstrate that effective magnetothermal insulation can occur in magnetized inertial confinement fusion schemes without the need for extremely large magnetic fields.

DOI: [10.1103/2rqf-hq77](https://doi.org/10.1103/2rqf-hq77)

I. INTRODUCTION

Heat transport in magnetized, collisional plasmas is a fundamental process in high-energy-density physics. As magnetic fields force electrons to gyrate around the field lines, cross-field heat flux can be strongly inhibited, inducing an increase in plasma temperature and introducing anisotropy that inherently modifies the system behavior, hydrodynamics, and energy dissipation [1–6]. At the same time, magnetic fields can be advected by the Nernst effect and suprathermal electrons [7–10], giving rise to structures such as magnetic transport barriers and magnetic cavities [2,3,11–14]. However, despite the development of multiple theoretical frameworks [15–25], the relationship between heat flux and magnetic fields remains poorly understood, largely due to a critical lack of experimental benchmarks.

Prior to this work, direct measurements of electron heat transport under imposed magnetic fields were scarce. In a notable experiment by Froula *et al.* [1], the authors observed the

quenching of heat flow in an N₂ gas-jet plasma with an applied 12 T external field, reporting a fourfold increase in electron temperature. However, kinetic simulations of this experiment were unable to fully reproduce their results [2], instead predicting a maximum temperature enhancement by only a factor of 2. More recently, Arran *et al.* [3] directly measured the influence of the Nernst effect in a similar laser-heated, magnetized plasma experiment using proton radiography, showing that it can dominate the field evolution at early times and remain important up to ~ 1 ns.

These interactions are of particular interest in the framework of inertial confinement fusion (ICF). Over the last few years, the idea of premagnetizing the imploding fuel has drawn significant interest, as both simulations [26–32] and experiments [33–38] have shown that a seed field can (1) improve the fusion yield and laser-target coupling, (2) modify perturbations, (3) reduce fuel-ablator mixing, and (4) better confine α particles. In particular, Sio *et al.* [37] reported hot-spot temperature and neutron-yield gains in magnetized indirect-drive implosions, with a sizable performance increase already observed at ~ 12 T and a smaller additional gain at ~ 26 T. Furthermore, in indirect drive, the Nernst effect can pile up the magnetic field along the hohlraum walls, where the accumulated field can strongly inhibit nonlocal transport and preheat [13,39].

In this paper, we report direct experimental evidence that magnetothermal insulation saturates in an underdense H₂ plasma beyond a threshold Hall parameter value $\chi_{\text{Hall}} = \omega_{ce} \tau_e \gtrsim 50$, where ω_{ce} is the electron gyrofrequency

*Present address: Graduate School of Engineering, Osaka University, Osaka, Japan.

†Contact author: angelos@eie.eng.osaka-u.ac.jp

‡Contact author: bruno.albertazzi@polytechnique.edu

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.




FIG. 1. A laser beam irradiated a gas jet, generating plasma. Part of the probe beam's photons were scattered at 90° (along the z axis) and collected by a spectrometer that resolved the electronic plasma wave feature in time and space. The magnetic field is oriented along the x axis.

and τ_e is the electron collision time. Using a spatially and temporally resolved Thomson-scattering (TS) diagnostic, we measure a plateau in temperature gain: A 10 T field induces a $1.4\times$ increase in electron temperature relative to 0 T, while doubling the field strength produces only a marginal rise. Unlike the Froula *et al.* [1] experiment, we show a more modest temperature increase, even though the heat conduction inhibition is orders of magnitude stronger in our experiment. Moreover, we follow this behavior in time and observe a density cavitation driven by strong thermal-pressure gradients. Finally, Vlasov-Fokker-Planck (VFP) simulations support our results and indicate that the temperature gain does not increase further even for arbitrarily large magnetic fields, potentially relaxing design constraints for magnetized ICF [33,35,36].

II. EXPERIMENTAL SETUP AND METHODOLOGY

The experimental setup is shown in Fig. 1. The experiment was performed at the LULI2000 laser facility (École Polytechnique, France), where a nanosecond laser beam ($\tau = 1.5$ ns, 200 ps of rise time, square profile) with energy $E_0 \approx 800$ J at the fundamental wavelength (1053 nm) irradiated a hypersonic cylindrical gas jet, generating plasma. The beam was defocused to a 500 μm diameter at the interaction plane, resulting in an average intensity of $I_0 = 2.5 \times 10^{14}$ W/cm 2 . The gas-jet nozzle was positioned 1.5 mm below the interaction region, where it had a Gaussian density profile with an FWHM of 950 μm (peak electron density of 4×10^{19} cm $^{-3}$ for H $_2$ at 14 bar). An external magnetic field up to 20 T, oriented along the laser direction (x axis), was generated by a Helmholtz coil system [40]; the field was spatially uniform to within a few percent over the >1 cm 3 interaction volume and varied by less than 2% over the ~ 2 μs interval around its peak. Its initial strength was measured with a Rogowski probe and used throughout the experimental analysis to calculate the dimensionless numbers reported in this work. Direct




FIG. 2. (Top) Spatially resolved electronic Thomson-scattering images at 1 ns for (a) 0 T and (b) 20 T. The heater beam irradiates the $-250 \mu\text{m} < y < 250 \mu\text{m}$ region and the dashed arrow shows the direction of the probe beam. (Bottom) Experimental data integrated over the shaded regions and overlaid with synthetic Thomson-scattering spectra: (c) 2.5×10^{19} cm $^{-3}$, 450 eV and (d) 2.3×10^{19} cm $^{-3}$, 760 eV.

measurements of the in-plasma field, such as those obtained using proton deflectometry [41–43], could not be performed because of geometric constraints imposed by the electromagnetic coil casing. The difference between nominal and instantaneous field values is addressed in the VFP simulations section (Sec. IV A).

A frequency-doubled probe beam (526.5 nm, $\lesssim 3$ J, $\tau = 1.5$ ns, 30 μm focal spot size) propagated along the y axis and interacted with the laser-produced plasma (see Appendix B for the probe's impact). An $f/6$ lens system collected the photons that scattered off at a 90° angle (z axis), imaging the x – y plane at the entrance of a spectrometer. There, they were dispersed (spectral resolution of 2.4 nm) and recorded by a gated optical imager and a streak camera, providing both spatially and temporally resolved electron Thomson-scattering spectra. Typical spatially resolved spectra, integrated over 125 ps, are shown in Figs. 2(a) and 2(b): The wavelength dispersion is along x , while the transverse spatial coordinate is mapped along y , producing a one-dimensional spatial slice with resolution of 40 μm . The heater-beam axis corresponds to $y = 0$ and the wavelengths around 526.5 nm were blocked by a notch filter.

Local plasma conditions were extracted by fitting the measured Thomson-scattering spectra with synthetic ones via a χ^2 -minimization procedure [see Figs. 2(c) and 2(d)] [44,45], assuming a Maxwellian electron velocity distribution function.




FIG. 3. Interferometry images for 0, 10, and 20 T at (top) 1 ns and (bottom) 2 ns with $P_B = 3.5$ bar (gas-jet backing pressure). The red arrow indicates the propagation direction of the laser. For 0 T, a structure that extends past the shock front can be seen (fringe shifts ahead of the shock front), attributed to preheating due to suprathermal electrons.

III. RESULTS

A. Inhibition of magnetized transport and saturation of the temperature gain

At 20 T, the Thomson satellites are noticeably broader than at 0 T, indicating a substantially higher electron temperature (T_e). Microscopically, this behavior is attributed to reduced electron mobility. Without any magnetization, electrons that transport most of the heat can stream across the temperature gradient with minimal collisions and deposit their energy ahead of it (preheat due to nonlocal transport) [9,18,21,46–48]: For 760 eV, electrons moving at the thermal velocity have a mean free path (mfp) of $\lambda_{\text{mfp}} = 103.6 \mu\text{m}$. In contrast, the magnetic field forces them to gyrate, effectively relocating transport by reducing the mfp to the Larmor radius ($\rho_L = 3.3 \mu\text{m}$ for $B = 20$ T).

This is seen in Fig. 3 by comparing the 0, 10, and 20 T images, showing a more spatially confined plasma morphology in the presence of the external field. In the unmagnetized case, fringe shifts extend ahead of the shock front, indicating strong preheating ahead of the laser-heated region, but even a 10 T field is sufficient to strongly suppress this behavior. As a result, the unmagnetized plasma experiences larger energy losses, leading to lower peak T_e values and less steep T_e profiles. We should note that the magnetic field is not dynamically important in this parameter range, as quantified by two dimensionless parameters. The ram-pressure ratio $\beta_{\text{ram}} = \rho v^2 / (B^2 / 2\mu_0) \sim 5\text{--}10$ (with $v \sim 100\text{--}200$ km/s at 1–2 ns for $B = 20$ T) shows that the field cannot redirect the bulk flow,

while the thermal-pressure ratio $\beta_{\text{th}} = n_e k_B T_e / (B^2 / 2\mu_0) \gtrsim 20$ shows that magnetic pressure is negligible in the energy balance; since both exceed unity, the plasma is dominated by the thermal pressure and the field is dynamically weak. The full parameter range and the diagnostics used to infer the dimensionless numbers are summarized in Appendix A.

Figure 4 shows the electron density (n_e) and T_e spatial profiles for $B = 0\text{--}20$ T, produced by sampling the Thomson-scattering spectra every $95 \mu\text{m}$. Starting with the top panels ($t = 1$ ns), the n_e curves are similar in all cases: Shock fronts develop at $y \approx \pm 400 \mu\text{m}$ as a direct result of laser heating, while the central feature is due to the gas-jet density profile. However, a clear temperature increase is observed when the plasma is subjected to an external magnetic field. For $y = 0$, the peak T_e rises from 580 eV in the unmagnetized case to 800 eV for $B = 20$ T, while at $y = -300 \mu\text{m}$, T_e shows the maximum enhancement (from 309 to 781 eV), both consistent with the suppression of cross-field losses.

Most notably, the 10 and 20 T temperature profiles are nearly indistinguishable at 1 ns [Fig. 4(b)] because, once $\chi_{\text{Hall}} \propto T_e^{3/2} B / n_e$ reaches ~ 50 , stronger B values only lead to a marginal rise in T_e . In terms of heat flux inhibition, the classical Braginskii model [16] predicts $\kappa_{\perp} / \kappa_{\parallel} \approx \chi_{\text{Hall}}^{-2} \sim 5 \times 10^{-4}$ for 10 T and $\sim 10^{-4}$ for 20 T at $y = 0$ (we neglect the $\kappa_{\wedge}$ term due to the system's cylindrical geometry), where $\kappa_{\parallel, \perp}$ are the thermal conductivity coefficients parallel and perpendicular to the field in the electron heat flux $\mathbf{q}_e$ expression:

$$\mathbf{q}_e = -\kappa_{\parallel} \nabla_{\parallel} T_e - \kappa_{\perp} \nabla_{\perp} T_e. \quad (1)$$




FIG. 4. Spatial n_e and T_e profiles of H_2 gas-jet plasma ($P_B = 14$ bar) for $B = 0$ –20 T at (top) 1 ns and (bottom) 2 ns, retrieved using a χ^2 -minimization fitting. The shaded uncertainty bands are displayed at half their actual width for visual clarity.

In other words, the magnetic field reduces cross-field heat losses, trapping energy that would otherwise escape. However, it cannot eliminate heat flow parallel to $\mathbf{B}$ so, in the strongly magnetized limit ($\chi_{\text{Hall}} \gg 1$), the system tends to a peak T_e asymptotically. We refer to this state of maximum heat-flux inhibition as *saturation*. In contrast, the 5 T curve in Fig. 4 ($T_{e,\text{max}} \sim 650$ eV, $\chi_{\text{Hall}} \sim 25$) lies below the saturation threshold: Cross-field transport is only partially inhibited, so additional magnetization still measurably raises the peak T_e .

B. Analytical estimates of the saturated gain

A simple way to estimate the T_e gain during saturation is the geometric argument used by Montgomery *et al.* [34], recognizing that conductive losses scale with the total surface area A through which heat can dissipate. For a cylindrical plasma morphology, heat is conducted through both the cylindrical side wall ($A_{\perp} = 2\pi RL$) and the bases ($A_{\parallel} = 2\pi R^2$) when $B = 0$ T, whereas in the strong magnetization limit it can only do so through $A_{\parallel}$. The laser-produced plasma has a radius $R \approx 0.5$ mm and length $L \approx 2$ mm at 1 ns (see Fig. 3), so $(A_{\parallel} + A_{\perp})/A_{\parallel} = 1 + L/R = 5$. Finally, the unmagnetized conductive heat losses scale as $AT_e^{7/2}$ [15], giving an upper bound estimate for the temperature increase of $5^{2/7} \approx 1.58$.

For plasmas that are directly irradiated by a laser, this estimate can be further refined by accounting for the temperature dependence of inverse bremsstrahlung (IB). In this laser intensity range ($I_0 \sim 10^{14}$ W/cm²), IB is the dominant heating mechanism [49,50], but its heating rate decreases sharply with increasing temperature. Assuming a fully ionized plasma,

$$\left(\frac{\partial T_e}{\partial t}\right)_{\text{IB}} \propto \frac{Z^* n_e \ln \Lambda}{T_e^{3/2}} v_{\text{osc}}^2, \quad (2)$$

where Z^* is the ionization state, $\ln \Lambda$ is the Coulomb logarithm, $v_{\text{osc}} = eE_L/(m_e \omega_L)$ is the electron oscillation velocity, $E_L = \sqrt{2I_0/(\epsilon_0 c)}$ is the laser electric field amplitude, e and m_e are the electron charge and mass, and ω_L is the laser angular frequency [51–53]. Therefore, for a constant n_e , the plasma becomes more transparent to the laser (lower absorption) as T_e grows, so the heating rate drops. The peak temperature is thus set when conductive losses are balanced by IB heating:

$$\frac{\partial(\rho_e \epsilon_e)}{\partial t} = P_{\text{IB}} - P_{q_e} = 0, \quad (3)$$

where ρ_e is the electron mass density, ϵ_e is the electron energy, P_{IB} is the IB heating power, and P_{q_e} denotes conductive losses ($\nabla \cdot \mathbf{q}_e$ in the unmagnetized case and $\nabla_{\parallel} \cdot \mathbf{q}_{e,\parallel}$ in the strong-magnetization limit). In the quasisteady state ($P_{\text{IB}} = P_{q_e}$) and for the same laser parameters, one obtains $T_e^5 \propto A^{-1}$, so the temperature gain scales as $(1 + L/R)^{1/5}$, giving ~ 1.38 for our system.

Consistent with this upper bound, we experimentally observe a T_e increase of $\sim 1.4\times$ in the central region relative to the unmagnetized case. For comparison, applying the same estimate to the Froula *et al.* experiment [1] yields a maximum temperature increase of about 1.8 – $2\times$, which is in good agreement with the simulations by Ridgers *et al.* [2], but well below the $4\times$ increase reported in that experiment. Part of this discrepancy may stem from an enhanced nonlocal behavior: The heater beam had a smaller focal spot diameter (150 μm versus 500 μm here) and used a shorter laser pulse (1 ns versus 1.5 ns here). These conditions should produce steeper gradients and stronger nonlocal transport in the unmagnetized plasma, leading to larger energy losses and a lower unmagnetized T_e , thereby increasing the observed T_e gain (see Appendix C for a more detailed comparison).

C. Late-time density cavitation

The energy balance of Eq. (3) depends on IB heating and heat-flux losses that evolve along with the plasma conditions. At 2 ns [Figs. 4(c) and 4(d)], the following features can be observed for $B \geq 10$ T: Two steep, dense fronts emerge on either side of the laser-irradiated region ($y \approx \pm 700$ μm) and a density cavity is formed at $y = 0$. The fronts are more than $1.5\times$ denser and the cavity more rarefied than at the same positions in the unmagnetized case. Moreover, the T_e profiles are similar for all B values, showing that heat has already dissipated over this time frame.

Figure 5 presents temporally resolved Thomson-scattering data at the target chamber center (TCC), i.e., $x = y = z = 0$, showing that the quasisteady state is sustained for ~ 0.5 ns (60 ps temporal resolution and integrated over a $30 \times 30 \times 50$ μm^3 volume). Initially, n_e evolves similarly for 0 and 20 T, but from 1.2 to 1.5 ns, a steep drop occurs in the magnetized case. Afterward, the unmagnetized plasma is consistently denser for the remainder of the probing window. As for T_e , it exhibits the familiar modification in the presence of a 20 T field; the peak T_e is reached at ~ 0.7 ns. Shortly after, the plasma begins to cool down even before the laser irradiation ends. This cooling coincides with the steep n_e drop in the 20 T case because IB absorption depends strongly on the electron density [see Eq. (2)]. The onset of density cavitation causes a reduction of




FIG. 5. Temporal evolution of (left) n_e and (right) T_e for H_2 at the TCC for 0 and 20 T. The curves are reconstructed from two separate shots (0–1.5 and 1.5–2.9 ns). The intensity temporal profiles of the heater and probe beams are overlaid on the right panel (arbitrary units; the dashed line denotes the probe profile for a delay of 1.5 ns). Beam power is not drawn to scale.

laser energy deposition in the irradiated region so, once the heating rate collapses, the heat losses dominate and T_e drops.

The density cavitation itself results from heat being trapped inside the laser-irradiated region. As cross-field heat flux is effectively stopped for 10 and 20 T, steep temperature gradients develop that drive strong pressure gradients. Plasma is therefore advected laterally away from the heated region, piling up at $y = \pm 700 \mu\text{m}$ and depleting the center, which produces the observed dense fronts and density cavity. Early time Nernst advection could also contribute indirectly by advecting the magnetic field, forming transport barriers and a magnetic cavity, as described in Refs. [2,3]. With $Re_m > 900$ and $\beta_{th} \gtrsim 20$, where Re_m is the magnetic Reynolds number and β_{th} is the ratio of the plasma's thermal to magnetic pressure, resistive diffusion and magnetic-pressure-driven terms are negligible (see the magnetic field induction equation in Refs. [54,55]). Thus, both ideal magnetohydrodynamics (MHD) advection and the Nernst effect can in principle contribute to the field evolution [3,7,12,13]. However, their contributions here cannot be separated quantitatively without directly probing the magnetic field, for example, by proton deflectometry or Zeeman splitting, which could be performed once the experimental geometry permits [42,56–58].

IV. SIMULATIONS AND DISCUSSION

A. Vlasov-Fokker-Planck simulations

To support our conclusions, we carried out simulations with the kinetic code ALADIN [59–61]. ALADIN is an implicit code that solves the VFP equation in the diffusion approximation, coupled to Ampère's and Faraday's laws, in two spatial dimensions (y and z) and one velocity dimension ($|v|$) [52]. The code advances the electromagnetic field components E_y , E_z , and B_x , and is coupled to a hydrodynamic module, enabling kinetic simulations over timescales relevant to the bulk plasma motion. In the simulation, a laser pulse with parameters matching the experimental conditions propagated through the y - z plane (along the x axis, not simulated) and irradiated an H_2 gas jet identical to the measured profile but with an initial $T_e = 20$ eV. For 40 T, numerical instabilities prevented us from running the simulation up to 2 ns.

ALADIN reproduces the system dynamics well at 1 ns, accurately capturing both the density profiles and the saturation of heat-flux suppression [see Figs. 6(a) and 6(b)]. Once saturation sets in, T_e plateaus at 836 eV for 20 T and a 40 T field only raises the temperature by 6 eV. Figure 6(e) plots the simulated T_e as a function of the calculated Hall parameter χ_{Hall} with

their experimental counterparts and shows a clear asymptotic behavior. The simulated χ_{Hall} values are systematically below the measurements partly because they are computed using the instantaneous magnetic field from ALADIN, whereas the experimental points use the nominal applied field. ALADIN predicts that the magnetic field cavitates in all magnetized runs as the system evolves, mainly via advection, but the relative reduction is largest for $B = 5$ T: Over 1 ns, the field at $y = 0$ decreases by 25% to 3.8 T, whereas for 20 T it drops only to 19.8 T. Thus, the use of the nominal field in the experimental data does not change our conclusions, but only slightly affects the precise value of χ_{Hall} .

Even though the simulations successfully predict many experimentally observed features, their two-dimensional geometry inherently prevents them from capturing the plasma cooling. In other words, since heat can only dissipate through




FIG. 6. (a)–(d) Simulated spatial n_e and T_e profiles for $t = 1, 2$ ns. (e) Comparison between the experimental and simulated values of T_e and χ_{Hall} during saturation ($t = 1$ ns) at $y = 0$. Experimental χ_{Hall} values assume the nominal field strength, whereas simulated ones use the B obtained from ALADIN.

the $A_{\perp}$ cylinder surface, T_e does not drop rapidly enough, leading to significantly higher T_e values at $t = 2$ ns compared to the experiment [see Figs. 6(c) and 6(d)]. The only exception is the unmagnetized case, where the simulated and experimental T_e are in closer agreement, consistent with our proposed explanation. For this reason, modeling the late-time cooling likely requires a three-dimensional treatment to account for the losses along B .

B. Implications for magnetized ICF

Regarding magnetized ICF, our results imply that the temperature gain from magnetothermal insulation *during the compression phase* is intrinsically limited. Once the Hall parameter exceeds a critical threshold ($\chi_{\text{Hall}} \sim 50$), cross-field conduction is effectively quenched and T_e can be enhanced by up to $\sim(1 + L/R)^{2/7}$ times, assuming a cylindrical geometry. For typical indirect-drive ICF parameters (3 keV, $2.5 \times 10^{24} \text{ cm}^{-3}$, and $2000\times$ target compression) [62], a seed magnetic field of $B_0 \approx 26$ T is required for $\chi_{\text{Hall}} = 50$ (assuming ideal-MHD compression of B and neglecting Nernst advection). Attaining this would deliver close to the largest feasible T_e increase, improving the prospects for ignition. Therefore, for magnetothermal insulation, a practical target design should aim at this value rather than at an arbitrarily large field, although stronger fields may still benefit the implosion in other ways; for example, by suppressing the growth of small-scale Rayleigh-Taylor perturbations [28], or by raising the ignition margin and thereby improving robustness to low-mode asymmetries [27].

This conclusion is consistent with the trends reported in magnetized ICF experiments [33–38]. Specifically, Sio *et al.* [37] examined the effect of an external magnetic field in indirectly driven implosions and observed notable gains in deuterium-deuterium neutron yield and hot-spot temperature even at a modest ~ 12 T field. Increasing the applied field further to ~ 26 T produced a smaller additional performance gain, suggesting diminishing returns as the initial magnetic field was increased. However, these measurements were integrated over the implosion duration and included the combined influence of numerous processes, such as heat-flow inhibition, hydrodynamic compression, and radiation. They therefore did not directly isolate the role of magnetothermal insulation during the compression phase, nor experimentally determine the maximum possible temperature gain associated with suppressing cross-field electron heat transport. In contrast, the present work reaches this state in a controlled geometry, as shown by the negligible difference between the 10 and 20 T cases, the agreement with simple analytical estimates, and the asymptotic behavior reproduced by the VFP simulations.

V. SUMMARY AND CONCLUSIONS

In summary, we report the first direct, spatially and temporally resolved measurement of the saturation of heat-flux inhibition by an external, longitudinal magnetic field in a laser-produced plasma. We quantify both the maximum temperature enhancement achievable through magnetothermal insulation and the Hall parameter at which this limit is reached, extending previous work into the strongly magne-

tized regime [1]. Spatially and temporally resolved Thomson scattering shows that the temperature gain reaches a plateau once the Hall parameter χ_{Hall} approaches ~ 50 , with only a marginal additional increase between 10 and 20 T. The measured gain agrees with simple geometrical estimates, and VFP simulations reproduce the observed asymptotic behavior. In the strongly magnetized cases, we also report pronounced density cavitation and dense fronts, consistent with pressure-driven advection after cross-field heat losses are suppressed. Finally, our results identify a field strength threshold above which only limited additional benefit is expected in magnetized ICF implosions, potentially easing design constraints and target complexity [31,35,36], while also providing a quantitative benchmark for transport models in the $\chi_{\text{Hall}} \gtrsim 10$ collisional regime.

ACKNOWLEDGMENTS

The authors would like to thank the team at LULI2000 for their work in obtaining the data presented here. A.T. would also like to thank Archie Bott for insightful discussions. This work was supported by grants managed by l'Agence Nationale de la Recherche under the Investissements d'Avenir programs ANR-18-EURE-0014, ANR-10-LABX-0039-PALM, and ANR-22-CE30-0044. This project was provided with computer and storage resources by GENCI at TGCC, thanks to the allocation A0170515693 on the supercomputer Joliot Curie's ROME partition. N.O. was supported by grants from Japan Society for the Promotion of Science (JSPS) KAKENHI (Grant No. 23K20038) and JSPS Core-to-Core program (Grant No. JPJSCCA20230003). This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (Grant Agreement No. 101052200—EUROfusion). The involved teams have operated within the framework of the Enabling Research project AWP24-ENR-IFE.02.CEA-01 "Magnetized ICF".

The authors have no conflicts to disclose.

Views and opinions expressed are those of the author(s) only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.

DATA AVAILABILITY

The data that support the findings of this article are not publicly available upon publication because it is not technically feasible and/or the cost of preparing, depositing, and hosting the data would be prohibitive within the terms of this research project. The data are available from the authors upon reasonable request.

APPENDIX A: DETAILED TABLE OF EXPERIMENTAL PARAMETERS AND QUANTITIES

The parameter space accessed in the experimental campaign described in this paper is summarized in Table I. TS, Mach-Zehnder interferometry, and streaked optical pyrometry (SOP) were used.

TABLE I. The parameter space explored in this work and the respective dimensionless numbers derived for the relevant time range. TS, Mach-Zehnder interferometry, and SOP were used to infer the relevant plasma quantities for H₂, He, and N₂ gas jets over a 0–3 ns probing window. The dimensionless numbers listed were calculated using the nominal magnetic field value, i.e., assuming no compression or cavitation.

Quantity	Parameter range	Diagnostic
Density (n_e)	$1\text{--}60 \times 10^{18} \text{ cm}^{-3}$	TS and interferometry
Temperature (T_e)	60–1100 eV	TS
Velocity (v)	100–400 km/s	SOP
Magnetic field (B)	0–20 T (along the x axis)	Rogowski probe
Gas backing pressure (P_B)	0.5–14 bar	Manometer
Knudsen number (Kn)	10^{-6} to 0.5	TS and interferometry
Electron Hall parameter (χ_{Hall})	0–250	TS and interferometry
Magnetic Reynolds number (Re_m)	950–9000	TS, interferometry, and SOP
Magnetosonic Mach number (M_{ms})	3–15	SOP

APPENDIX B: IMPACT OF THE PROBE BEAM ON THE THOMSON-SCATTERING SPECTRA

We assessed and quantified the influence of the probe beam on the gas jet by conducting multiple probe-only shots and collecting the Thomson-scattering spectra. Figure 7 shows spatially and temporally resolved Thomson-scattering results for an H₂ gas jet ($P_B = 14$ bar and $B = 0$ T). From these measurements, the probe beam clearly ionizes the gas jet. Over ~ 1.5 ns, it produces electron densities $n_e = (0.5\text{--}2.75) \times 10^{19} \text{ cm}^{-3}$ and electron temperatures $T_e = 80\text{--}230$ eV, with the peak values occurring at early times and near $y = 0$. Notably, n_e never reaches the maximum value of $4 \times 10^{19} \text{ cm}^{-3}$ (see Fig. 5), indicating that the probe beam does not fully ionize the gas.

The maximum T_e in the probe-only case is 227 eV (at 0.3 ns and $y = 0$). This is far smaller than in heater-probe shots, where $T_e > 600$ eV in the center of the plasma and can reach up to ~ 800 eV (see Fig. 4). The region where the probe beam could, in principle, matter more is the edge of the plasma ($y < -500 \mu\text{m}$ and $y > 500 \mu\text{m}$), where the heater beam does not irradiate the gas jet. However, the probe-only data show only modest edge heating, with $T_e \lesssim 110$ eV at $t = 1$ ns [see Fig. 7(c)]. In contrast, Fig. 4 indicates T_e values of about 200–400 eV in the same regions during heater-probe shots. This implies that heat conduction is the dominant mechanism shaping the temperature profile there, rather than direct probe-beam heating. In fact, nonlocal heat-conduction effects (such as preheating) are expected and observed there, but a




FIG. 7. Spatially and temporally resolved, probe-beam-only Thomson-scattering spectra for H₂ ($P_B = 14$ bar), $B = 0$ T, and $E = 3.1$ J. The spatial profile is recorded at $t = 1$ ns and the temporal evolution is captured at $y = 0$.

detailed discussion is outside the scope of this work and will be presented in a future publication.

Taken together, the probe-only measurements provide an upper bound on the probe's ability to perturb the plasma. In heater-probe shots, the probe is fired simultaneously with or after the heater beam, so it propagates through already heated plasma; its inverse-bremsstrahlung heating therefore decreases as $(\partial T_e / \partial t)_{IB} \propto T_e^{-3/2}$ [see Eq. (2)] [51–53]. Accordingly, the probe's impact in heater-probe shots is even smaller than what the probe-only results suggest, at least in the central region. This is important for interpreting the Thomson-scattering spectra, because it confirms that the probe beam does not meaningfully disturb the plasma it is intended to diagnose.

APPENDIX C: MAGNETOTHERMAL CONDUCTION INHIBITION IN N₂

In addition to the H₂ shots discussed in the main text, we also irradiated N₂ gas jets with the same heater and probe beams. The N₂ results support the same overall conclusions, but the collected electronic Thomson-scattering spectra are noisier because the background signal (bremsstrahlung radiation) increases with the atomic number Z . For this reason, this work focuses on the H₂ dataset.

Compared to H₂, N₂ reaches a higher T_e at 20 T because inverse-bremsstrahlung heating is more efficient at larger Z^* [see Eq. (2)] [51–53]. This dataset also allows us to more directly compare our findings with the results reported by Froula *et al.* [1], who studied an N₂ gas jet driven by a pump beam similar to the LULI2000 laser system and applied magnetic fields of 4 and 12 T. Their main conclusions were an increase in T_e from ~ 200 to ~ 800 eV at $t = 0.5$ ns and 12 T.

While Fig. 8 does exhibit a substantial increase in T_e when an external magnetic field is applied, it is notably less than what Froula *et al.* [1] found from the ion-acoustic feature of




FIG. 8. (Top) Spatially resolved electronic Thomson-scattering images of an N₂ gas-jet plasma ($P_B = 0.5$ bar) at 1 ns for (a) 0 T and (b) 20 T. (Bottom) The respective spatial (c) n_e and (d) T_e profiles. The shaded uncertainty bands are displayed at half their actual width for visual clarity.

the Thomson-scattering spectrum. We observe a T_e increase from ~ 750 eV up to ~ 1050 eV ($\sim 1.4\times$), whereas Ref. [1] reported a $4\times$ increase in Fig. 4 of their work. As discussed in the main text, these differences may be reconciled by the more pronounced nonlocal behavior in the unmagnetized case due to the smaller focal spot in Ref. [1]. That would lower the unmagnetized electron temperature and subsequently raise the gain once the magnetic field suppresses cross-field transport.

- [1] D. H. Froula, J. S. Ross, B. B. Pollock, P. Davis, A. N. James, L. Divol, M. J. Edwards, A. A. Offenberger, D. Price, R. P. J. Town, *et al.*, Quenching of the nonlocal electron heat transport by large external magnetic fields in a laser-produced plasma measured with imaging Thomson scattering, *Phys. Rev. Lett.* **98**, 135001 (2007).
- [2] C. P. Ridgers, R. J. Kingham, and A. G. R. Thomas, Magnetic cavitation and the reemergence of nonlocal transport in laser plasmas, *Phys. Rev. Lett.* **100**, 075003 (2008).
- [3] C. Arran, P. Bradford, A. Dearling, G. S. Hicks, S. Al-Atabi, L. Antonelli, O. C. Ettlinger, M. Khan, M. P. Read, K. Glize, *et al.*, Measurement of magnetic cavitation driven by heat flow in a plasma, *Phys. Rev. Lett.* **131**, 015101 (2023).
- [4] Z. Wang, R. Cheng, G. Wang, X. Jin, Y. Tang, Y. Chen, Z. Zhou, L. Shi, Y. Wang, Y. Lei, *et al.*, Observation of plasma dynamics in a theta pinch by a novel method, *Matter Radiat. Extremes* **8**, 045901 (2023).
- [5] A. Triantafyllidis, J.-R. Marquès, Y. Benkadoum, Y. De León, A. Ciardi, J. Béard, J.-M. Lagarrigue, A. Dearling, N. Ozaki, M. Koenig, *et al.*, Dynamics and energy dissipation of collisional blast waves in a perpendicular magnetic field, *Phys. Plasmas* **32**, 022102 (2025).
- [6] Z. Wang, R. Cheng, X. Jin, Y. Chen, L. Shi, G. Wang, Z. Zhou, Z. Iqbal, Y. Chen, J. Zhang, *et al.*, Magnetic transport and radiation properties during compression of a magnetized plasma, *Matter Radiat. Extremes* **10**, 037401 (2025).
- [7] A. Nishiguchi, T. Yabe, M. G. Haines, M. Psimopoulos, and H. Takewaki, Convective amplification of magnetic fields in laser-produced plasmas by the Nernst effect, *Phys. Rev. Lett.* **53**, 262 (1984).
- [8] T. H. Kho and M. G. Haines, Nonlinear kinetic transport of electrons and magnetic field in laser-produced plasmas, *Phys. Rev. Lett.* **55**, 825 (1985).
- [9] J. F. Luciani, P. Mora, and A. Bendib, Magnetic field and nonlocal transport in laser-created plasmas, *Phys. Rev. Lett.* **55**, 2421 (1985).
- [10] T. H. Kho and M. G. Haines, Nonlinear electron transport in magnetized laser plasmas, *Phys. Fluids* **29**, 2665 (1986).
- [11] L. Willingale, A. G. R. Thomas, P. M. Nilson, M. C. Kaluza, S. Bandyopadhyay, A. E. Dangor, R. G. Evans, P. Fernandes,

- M. G. Haines, C. Kamperidis, *et al.*, Fast advection of magnetic fields by hot electrons, *Phys. Rev. Lett.* **105**, 095001 (2010).
- [12] L. Lancia, B. Albertazzi, C. Boniface, A. Grisollet, R. Riquier, F. Chaland, K.-C. Le Thanh, P. Mellor, P. Antici, S. Buffechoux, *et al.*, Topology of megagauss magnetic fields and of heat-carrying electrons produced in a high-power laser-solid interaction, *Phys. Rev. Lett.* **113**, 235001 (2014).
- [13] A. S. Joglekar, C. P. Ridgers, R. J. Kingham, and A. G. R. Thomas, Kinetic modeling of Nernst effect in magnetized hohlraums, *Phys. Rev. E* **93**, 043206 (2016).
- [14] J. P. Brodrick, R. J. Kingham, M. M. Marinak, M. V. Patel, A. V. Chankin, J. T. Omotani, M. V. Umansky, D. Del Sorbo, B. Dudson, J. T. Parker, *et al.*, Testing nonlocal models of electron thermal conduction for magnetic and inertial confinement fusion applications, *Phys. Plasmas* **24**, 092309 (2017).
- [15] L. Spitzer Jr. and R. Härm, Transport phenomena in a completely ionized gas, *Phys. Rev.* **89**, 977 (1953).
- [16] S. I. Braginskii, Transport processes in a plasma, *Rev. Plasma Phys.* **1**, 205 (1965).
- [17] D. Shvarts, J. Delettrez, R. L. McCrory, and C. P. Verdon, Self-consistent reduction of the Spitzer-Härm electron thermal heat flux in steep temperature gradients in laser-produced plasmas, *Phys. Rev. Lett.* **47**, 247 (1981).
- [18] J. F. Luciani, P. Mora, and J. Virmont, Nonlocal heat transport due to steep temperature gradients, *Phys. Rev. Lett.* **51**, 1664 (1983).
- [19] Y. T. Lee and R. M. More, An electron conductivity model for dense plasmas, *Phys. Fluids* **27**, 1273 (1984).
- [20] E. M. Epperlein and M. G. Haines, Plasma transport coefficients in a magnetic field by direct numerical solution of the Fokker-Planck equation, *Phys. Fluids* **29**, 1029 (1986).
- [21] G. P. Schurtz, P. D. Nicolai, and M. Busquet, A nonlocal electron conduction model for multidimensional radiation hydrodynamics codes, *Phys. Plasmas* **7**, 4238 (2000).
- [22] P. D. Nicolai, J.-L. A. Feugeas, and G. P. Schurtz, A practical nonlocal model for heat transport in magnetized laser plasmas, *Phys. Plasmas* **13**, 032701 (2006).
- [23] J. D. Sadler, H. Li, and K. A. Flippo, Parameter space for magnetization effects in high-energy-density plasmas, *Matter Radiat. Extremes* **6**, 065902 (2021).
- [24] J. D. Sadler, C. A. Walsh, and H. Li, Symmetric set of transport coefficients for collisional magnetized plasma, *Phys. Rev. Lett.* **126**, 075001 (2021).
- [25] H. Zhao, Z. Sheng, and S. Weng, Nonlocal thermal transport in magnetized plasma along different directions, *Matter Radiat. Extremes* **7**, 045901 (2022).
- [26] S. A. Slutz and R. A. Vesey, High-gain magnetized inertial fusion, *Phys. Rev. Lett.* **108**, 025003 (2012).
- [27] L. J. Perkins, D. D.-M. Ho, B. G. Logan, G. B. Zimmerman, M. A. Rhodes, D. J. Strozzi, D. T. Blackfield, and S. A. Hawkins, The potential of imposed magnetic fields for enhancing ignition probability and fusion energy yield in indirect-drive inertial confinement fusion, *Phys. Plasmas* **24**, 062708 (2017).
- [28] C. A. Walsh, K. McGlinchey, J. K. Tong, B. D. Appelbe, A. Crilly, M. F. Zhang, and J. P. Chittenden, Perturbation modifications by pre-magnetisation of inertial confinement fusion implosions, *Phys. Plasmas* **26**, 022701 (2019).
- [29] C. A. Walsh, S. O'Neill, J. P. Chittenden, A. J. Crilly, B. Appelbe, D. J. Strozzi, D. Ho, H. Sio, B. Pollock, L. Divol, *et al.*, Magnetized ICF implosions: Scaling of temperature and yield enhancement, *Phys. Plasmas* **29**, 042701 (2022).
- [30] D. J. Strozzi, H. Sio, G. B. Zimmerman, J. D. Moody, C. R. Weber, B. Z. Djordjević, C. A. Walsh, B. A. Hammel, B. B. Pollock, A. Povilus, *et al.*, Design and modeling of indirectly driven magnetized implosions on the NIF, *Phys. Plasmas* **31**, 092703 (2024).
- [31] C. A. Walsh, S. T. O'Neill, D. J. Strozzi, L. S. Leal, R. Spiers, A. J. Crilly, B. Pollock, H. Sio, B. Hammel, B. Z. Djordjević, *et al.*, Magnetized ICF implosions: Ignition at low laser energy using designs with more ablator mass remaining, *Phys. Plasmas* **32**, 072715 (2025).
- [32] S. T. O'Neill, B. D. Appelbe, A. J. Crilly, C. A. Walsh, D. J. Strozzi, J. D. Moody, and J. P. Chittenden, Burn propagation in magnetized high-yield inertial fusion, *Phys. Plasmas* **32**, 022703 (2025).
- [33] P. Y. Chang, G. Fiksel, M. Hohenberger, J. P. Knauer, R. Betti, F. J. Marshall, D. D. Meyerhofer, F. H. Séguin, and R. D. Petrasso, Fusion yield enhancement in magnetized laser-driven implosions, *Phys. Rev. Lett.* **107**, 035006 (2011).
- [34] D. S. Montgomery, B. J. Albright, D. H. Barnak, P. Y. Chang, J. R. Davies, G. Fiksel, D. H. Froula, J. L. Kline, M. J. MacDonald, A. B. Sefkow, *et al.*, Use of external magnetic fields in hohlraum plasmas to improve laser-coupling, *Phys. Plasmas* **22**, 010703 (2015).
- [35] J. D. Moody, B. B. Pollock, H. Sio, D. J. Strozzi, D. D.-M. Ho, C. A. Walsh, G. E. Kemp, B. Lahmann, S. O. Kucheyev, B. Kozioziemski, *et al.*, Increased ion temperature and neutron yield observed in magnetized indirectly driven D₂-filled capsule implosions on the National Ignition Facility, *Phys. Rev. Lett.* **129**, 195002 (2022).
- [36] J. D. Moody, B. B. Pollock, H. Sio, D. J. Strozzi, D. D.-M. Ho, C. Walsh, G. E. Kemp, S. O. Kucheyev, B. Kozioziemski, E. G. Carroll, *et al.*, The magnetized indirect drive project on the National Ignition Facility, *J. Fusion Energy* **41**, 7 (2022).
- [37] H. Sio, J. D. Moody, B. B. Pollock, D. J. Strozzi, D. D.-M. Ho, C. A. Walsh, G. E. Kemp, B. Lahmann, S. O. Kucheyev, B. Kozioziemski, *et al.*, Performance scaling with an applied magnetic field in indirect-drive inertial confinement fusion implosions, *Phys. Plasmas* **30**, 072709 (2023).
- [38] M. Bailly-Grandvaux, R. Florido, C. A. Walsh, G. Pérez-Callejo, F. N. Beg, *et al.*, Impact of strong magnetization in cylindrical plasma implosions with applied *B*-field measured via X-ray emission spectroscopy, *Phys. Rev. Res.* **6**, L012018 (2024).
- [39] R. Riquier, Magnetic field in laser plasmas: Non-local electron transport and reconnection, Ph.D. thesis, Université Paris Saclay (COMUE), 2016.
- [40] B. Albertazzi, J. Béard, A. Ciardi, T. Vinci, J. Albrecht, J. Billette, T. Burris-Mog, S. N. Chen, D. Da Silva, S. Dittrich, *et al.*, Production of large volume, strongly magnetized laser-produced plasmas by use of pulsed external magnetic fields, *Rev. Sci. Instrum.* **84**, 043505 (2013).
- [41] B. Albertazzi, S. N. Chen, P. Antici, J. Böker, M. Borghesi, J. Breil, V. Dervieux, J. L. Feugeas, L. Lancia, M. Nakatsutsumi, *et al.*, Dynamics and structure of self-generated magnetic fields on solids following high contrast, high intensity laser irradiation, *Phys. Plasmas* **22**, 123108 (2015).
- [42] D. B. Schaeffer, A. F. A. Bott, M. Borghesi, K. A. Flippo, W. Fox, J. Fuchs, C. Li, F. H. Séguin, H.-S. Park, P. Tzeferacos,

- et al.*, Proton imaging of high-energy-density laboratory plasmas, *Rev. Mod. Phys.* **95**, 045007 (2023).
- [43] J. A. Percy, G. D. Sutcliffe, T. M. Johnson, B. L. Reichelt, S. G. Dannhoff, Y. Lawrence, J. Frenje, M. Gatu-Johnson, R. D. Petrasso, and C. Li, Hohlraum fields with monoenergetic proton radiography at OMEGA, *Appl. Opt.* **63**, A98 (2024).
- [44] J. Zheng, C. X. Yu, and Z. J. Zheng, Effects of non-Maxwellian (super-Gaussian) electron velocity distribution on the spectrum of Thomson scattering, *Phys. Plasmas* **4**, 2736 (1997).
- [45] B. C. Foo, D. B. Schaeffer, and P. V. Heuer, Recovering non-Maxwellian particle velocity distribution functions from collective Thomson-scattered spectra, *AIP Adv.* **13**, 115328 (2023).
- [46] R. C. Malone, R. L. McCrory, and R. L. Morse, Indications of strongly flux-limited electron thermal conduction in laser-target experiments, *Phys. Rev. Lett.* **34**, 721 (1975).
- [47] A. R. Bell, R. G. Evans, and D. J. Nicholas, Electron energy transport in steep temperature gradients in laser-produced plasmas, *Phys. Rev. Lett.* **46**, 243 (1981).
- [48] R. J. Henchen, M. Sherlock, W. Rozmus, J. Katz, D. Cao, J. P. Palastro, and D. H. Froula, Observation of nonlocal heat flux using Thomson scattering, *Phys. Rev. Lett.* **121**, 125001 (2018).
- [49] E. Fabre, F. Amiranoff, R. Fabbro, C. Garban-Labaune, J. Virmont, M. Weinfeld, F. David, and R. Pellat, Effect of laser wavelength on interaction studies for inertial fusion, in *Plasma Physics and Controlled Nuclear Fusion Research 1980* (IAEA, Vienna, 1981), Vol. 2, pp. 263–272.
- [50] C. Garban-Labaune, E. Fabre, C. E. Max, R. Fabbro, F. Amiranoff, J. Virmont, M. Weinfeld, and A. Michard, Effect of laser wavelength and pulse duration on laser-light absorption and back reflection, *Phys. Rev. Lett.* **48**, 1018 (1982).
- [51] A. B. Langdon, Nonlinear inverse bremsstrahlung and heated-electron distributions, *Phys. Rev. Lett.* **44**, 575 (1980).
- [52] R. J. Kingham and A. R. Bell, An implicit Vlasov–Fokker–Planck code to model non-local electron transport in 2-D with magnetic fields, *J. Comput. Phys.* **194**, 1 (2004).
- [53] C. P. Ridgers, A. G. R. Thomas, R. J. Kingham, and A. P. L. Robinson, Transport in the presence of inverse bremsstrahlung heating and magnetic fields, *Phys. Plasmas* **15**, 092311 (2008).
- [54] R. P. Drake, *Introduction to High-Energy-Density Physics* (Springer, 2018), p. 127.
- [55] A. Triantafyllidis, Laser-produced magnetized plasmas: Heat transport and blast waves, Ph.D. thesis, Institut Polytechnique de Paris, 2025.
- [56] C. Arran, C. P. Ridgers, and N. C. Woolsey, Proton radiography in background magnetic fields, *Matter Radiat. Extremes* **6**, 046904 (2021).
- [57] M. Borghesi, A. J. MacKinnon, A. R. Bell, R. Gaillard, and O. Willi, Megagauss magnetic field generation and plasma jet formation on solid targets irradiated by an ultraintense picosecond laser pulse, *Phys. Rev. Lett.* **81**, 112 (1998).
- [58] A. Triantafyllidis, J.-R. Marquès, S. Ferri, A. Calisti, Y. Benkadoum, Y. De León, A. Dearling, A. Ciardi, J. Béard, J.-M. Lagarrigue, *et al.*, Zeeman splitting observations in laser-produced magnetized blast waves, *Matter Radiat. Extremes* **10**, 047603 (2025).
- [59] Private communication. ALADIN is a Vlasov-Fokker-Planck code developed at CEA by D. Deck and K.-C. Le Thanh based on Refs. [52,63].
- [60] A. Chrisment, P. Loiseau, J.-L. Feugeas, P.-E. Masson-Laborde, J. Mathiaud, V. Tikhonchuk, and P. Nicolai, Analysis of a kinetic model for electron heat transport in inertial confinement fusion plasmas, *Phys. Plasmas* **29**, 062301 (2022).
- [61] A. Chrisment, A. Debayle, J.-L. Feugeas, P. Loiseau, P.-E. Masson-Laborde, J. Mathiaud, P. Nicolai, and V. Tikhonchuk, A deterministic numerical scheme for an electron heat transport model, *Math. Comput. Simul.* **205**, 78 (2023).
- [62] L. Gao, B. F. Kraus, K. W. Hill, M. B. Schneider, A. Christopherson, B. Bachmann, M. Bitter, P. Efthimion, N. Pablant, R. Betti, *et al.*, Hot spot evolution measured by high-resolution X-ray spectroscopy at the National Ignition Facility, *Phys. Rev. Lett.* **128**, 185002 (2022).
- [63] C. P. Ridgers, Magnetic fields and non-local transport in laser plasmas, Ph.D. thesis, Department of Physics, The Blackett Laboratory, Imperial College, 2008.