

Quantum-selected configuration interaction: Classical diagonalization of Hamiltonians in subspaces selected by quantum computers

Keita Kanno ^{1,*} Masaya Kohda,^{1,†} Ryosuke Imai ^{1,‡} Sho Koh ¹ Kosuke Mitarai,^{2,3}
Wataru Mizukami,^{2,3} and Yuya O. Nakagawa ¹

¹*QunaSys Inc., Aqua Hakusan Building 9F, 1-13-7 Hakusan, Bunkyo, Tokyo 113-0001, Japan*

²*Graduate School of Engineering Science, Osaka University, 1-3 Machikaneyama, Toyonaka, Osaka 560-8531, Japan*

³*Center for Quantum Information and Quantum Biology, Osaka University, 1-2 Machikaneyama, Toyonaka, Osaka 560-0043, Japan*



(Received 17 March 2023; accepted 9 May 2026; published 9 June 2026)

We propose quantum-selected configuration interaction, a class of hybrid quantum-classical algorithms for calculating the ground- and excited-state energies of many-electron Hamiltonians on noisy quantum devices. Suppose that an approximate ground state can be prepared on a quantum computer either by variational quantum eigensolver or by some other method. Then, by sampling the state in the computational basis, which is hard for classical computation in general, one can identify the electron configurations that are important for reproducing the ground state. The Hamiltonian in the subspace spanned by those important configurations is diagonalized on classical computers to output the ground-state energy and the corresponding eigenvector. The excited-state energies can be obtained similarly. The result is robust against statistical and physical errors because the noisy quantum devices are used only to define the subspace, and the resulting ground-state energy strictly satisfies the variational principle even in the presence of such errors. The expectation values of various other operators can also be estimated for obtained eigenstates with no additional quantum cost, since the explicit eigenvectors in the subspaces are known. We verified our proposal by numerical simulations and demonstrated it on a quantum device for an eight-qubit molecular Hamiltonian. The proposed algorithms are potentially feasible to tackle some challenging molecules by exploiting quantum devices with several tens of qubits, assisted by high-performance classical computing resources for diagonalization.

DOI: [10.1103/dmn4-snfx](https://doi.org/10.1103/dmn4-snfx)

I. INTRODUCTION

Recent years have seen a rapid development of quantum computers toward their practical use. Although current quantum devices are prone to errors due to physical noise, ways to achieve *quantum advantage* over classical computations have been explored experimentally [1–3], and such noisy intermediate-scale quantum (NISQ) devices are believed to become useful in the near future [4]. Quantum chemistry is at the top of the list of such useful applications (see, e.g., Refs. [5–9]): For instance, energy eigenvalues of a molecular Hamiltonian can be calculated by quantum algorithms developed for NISQ devices, where the most notable is the variational quantum eigensolver (VQE) [10] to find the ground-state energy.

However, VQE faces several challenges to be overcome for practical use. The major obstacle comes from errors caused

by statistical fluctuation and physical noise inherent in the noisy devices. Suppressing the statistical error to a practically acceptable level needs a prohibitively large number of samples [11], and error-mitigation techniques [12–21] for reducing physical noise require even more samples to compensate for the additional statistical error they introduce [22–25]. In particular, the effect of the errors can spoil the *variational* nature of VQE: That is, the energy estimated by quantum devices is not guaranteed to give an upper bound on the exact ground-state energy. Violations of the variational principle arise not only from the statistical fluctuations [11] but also from error-mitigation techniques such as zero-noise extrapolation [13]. It is problematic because lowering the resulting energy of VQE does not necessarily mean approaching the exact ground state. While such violations may not be as severe as those seen in, for example, coupled-cluster theory [26,27], they still pose a challenge when aiming for high-accuracy results. Besides, there are other challenges for VQE such as the barren plateau problem, which can interrupt the optimization [28].

In this paper, we propose a class of hybrid quantum-classical methods, which we call quantum-selected configuration interaction (QSCI), to find low-lying eigenvalues and eigenstates of a many-electron Hamiltonian [29]. QSCI is noise resilient and, in principle, free of costly optimization of parametrized quantum circuits. In particular, QSCI sets rigorous upper bounds on the ground-state energy [30] even under the effect of physical and statistical errors. Here, we outline a version of QSCI for finding a ground state: Suppose

*Contact author: kanno@qunasys.com

†Contact author: kohda@qunasys.com

‡Present address: QuEL, Inc., Hachioji ON Building 5F, 4-7-14 Myojincho, Hachioji, Tokyo 192-0046, Japan

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

that an approximate ground state, which we call an *input state* in this paper, can be prepared on a quantum computer; one then repeats a measurement of the state to identify the computational basis states (or electron configurations) that are important to express the ground state [31]; one then diagonalizes, on classical computers, the truncated Hamiltonian matrix in the subspace spanned by the identified configurations to obtain the smallest eigenvalue and eigenvector. The resulting eigenvalue approximates the ground-state energy. The diagonalization is classically tractable unless the number of selected configurations is exponentially large in the system size. The algorithm can be extended to find excited states by enlarging the subspace or by repeating the procedure for each energy eigenstate.

Since the matrix elements of the Hamiltonian in the computational basis can be exactly calculated on classical computers, the diagonalization results in an energy that gives a definite upper bound on the exact ground-state energy, regardless of the quality of the subspace spanned by the identified configurations; the quality only affects how tight the bound is. The states need to be measured only in the computational basis, and thus no additional gate operation is required for the measurement. In the presence of symmetries with conserved quantities such as the particle number, the postselection of the computational basis states in the sampling outcome allows one to mitigate the bit-flip errors. We experimentally demonstrate the effectiveness of the postselection in this paper. The algorithm may take any quantum states as the input states if they share the relevant electronic configurations with the true eigenstates and can be efficiently prepared on quantum devices. Such input states may be prepared, e.g., by parametrized quantum circuits moderately optimized via VQE and its variants [9], and other preparation schemes are discussed in Sec. V C. Sampling from such quantum states can be hard for classical computers [32–34], thereby providing a potential quantum speedup in QSCI.

QSCI is particularly advantageous for estimating expectation values of multiple observables with respect to energy eigenstates, as additional observables can be evaluated without extra quantum computational cost. This is achieved by constructing classical representations of approximate eigenstates via diagonalization of the Hamiltonian in a reduced, problem-adapted subspace, which is spanned by configurations sampled on a quantum computer. Unlike tomography techniques such as classical shadows [35,36], neural network tomography [37], and tensor network tomography [38], QSCI explicitly assumes that the target state is an eigenstate of a given Hamiltonian, which allows it to refine eigenstate approximations effectively without requiring full state tomography.

As the name suggests, QSCI can be viewed as a configuration interaction (CI), where the many-body basis set is determined by quantum computers via sampling of an input state. There are established techniques [39] that choose fixed basis sets. A common approach in electronic structure theory is to select only one- and two-particle excitations from a reference wavefunction. When the reference wavefunction is chosen to be Hartree-Fock, the resulting method is known as CI with singles and doubles (CISD). If the reference wavefunction is a correlated wavefunction beyond the mean-field

approximation, the method is called multireference CISD (MR-CISD). In the context of quantum computing, MR-CISD has sometimes been called as quantum subspace expansion (QSE) [40–42].

Another approach is the adaptive selection of a suitable basis set for a target system. In quantum chemistry, a systematic selection of important bases has a long history [43–52]. And there are recently active studies along with such a systematic selected CI [53–74]. Thanks to such developments, systematic selected CIs are now gradually being considered as a promising approach for large-scale quantum chemical simulations. The likely reason for this revival is that selected CI is an algorithm that can be adapted to current classical computer architectures with sufficient memory. QSCI may be seen as a systematic selected CI that utilizes quantum computers.

Our methods are capable of selecting electron configurations that are necessary to describe the eigenstates to some accuracy but are missed in the conventional methods with a fixed basis set. Note that our methods call the diagonalization procedure at most only once for each eigenstate, while the adaptive methods iteratively repeat the diagonalization to search for a configuration to be added in the basis set; our methods require much less classical computational time compared to those adaptive methods.

The classical diagonalization is already utilized in various hybrid quantum-classical algorithms to find energy eigenstates. Most notable is QSE, which spans the subspace by states built upon the reference VQE state, and is widely used for various applications, e.g., excited-state calculations [40], band structure calculations [75], and noise reduction [19,40,41,76–78]. More generally, one can span the subspace by various methods [79–86], which are sometimes collectively called as the quantum subspace diagonalization. In those methods, however, the matrix elements of the subspace Hamiltonian are calculated on quantum computers and thus are subject to the physical and statistical errors. There is a proposal [87] where some of the matrix elements are classically calculated, but the method still requires some matrix elements that are efficiently computable only by quantum computers for a possible quantum speedup. In QSCI, on the other hand, all the matrix elements are classically computed, giving up the use of more complex and physically motivated states as basis states that define the subspace.

The rest of the paper is organized as follows. The proposed methods are introduced in Sec. II and numerically tested in Sec. III. A demonstration on a quantum device is presented in Sec. IV, along with a noisy simulation as a preparatory study. We discuss aspects of the proposed methods in Sec. V and finally conclude in Sec. VI. Details of the algorithms, numerical simulations, and experiment, as well as supplemental numerical results, are given in the Appendixes.

II. METHODS

In this section, we present the methods of QSCI. Two ways of implementation are introduced: single diagonalization scheme in Sec. II C 1 and sequential diagonalization scheme in Sec. II C 2. They are designed for finding multiple energy eigenstates and reduce to the same simplified method when used for finding the ground state alone. After intro-

ducing necessary ingredients, we begin with the algorithm specific to finding the ground state, which is simple and illustrative, and then proceed to the two methods that can also find excited states.

A. Preliminary

We consider electronic structure problems of molecules in the second-quantization formalism with the Born-Oppenheimer approximation. A Hamiltonian and wavefunctions for electrons, in this setup, can be mapped onto N_q qubits such that the Slater determinants [88] for the Hartree-Fock state and its excitations are associated with the computational basis states $|x\rangle$, where $x \in \{0, 1\}^{N_q}$ is an N_q -bit string (see, e.g., Refs. [5,6]). In the Jordan-Wigner mapping, which we adopt in the numerical study, N_q corresponds to the number of spin orbitals, and “1” or “0” represents whether each spin orbital is occupied or not. The methods can work with other mapping schemes such as the Bravyi-Kitaev mapping [89], although the fermion-qubit correspondence is less intuitive and the error mitigation (discussed later) is less effective. We denote the qubit Hamiltonian by $\hat{H}$. A linear combination of all the computational basis states,

$$|\psi\rangle = \sum_{x \in \{0,1\}^{N_q}} \alpha_x |x\rangle, \quad (1)$$

encompasses the full-CI wavefunction. Note that for a fixed number of electrons only a subset of the computational basis states is needed.

In the full-CI method, sets of the CI coefficients $\{\alpha_x\}$ that correspond to energy eigenstates are found by diagonalizing the Hamiltonian in the full Fock space. The method is costly due to the combinatorial growth of the Fock-space dimension as the number of spin orbitals increases. For reducing the computational cost, there exist various classical approaches that truncate the Fock space and approximate the sum in Eq. (1) using a fixed or adaptively selected basis set, as mentioned in the previous section. In line with these efforts, but from a different viewpoint, we propose methods that harness quantum computers to identify important computational basis states, or electron configurations, for truncating the Fock space.

B. QSCI for ground state

We now describe the explicit algorithms. We begin with the algorithm for finding the lowest eigenvalue and the corresponding eigenstate (ground state) of an electronic Hamiltonian $\hat{H}$ on N_q qubits. For simplicity, we assume that the ground state is unique. When the degeneracy exists, the algorithms given in the next subsection, which are aimed at finding multiple eigenstates, can be straightforwardly applied. Indeed, the algorithm introduced in this subsection is a special case of each of the two algorithms in the next subsection.

Let $|\psi_{\text{in}}\rangle$ be an input state, which approximates the ground state, and suppose $|\psi_{\text{in}}\rangle$ can be prepared by a quantum circuit with N_q qubits. Then, one prepares the input state on a quantum computer and measures the state in the computational basis, which results in an outcome bit string $x \in \{0, 1\}^{N_q}$. Repeating such a sampling procedure (or shot) for N_{shot} times, one counts how many times each x appears. Based on the

total sampling result, the most frequent R computational basis states are selected to define the set

$$\mathcal{S}_R = \{|x\rangle | x \in \{0, 1\}^{N_q}, R \text{ most frequent}\}, \quad (2)$$

where R is a positive integer manually determined. This is to truncate the Fock space. One may in principle include all the computational basis states appeared in the measurements, while choosing an appropriately small R can reduce the computational cost for diagonalization.

One then solves the eigenvalue problem in the subspace spanned by $\mathcal{S}_R$:

$$\mathbf{H}_R \mathbf{c} = E_R \mathbf{c}, \quad (3)$$

where $\mathbf{H}_R$ is the $R \times R$ Hermitian matrix defined by

$$(\mathbf{H}_R)_{xy} = \langle x | \hat{H} | y \rangle \quad \text{for } |x\rangle, |y\rangle \in \mathcal{S}_R, \quad (4)$$

and $\mathbf{c}$ is an eigenvector with eigenvalue E_R , satisfying $\mathbf{c}^\dagger \mathbf{c} = 1$. This step of the algorithm proceeds via classical computations: calculations of the matrix elements $\langle x | \hat{H} | y \rangle$ and the diagonalization of $\mathbf{H}_R$. The former calculations can be efficiently done by some classical method, e.g., by the Slater-Condon rules in the fermionic basis. The latter diagonalization is performed to obtain the smallest eigenvalue E_R and the eigenvector $\mathbf{c}$, which are output of the algorithm. See Sec. V A for further discussion on costs of these classical computations. Here, E_R approximates the exact ground-state energy of $\hat{H}$, while $\mathbf{c}$ approximately gives the (normalized) CI coefficients, or the vector representation of the ground state, respectively. The corresponding quantum state, which we call the *output state*, is constructed as

$$|\psi_{\text{out}}\rangle = \sum_{|x\rangle \in \mathcal{S}_R} c_x |x\rangle, \quad (5)$$

where c_x is an element of the eigenvector $\mathbf{c}$. The output state $|\psi_{\text{out}}\rangle$ approximates the true ground state of $\hat{H}$. We remark that one does not need to realize the output state on quantum computers. Retaining the eigenvector $\mathbf{c}$ as classical data is enough for the application explained below.

The output state can be used to estimate the expectation values of observables other than the Hamiltonian for the ground state, solely based on classical computations. Specifically, suppose that an observable in question is represented by a qubit operator $\hat{O}$. If the matrix elements $\langle x | \hat{O} | y \rangle$ can be efficiently computed on classical computers, so does the expectation value $\langle \psi_{\text{out}} | \hat{O} | \psi_{\text{out}} \rangle$, which is expected to give an approximation to the expectation value for the true ground state. In particular, if $\hat{O}$ can be expressed as a linear combination of poly(N_q) Pauli strings, which is the case in many physical quantities, its expectation value can be efficiently computed on classical computers.

Comments are in order for technical details. We identify the set $\mathcal{S}_R$ to span the subspace by sampling the input state. In this way, we expect that important computational basis states, or Slater determinants, to describe the ground-state wavefunction can be selected. This is because in the sampling procedure a bit string x occurs with the probability $|\langle x | \psi_{\text{in}} \rangle|^2$, while $\langle x | \psi_{\text{in}} \rangle$ gives the CI coefficient of the corresponding Slater determinant in the input wavefunction $|\psi_{\text{in}}\rangle$ [90]. Indeed, $\mathcal{S}_R$ gives the R Slater determinants with the largest

coefficients $|\langle x | \psi_{\text{in}} \rangle|$ in $|\psi_{\text{in}}\rangle$, under the ideal situation where physical noise can be ignored and the sampling is performed with an infinite number of shots. In passing, we sometimes adopt a method equivalent to this ideal situation to define $\mathcal{S}_R$ in the numerical study: That is, we just pick up the R Slater determinants with the largest absolute values of the CI coefficients in the input state, instead of performing actual sampling procedures. We call this method as the *idealized sampling* in this paper. Note that we assume the input state approximates the true ground state. This is just to ensure the two states share the important computational basis states, and there is no need for a precise agreement between the CI coefficients of the two states. Such an input state can be prepared, e.g., by a parametrized quantum circuit moderately optimized via VQE. In Sec. VC, we discuss methods to prepare the input state, including non-VQE-based ways.

The set $\mathcal{S}_R$ is defined in Eq. (2) by specifying R , the number of the computational basis states retained in the subspace. However, this is not the unique choice. For instance, one may define the set by taking all the computational basis states in the measurement outcome, as already mentioned. Or, one may instead set a threshold on the rate of occurrence f_x for an outcome x in the total sampling result, and then define an alternative set $\mathcal{S}_\epsilon = \{|x\rangle | f_x \geq \epsilon\}$ with a threshold parameter ϵ . For a proof-of-principle demonstration, we adopt Eq. (2) to define the subspace for diagonalizing the Hamiltonian in the rest of the paper.

In reality, physical noise and statistical fluctuation, the latter due to a finite number of shots, cannot be ignored, causing some errors in the output. However, the effect is only indirect and the method is robust against those errors: That is, the errors can degrade the quality of the selected subspace by missing important configurations or by picking up irrelevant configurations in the sampling procedures, but the lowest eigenvalue and eigenvectors are exact within the subspace. The latter point, the exactness within the subspace, results from the use of diagonalization for the matrix $\mathbf{H}_R$, whose elements are exactly computed. Consequently, the obtained energy E_R sets an upper bound on E_{exact} , the true ground-state energy of $\hat{H}$:

$$E_{\text{exact}} \leq E_R. \quad (6)$$

Note that this variational inequality holds even under statistical fluctuation and physical noise. The situation is in contrast with VQE, where such an inequality is not guaranteed, as the energy is directly measured on quantum computers and hence is susceptible to the errors [91]. It is also worth mentioning that, for a given sampling outcome, increasing R , the subspace size, always leads to a better approximation of the ground-state energy: $E_{\text{exact}} \leq E_{R_a} \leq E_{R_b}$ for $R_a > R_b$. This can be used to see if the calculation converges. On the other hand, smaller R can reduce the classical computational cost. Such a trade-off between the accuracy and cost is discussed in Sec. IIIC.

The algorithm finds the lowest-energy state in the subspace $\mathcal{S}_R$, which gives an approximation to the ground state in the *full* Fock space. When there exists symmetry in the Hamiltonian, there are associated conserved quantities, e.g., the total electron number N_e (or the charge of molecule) and the z component of total electron spin S_z . Given this, one may wish

to find the lowest-energy state in a specific symmetry sector. In such a case, the method can be similarly applied but by relying on the subspace with fixed conserved quantities. For N_e and S_z , this can be easily achieved as follows since each computational basis state corresponds to a Slater determinant with definite N_e and S_z : One prepares an input state with the desired values of (N_e, S_z) , for which the sampling results in configurations each with the desired (N_e, S_z) ; or, if such an input state cannot be prepared, one may postselect the sampling outcome, where one discards an outcome $x \in \{0, 1\}^{N_q}$ if it conflicts with the desired (N_e, S_z) . It is worth noting that the variational inequality (6) still holds in each sector of Fock space specified by (N_e, S_z) .

Physical noise can cause a contamination of symmetry sectors: For an input state with fixed (N_e, S_z) , sampling on a noisy device can result in electron configurations with unwanted values of (N_e, S_z) , due to the bit-flip noise [92] or readout error. Nevertheless, one can mitigate such errors by postselecting the sampling outcome according to the conserved quantities, as described above. One then diagonalizes the Hamiltonian in the postselected subspace. We find that the postselection is particularly effective to mitigate the readout error in the Jordan-Wigner mapping, while it is also applicable to other fermion-qubit mapping schemes (see Appendix A for discussions).

The algorithm is schematically summarized in Fig. 1.

C. QSCI for multiple energy eigenstates

We now extend the algorithm to find multiple energy eigenstates, including low-lying excited states. For this, we note that the previous algorithm can output multiple eigenvectors, which can be taken to approximate excited states as well as the ground state. Yet, the quality of the approximation would not be satisfactory for the excited states, as the subspace is tailored for the ground state. Hence, we introduce extra input states to construct subspace(s) that can capture the excited states. In the following, we present two distinct algorithms to find multiple energy eigenstates and energies, schematically shown in Fig. 2. The first algorithm, which we call the single diagonalization scheme, constructs a common subspace for both ground and excited states of interest, and performs the diagonalization in the subspace to simultaneously obtain all the desired eigenstates and energies. On the other hand, the second algorithm, dubbed as the sequential diagonalization scheme, constructs multiple subspaces, each tailored for each energy eigenstate, and sequentially diagonalizes the Hamiltonian in each subspace. Both of the algorithms contain the algorithm specific to the ground state, introduced in the preceding subsection, as a special case.

1. Single diagonalization scheme

Here, we describe the single diagonalization scheme. Suppose one seeks for N_s low-lying eigenstates of $\hat{H}$, which consist of the ground state(s) and subsequent excited states. In this case, one prepares multiple input states $|\psi_{\text{in}}^{(i)}\rangle$ ($i = 0, 1, \dots, N_{\text{in}} - 1$), which correspond to the low-lying energy eigenstates. Here, we allow $N_{\text{in}} \leq N_s$, although the natural choice would be $N_{\text{in}} = N_s$. For each of the input states, one repeats the sampling procedure as in the previous subsection.




FIG. 1. Schematic description of the QSCI algorithm for finding the ground state. When selecting the configurations, one may postselect the configurations by using conserved quantities such as the electron number or spin S_z to mitigate the errors.

One then obtains the set of important configurations $\mathcal{S}_{R_i}^{(i)}$, formed by most frequent R_i bit strings in the total sampling outcome for the i th input state. Combining all the sets $\mathcal{S}_{R_i}^{(i)}$, one constructs the common subspace [93]:

$$\mathcal{S}_R = \mathcal{S}_{R_0}^{(0)} \cup \mathcal{S}_{R_1}^{(1)} \cup \dots \cup \mathcal{S}_{R_{N_{\text{in}}-1}}^{(N_{\text{in}}-1)}. \quad (7)$$

In this case, the parameters R_i may be eigenstate dependent, while R is the number of elements in the common subspace $\mathcal{S}_R$. $R \geq N_s$ is required to yield at least N_s eigenvectors in the diagonalization procedure, as shortly explained.

One may treat all R_i as free parameters, which determine R in turn. Or, one may first choose a value for R and, then, decide each R_i following some strategy. There are various ways for the latter strategy, depending on the purpose of using

the algorithm. For example, if one prioritizes the ground state in terms of accuracy, a possible choice would be $R_0 = R$ and $R_i = 0$ for $i \neq 0$, albeit extreme. Or, if one wishes to treat all the input states on equal footing, each of R_i can be chosen as equal as possible [94].

With the common subspace $\mathcal{S}_R$ constructed, one then diagonalizes the Hamiltonian in $\mathcal{S}_R$ as in the previous subsection: One constructs the $R \times R$ Hermitian matrix $\mathbf{H}_R$, solves the eigenvalue equation $\mathbf{H}_R \mathbf{c} = E_R \mathbf{c}$, and then picks up N_s low-lying eigenvectors and eigenvalues, $(\mathbf{c}^{(0)}, E_R^{(0)})$, $(\mathbf{c}^{(1)}, E_R^{(1)})$, $\dots$, $(\mathbf{c}^{(N_s-1)}, E_R^{(N_s-1)})$, where $\mathbf{c}^{(i)\dagger} \mathbf{c}^{(j)} = \delta_{ij}$. Here, $E_R^{(i)}$ ($E_R^{(0)}$) approximates the true energy of the i th excited state (ground state), when the ground state is unique, for instance. The corresponding output states can be




FIG. 2. Schematic descriptions of the QSCI algorithms for finding the ground state and the first excited state: (a) single diagonalization scheme and (b) sequential diagonalization scheme. In both panels, $|\psi_{\text{in}}^{(0)}\rangle$ ($|\psi_{\text{in}}^{(1)}\rangle$) is the input state for the ground (first excited) state. In panel (b), the overlap term is constructed from the preobtained output state $|\psi_{\text{out}}^{(0)}\rangle$ to define the effective Hamiltonian for the first excited state, $\hat{H}^{(1)} = \hat{H} + \beta_0 |\psi_{\text{out}}^{(0)}\rangle \langle \psi_{\text{out}}^{(0)}|$.

constructed as

$$|\psi_{\text{out}}^{(i)}\rangle = \sum_{|x\rangle \in \mathcal{S}_R} c_x^{(i)} |x\rangle, \quad (8)$$

for $i = 0, 1, \dots, N_s - 1$.

Note that the algorithm in the previous subsection is a special case of the single diagonalization scheme with a single input state ($N_{\text{in}} = 1$). In this method, one can apply the same error-mitigation technique by the postselection as described in the previous subsection. The variational inequality now holds for each of energy eigenstates by Cauchy's interlace theorem [39] (see also Refs. [95,96]):

$$E_{\text{exact}}^{(i)} \leq E_R^{(i)}, \quad (9)$$

for $i = 0, 1, \dots, N_s - 1$, where $E_{\text{exact}}^{(i)}$ is the i th eigenvalue (in ascending order) by the exact diagonalization. We remark that QSE [40] and multistate-contracted VQE [83], which also rely on the subspace diagonalization to obtain excited states, need to measure matrix elements, while the current method exactly calculates the matrix elements. Hence, we expect our method to be more noise robust with the guarantee of the variational inequality.

2. Sequential diagonalization scheme

We now give another scheme of QSCI to find excited states. The sequential diagonalization finds the ground state(s) and subsequent excited states by sequential diagonalization procedures of the Hamiltonian $\hat{H}$ in distinct subspaces. The algorithm is similar to the variational quantum deflation (VQD) [97], a variant of VQE for excited states.

Suppose one seeks the k th excited state [98] given that preceding $(k - 1)$ excited states and ground state are already obtained by this method with the output states $|\psi_{\text{out}}^{(i)}\rangle$ ($i = 0, 1, \dots, k - 1$). As in the previous methods, one repeats the preparation and measurement of the input state $|\psi_{\text{in}}^{(k)}\rangle$ to obtain the set of important configurations:

$$\mathcal{S}_{R_k}^{(k)} = \{|x\rangle | x \in \{0, 1\}^{N_q}, R_k \text{ most frequent}\}. \quad (10)$$

One then has to find the lowest-energy state of $\hat{H}$ in this subspace, under the restriction that this state is orthogonal to the states already found, $|\psi_{\text{out}}^{(i)}\rangle$ ($i = 0, 1, \dots, k - 1$). This can be achieved by diagonalizing the following effective Hamiltonian [99] in the subspace spanned by $\mathcal{S}_{R_k}^{(k)}$:

$$\hat{H}^{(k)} = \hat{H} + \sum_{i=0}^{k-1} \beta_i |\psi_{\text{out}}^{(i)}\rangle \langle \psi_{\text{out}}^{(i)}|, \quad (11)$$

where β_i are real parameters, which need to be sufficiently large for ensuring the orthogonality. The additional terms correspond to the overlap terms in VQD. This is equivalent to solving the eigenvalue equation

$$\mathbf{H}_{R_k}^{(k)} \mathbf{c}^{(k)} = E_{R_k}^{(k)} \mathbf{c}^{(k)}, \quad (12)$$

and then pick up the smallest eigenvalue $E_{R_k}^{(k)}$ and eigenvector $\mathbf{c}^{(k)}$, normalized by $\mathbf{c}^{(k)\dagger} \mathbf{c}^{(k)} = 1$. Here, $\mathbf{H}_{R_k}^{(k)}$ is the $R_k \times R_k$ Hermitian matrix defined by

$$(\mathbf{H}_{R_k}^{(k)})_{xy} = \langle x | \hat{H}^{(k)} | y \rangle \quad \text{for } |x\rangle, |y\rangle \in \mathcal{S}_{R_k}^{(k)}, \quad (13)$$

whose matrix elements can be efficiently calculated by classical computations based on the expression

$$(\mathbf{H}_R^{(k)})_{xy} = \langle x | \hat{H} | y \rangle + \sum_{i=0}^{k-1} \beta_i c_x^{(i)} c_y^{(i)*}. \quad (14)$$

One then constructs the output state

$$|\psi_{\text{out}}^{(k)}\rangle = \sum_{|x\rangle \in \mathcal{S}_{R_k}^{(k)}} c_x^{(k)} |x\rangle, \quad (15)$$

which approximates the k th excited state.

Note that the expressions are specific to the k th excited state. In order to find entire (low-lying) spectrum, one has to repeat the above procedure sequentially, starting from $k = 0$, the ground state, which can be found by the ground-state algorithm already explained. This is similar to VQD, but the QSCI method does not require extra circuits to calculate the overlap terms.

The coefficients β_i can be chosen in the same manner as VQD. We want the smallest eigenvalue of $\mathbf{H}_{R_k}^{(k)}$ to approximate $E_{\text{exact}}^{(k)}$, the k th eigenvalue of $\hat{H}$. Following the discussion in Ref. [97], it suffices to choose $\beta_i > E_{\text{exact}}^{(k)} - E_{\text{exact}}^{(i)}$ for $i = 0, \dots, k - 1$; or, one may apply the looser condition of $\beta_i > 2 \sum_j |c_j|$, where c_j are coefficients in the qubit Hamiltonian $\hat{H} = \sum_j c_j P_j$, expressed by the Pauli strings P_j (see Appendix B 1 for details). In practice, the condition $\beta_i > E_{\text{exact}}^{(k)} - E_{\text{exact}}^{(i)}$ can be utilized if one has prior knowledge of the energy spectrum, e.g., based on variational quantum algorithms. Even if such information is not available, one may still rely on the looser condition $\beta_i > 2 \sum_j |c_j|$. Note that in the sequential diagonalization scheme, the variational inequality like Eq. (9) is not guaranteed due to the inexactness of the effective Hamiltonian, i.e., as Eq. (11) would be constructed only by approximate eigenstates in practice (see Appendix B 1 for further discussion).

III. BENCHMARK OF QSCI WITH NOISELESS SIMULATIONS

In this section, we test various aspects of QSCI for small molecules by noiseless numerical simulations, where the effects of physical noise are not included. In Secs. III A and III B, QSCI calculations are performed for ground states and excited states, using input states prepared by VQE and VQD [97], a variant of VQE for excited states. Then, the scalability of QSCI is examined in Sec. III C, and finally, the effect of the statistical error in QSCI is studied in Sec. III D. For the numerical simulations, a quantum-circuit simulation library Qulacs [101] is used with the help of QURI Parts [102], a library for developing quantum algorithms. The simulations in Sec. III D are carried out by the sampling simulator that takes into account the statistical error, while all the other simulations are performed by the state-vector simulator, where the expectation values are exactly calculated without errors.

For each simulation and experiment in this paper, the molecular Hamiltonian is first prepared as the second-quantized electronic Hamiltonian using the Born-Oppenheimer approximation with Hartree-Fock orbitals using the Slater-type orbitals (STO)-3G basis unless otherwise

stated, and converted to the qubit one by the Jordan-Wigner mapping. Active spaces are explicitly specified when employed, otherwise the full-space Hamiltonians are used. The electronic Hamiltonians are generated by OPENFERMION [103] interfaced with PYSCF [104]. The molecular geometries and other details are shown in Appendix C. Stable geometries are chosen for all the molecules except for the hydrogen chains, and a potential impact of unstable geometry is briefly analyzed in Appendix D 4.

A. QSCI for ground state

We first show the result of numerical simulation for ground state with input states prepared by noiseless VQE. We choose H_2O molecule with five active spatial orbitals and six active electrons as our problem, which leads to a ten-qubit Hamiltonian after the Jordan-Wigner mapping. In the VQE calculation, the parametrized quantum circuit is constructed by the real-valued symmetry-preserving ansatz [105,106] with the depth 10 and is optimized by the Broyden-Fletcher-Goldfarb-Shanno (BFGS) optimizer in the scientific library SCIPY [107]. See Appendix C 1 for details.

The QSCI calculation is performed for each iteration of the VQE optimization: Given the values of ansatz parameters obtained at the iteration, the input state is prepared by the parametrized quantum circuit with those values assigned; then, the QSCI calculation with the idealized sampling introduced in Sec. II B is performed to estimate the ground-state energy E_R for a given R , the number of configurations in the subspace $\mathcal{S}_R$. This calculation is repeated for all the iterations of VQE with different values of R . The effect of the uncertainty due to the finiteness of the number of shots is addressed later in Secs. III D, IV, and Appendix D 2.

In Fig. 3, the result is shown along with the optimization history of VQE: $E_R - E_{\text{exact}}$ is plotted (in hartree) for each optimization step of VQE, where E_{exact} is the ground-state energy obtained by the exact diagonalization in the active space, called the complete active space configuration interaction (CASSI). The energies obtained by VQE are shown in the same way.

Comparing the results at the last iteration in the plot, one can see that QSCI gives a lower energy than VQE for $R \gtrsim 16$. This shows that the method is able to improve the results of VQE even in the noiseless setting, where the effect of error mitigation is not present. We emphasize that, as discussed in Sec. II B, a lower energy by QSCI means that the energy is closer to the exact ground-state energy, which is manifested in the plot, where $E_R - E_{\text{exact}}$ is always positive. It is notable that we can already achieve the chemical accuracy [108] of 1.6×10^{-3} hartree with $R \sim 16$ while the CASSI in this case uses 100 determinants to express the ground state [109].

A similar tendency is observed for iterations of $\gtrsim 200$. Note, in this case, the VQE results already achieve the chemical accuracy. On the other hand, for intermediate iterations of 70–200, the VQE results do not reach the chemical accuracy, while QSCI can improve them to meet the chemical accuracy if $R \gtrsim 16$. This suggests that an intermediate result of VQE, which is not seeing convergence in the optimization yet, is already useful as an input state of QSCI, and that one can reduce the number of optimization steps for VQE by employing




FIG. 3. The result of QSCI, the proposed method, for the ground state of H_2O molecule by noiseless simulation, shown with optimization history of VQE, which is used to prepare the input states of QSCI. Each of the resulting energies is shown as the difference to the CASSI result E_{exact} in hartree. The dash-dotted line shows the result by the state-vector simulation of VQE. The lines specified by the parameter R show the results of QSCI, $E_R - E_{\text{exact}}$, for the given value of R , using the parametrized state at each iteration of VQE as the input state. The parameter R determines the classical computational cost for QSCI, as explained in the main text.

QSCI as a postprocessing. We note that the QSCI results do not monotonically decrease, as a QSCI calculation for an input state with a lower energy does not necessarily result in a lower output energy.

We emphasize again that QSCI is able to improve the energy accuracy significantly over that of the input state, demonstrating its potential to extract meaningful determinants even from imperfect input states. This supports the claim that a moderately accurate input state is sufficient to enable QSCI to yield sufficiently accurate results.

B. QSCI for excited states

We next show the results of noiseless simulations for excited states of H_2O using the two distinct implementations of QSCI presented in Sec. II C, namely, the single diagonalization and sequential diagonalization schemes, which take the input states for excited states as well as the ground state. The numerical setup is similar to the previous subsection, with some differences explained below. In the input-state preparation, we employ VQE for the ground state and VQD for excited states, each with the same ansatz and optimizer as in the previous subsection; we use the same ten-qubit Hamiltonian, but with the overlap terms and penalty terms [110–112] in VQD, for orthogonality between the eigenstates and for symmetry restrictions (charge neutrality for the molecule and $S_z = 0$ for the total electron spin) on the excited states. Un-

der the same symmetry restrictions, the first excited state is a triplet state (T_1) and the second excited state is a singlet state (S_1), according to the exact diagonalization. The VQD calculation for T_1 requires information of the ground state (S_0) to generate the overlap term, for which the ansatz state is used with the converged parameters in VQE. A similar procedure is applied for S_1 , but with the extra overlap term for T_1 added. With the input states for S_0 , T_1 , and S_1 , we perform QSCI calculations to find T_1 and S_1 , where the idealized sampling is used. See Appendix C 2 for details.

The results are shown in Fig. 4 along with the optimization history of VQD, in the similar way as Fig. 3, but for $|E_R^{(i)} - E_{\text{exact}}|$ ($i = 1$ for T_1 and $i = 2$ for S_1). At each iteration, three types of QSCI calculations are performed: sequential, single-ground-state, and single-mixed. Sequential diagonalization uses the T_1 (S_1) state at the iteration, and one (two) lower-energy state(s) at their final iterations as input states. Two single diagonalization methods use different input states: “single-ground-state” uses the ground state prepared by the converged VQE calculation, and that is why they are constant in the plot; on the other hand, “single-mixed” uses the two (three) states as input states, and selects R configurations so that each of the two (three) states contributes as equally as possible, as explained in Sec. II C 1. Note that R is the dimension of the common subspace $\mathcal{S}_R$ in Eq. (7). The coefficient(s) β_0 (and β_1 for S_1 state) of the overlap term(s) for orthogonality is set to $\beta_0 = \beta_1 = 1$ hartree, which is sufficiently larger than the energy gaps between the states in question. For sequential diagonalization, the values of R_i are fixed to $R_i = 16$ for $i = 0, 1$, and 2 , corresponding to S_0 , T_1 , and S_1 states, respectively; for single diagonalization, the value of R is set to $R = 16$, so that the sizes of the subspace Hamiltonian matrices to be diagonalized are the same among all the setups.

Comparing the three QSCI results for excited states, the sequential diagonalization performs the best except for the initial steps of iterations, where the quality of the input state is significantly low. Moreover, the sequential diagonalization outperforms the VQD calculation, even with a moderate value of $R_i = 16$. For some larger R , the single diagonalization is also expected to improve and eventually outperform the VQD result at the same iteration as it can achieve the same representability as the sequential one [113]. Although the sequential diagonalization seems to be better in terms of performance, it should be noted that there is no guarantee for the variational inequality in the sequential diagonalization. The inequality for excited states holds in the single diagonalization, as explained in Sec. II C.

C. Scaling of computational costs

We now investigate the scalability of the proposed method by estimating the classical and quantum computational costs to calculate the ground states for molecular Hamiltonians of various sizes. More concretely, we estimate the minimum value for R and the required number of shots N_{shot} to obtain the ground-state energy within an error ϵ for those Hamiltonians.

For this sake, we employ the chromium dimer Cr_2 with various active spaces and the linear hydrogen chains with different numbers of atoms. Both Cr_2 and hydrogen chains are known to be challenging molecules in quantum chemistry




FIG. 4. Same as Fig. 3 but for (a) (T_1) and (b) (S_1) excited states of H_2O with $S_z = 0$, along with optimization histories of VQD, shown by dotted lines, for input-state preparation; the energy differences are plotted by the absolute values. For the QSCI calculation of the T_1 (S_1) state, the input state(s) corresponding to the lower energy states, i.e., S_0 (S_0 and T_1) state(s), are prepared by converged sets of parameters of VQD. QSCI results are shown for the sequential diagonalization and the single diagonalization with two types of configuration selection, as described in the main text. The QSCI calculations with sequential diagonalization are done with $R_i = 16$ for $i = 0, 1, 2$, while the value of R is set to $R = 16$ for single diagonalization.

(see, e.g., Refs. [114,115] and references therein), while the hydrogen chains are also expected to show a clear scaling in the number of atoms. For Cr_2 , the correlation-consistent polarized valence quadruple-zeta (cc-pVQZ) basis set is used with n active orbitals and n active electrons with $n = 2, 4, \dots, 12$; the Jordan-Wigner mapping produces 4-, 8-, $\dots$, 24-qubit Hamiltonians, respectively. For the linear hydrogen chains, we consider 4, 6, $\dots$, 12 hydrogen atoms equally separated by a distance 1.0 Å; we use the STO-3G basis set without specifying the active space, corresponding to full-space Hamiltonians




FIG. 5. Estimated R required for a given energy error ϵ . Results with (a) expanding active spaces (Cr_2) or (b) various numbers of atoms (hydrogen chain) are shown by markers, along with the linear fit of each plot.

of 8-, 12-, ..., 24-qubit after the Jordan-Wigner mapping, respectively.

For each setup, the exact ground state of the Hamiltonian is prepared as the input state, and the QSCI calculation is performed by the idealized sampling introduced in Sec. II B, which picks up the R Slater determinants with the largest absolute values of CI coefficients in the input-state wavefunction. Then, for a given accuracy ϵ , the minimal R that satisfies $|E_R - E_{\text{exact}}| \leq \epsilon$ is determined, where E_R is the energy obtained by QSCI with the R configurations and E_{exact} by the exact diagonalization. In Fig. 5, the results are plotted for each molecule by varying the number of qubits, for $\epsilon = 0.1, 0.01$, and 0.001 hartree; they are extrapolated by fitting (shown by lines) to discuss the feasibility for larger system sizes.

As detailed in Sec. V A, we infer that the diagonalization with $R \simeq 5 \times 10^7$ configurations is achievable by the current state-of-the-art classical computing according to the reports [116,117]. The result for Cr_2 suggests that R is expected to

be manageable even when we require $\epsilon = 0.001$ hartree for a system larger than 50 qubits, where the exact diagonalization in the whole Fock space, i.e., CASCI, is challenging for classical computers. In the case of the hydrogen chains, on the other hand, the exponential growth of R is more clearly observed, and it may become hard to achieve $\epsilon = 0.001$ hartree for a system much larger than 50 qubits due to the limitation of classical computing. Note that the two scalings have slightly different meanings: The active space is enlarged for Cr_2 while fixing the molecule, i.e., the system size, while the system size itself is enlarged for the hydrogen chains. The results may suggest that our method is more suited to a localized system with many electrons involved, rather than a spatially extended system. For similar studies and results for several diatomic and aromatic molecules, see Appendix D 1.

We next estimate the number of shots for sampling required to achieve an error of ϵ by using the value of $1/|c_R|^2$ for each setup (Fig. 6). Here, c_R is the CI coefficient that has the R th largest absolute value in the input state, where R is taken to be the values shown in Fig. 5. When the state is sampled $1/|c_R|^2$ times, the probability of obtaining the R th most significant configuration is $O(1)$, and in that sense, $1/|c_R|^2$ gives a rough estimator for the number of shots required to sample R most significant configurations. We see in the next section, especially in Fig. 7, that this gives a ballpark estimate of the required number of shots for a given accuracy.

For comparison, the total number of shots required in a conventional expectation-value estimation is also estimated. More precisely, we analytically estimate the number of shots for which the standard deviation of the expectation-value estimations equals ϵ for the exact ground state (see, e.g., Ref. [31]). In the conventional methods, the expectation value of the Hamiltonian, which is expressed as a linear combination of Pauli strings, is estimated by directly measuring the quantum state in the basis of the Pauli strings multiple times and taking the average of the measurement outcome. To reduce the number of measurements, we employ the qubitwise commuting (QWC) grouping [110] with the sorted insertion algorithm [118]. The total shot is distributed to each of the groups with the shot allocation optimized for the exact ground state [119–121]. Note that, although there are methods that are capable of reducing the number of measurements better than QWC, they require more gate operations for measurements than QWC does; QWC requires a layer of single-qubit rotations after the state preparation, which is minimal for methods that measure the Pauli strings directly, while QSCI requires no gate operation. Most of the other grouping methods are thus expected to be more vulnerable to noise, and QWC is chosen for a fair comparison in this study.

Figure 6 shows the values of $1/|c_R|^2$ in QSCI for various numbers of qubits, along with the estimated numbers of shots in QWC. For the hydrogen chains, the results of QWC are fitted by a function $a(N_q)^b$ with parameters a and b as they are expected to be polynomial in the number of qubits [122], while the scaling of QSCI is unclear and fitting is not performed except in the case of Cr_2 with $\epsilon = 0.001$ hartree. In the case of Cr_2 , the number of shots for the proposed method seems to be consistently smaller than that of QWC, and by extrapolating the fitted results up to 50 qubits, one can see that the sampling cost is still within one day of execution time




FIG. 6. Estimated number of shots for a given energy error ϵ . For QSCI, the number of shots is approximated by $1/|c_R|^2$, where R for each ϵ is obtained in Fig. 5 and c_R is the CI coefficient of the input state with R th largest absolute value. The reasoning for this approximation is explained in the main text. For comparison, the results of the conventional expectation-value estimation with QWC grouping are plotted for each ϵ , fitted by a logarithmic function in the plot: The required numbers of shots for evaluating the energy are shown by dashed curves; for hydrogen chains, the ones for evaluating the gradients and Hessians in addition to the energy are shown by dash-dotted curves; the precision of gradients and Hessians are set to be $\epsilon/\text{\AA}$ and $\epsilon/\text{\AA}^2$, respectively. The number of shots that can be executed on Google's Sycamore processor [1], which can perform a million shots in about 200 s, is plotted in black dashed line. For panel (a), we also plot a linear fit of the QSCI results with $\epsilon = 0.001$ hartree, for reference.

by Google's Sycamore processor [1]. The result shows that, while the scaling may be exponential, QSCI can be advantageous at around 50 qubits, where the exact diagonalization is not possible but the quantum sampling cost is manageable. The advantage of QSCI, in terms of reducing the effect of statistical fluctuation, is more nontrivial in hydrogen chains with the number of qubits $N_q \gtrsim 30$.

The more operators are evaluated with the same output state, the more advantageous QSCI becomes; as we al-




FIG. 7. Energy error results for both QSCI and conventional QWC in sampling simulations. For each set of ten trials for each method, the standard deviation and the average of the absolute error to the exact value obtained by exact diagonalization are shown. QSCI energy error using $1/|c_R|^2$ shots obtained in Fig. 6 for $\epsilon = 0.1, 0.01, 0.001$ hartree is also plotted for reference. The horizontal line indicates the chemical accuracy, 1.6 mhartree.

ready noted in the previous section, QSCI does not require any additional quantum computation to evaluate additional observables, because QSCI outputs the classical vector representation of the state, and the expectation values are evaluated classically. On the other hand, in the conventional methods, quantum computational cost becomes more expensive, e.g., to measure additional Pauli strings introduced by the extra operators. To exploit this feature, we explore a scenario where the nuclear gradient and Hessian are evaluated along with the energy in the case of the hydrogen chains. For the shot allocation in the QWC grouping, we developed a method that is optimized for measuring multiple operators at once and is used in the simulation (see Appendix B 2 for details). The result, shown also in Fig. 6(b), implies that such a scenario makes QSCI much more advantageous [123], as the number of shots for QWC significantly increases.

QSCI generally outperforms QWC in terms of the sampling cost within the range of the system size that we studied. Although the scaling of QSCI seems to be worse than that of QWC in hydrogen chains, we should emphasize here that, even if QWC outperforms QSCI at, say, 50 qubits, it does not mean that QSCI is not useful for Hamiltonians with more than 50 qubits: QSCI has various features, such as error mitigation and the explicit representation of the output state, over the conventional methods, in addition to the reduction of the number of shots. The result should be interpreted as an implication that QSCI can be advantageous in moderately smaller but still classically challenging systems, even when we only consider the effect of reduction of the number of shots.

D. Sampling simulation

For assessing the effect of the statistical error during the sampling in QSCI, sampling simulation with different numbers of shots is performed. The result for a linear H_6 molecule

(12 qubits) is shown in Fig. 7, and results for other molecules are provided in Appendix D 2. For this simulation, the exact ground state is used as the input state, and we include all the configurations obtained in the sampling into the basis set $\mathcal{S}_R$ and we do not specify R beforehand. For comparison, we also performed a conventional sampling estimation for the exact ground state with QWC grouping and a shot allocation optimized for Haar random states.

For both QSCI and QWC, we perform ten trials of sampling simulation for each number of shots, and the average of the absolute differences to the exact ground-state energy is plotted along with the standard deviation of the ten trials. The absolute differences to the exact value are much smaller in QSCI compared to the conventional sampling with QWC grouping.

It is worth noting that the standard deviation of QSCI energy is smaller than its average difference, while that of QWC sampling is roughly equal. Energy values obtained by QSCI are biased estimators for the exact expectation values even when using the exact ground states as input states. Thus, the absolute difference can roughly be calculated as a sum of the intrinsic bias existing in QSCI and the standard deviation, which comes from statistical fluctuation of the subspace $\mathcal{S}_R$. In QWC, on the other hand, the statistical error is the only source of error. One can say that the QSCI result is much less affected by the statistical error compared to the conventional method.

Furthermore, as one can see in Fig. 7, $1/|c_R|^2$ calculated in the previous simulation gives a relatively accurate estimation of the total shots that gives an average error close to ϵ . Thus, the plots in Fig. 6 for both QWC and QSCI give reasonable estimations of the number of shots with expected average error ϵ , and the comparison is fair in this sense.

IV. BENCHMARK OF QSCI WITH NOISY SIMULATION AND EXPERIMENT

In this section, we describe the result of the experiment for the ground state of the hydrogen chain H_4 (eight qubits), conducted on the IonQ 11-qubit device through Amazon Braket service, along with the result of noisy sampling simulation using Qulacs with the identical setup. We first run a VQE calculation of a linear H_4 molecule with bond lengths 1.0 Å on a noiseless state-vector simulator. We use the STO-3G basis set without freezing any orbitals, and thus the problem Hamiltonian is eight-qubit. The so-called Ry ansatz with depth 8 is employed for the VQE calculation. See Appendix C 3 for details, including the circuit diagram of the ansatz. Then, we perform QSCI calculations on the quantum hardware and the noisy simulator using four sets of parameters at four distinct iterations of the VQE calculation. We use 10 000 shots for each sampling, and the most frequent R configurations are selected to define the subspace, with and without the postselection. The postselection of the sampling result is performed using the number of electrons $N_e = 4$ and the spin $S_z = 0$. For noisy simulation, to simulate the physical noises on the device, single-qubit depolarizing noise is added after each gate, and bit-flip noises are added at the end of the circuit to mimic the measurement error. The level of each type of noise is determined by the single-qubit and two-qubit gate fidelities,

and the measurement fidelity of the actual device: 99.61%, 96.868%, and 99.824%, respectively [124].

For comparison, on the quantum device and the noisy simulator, the calculation of the expectation value of the energy using a conventional method is performed with 10 000 shots. The QWC grouping and the shot allocation optimized for Haar random states are employed. Error-mitigation techniques, which may improve the result at the cost of additional quantum resources, are not employed in this study.

The results are presented in Fig. 8. By comparing the results from the noisy simulator and the quantum device, one can see that they have a reasonable agreement, although the result from the quantum device seems to be more affected by the errors. Moreover, it is clear that the postselection is powerful in both simulation and experiment. We also provide a comparison with classical CI methods such as CISD. It is particularly worth noting that even on the physical device, some of the QSCI calculations with $R = 27$ do outperform the result of CISD, which also diagonalizes the subspace Hamiltonian with 27 Slater determinants, and achieve the chemical accuracy for this eight-qubit system. The result of the configuration interaction perturbatively selected iteratively (CIPSI) [45] is also presented as more optimal result for selected CI. It is sufficient to use 16 Slater determinants to obtain an accuracy of 0.1 mhartree at the cost of multiple diagonalizations during the iterative selection process.

Some minor comments are in order: First, at the earlier iterations, the number of sampled (and postselected) configurations was sometimes less than the given R , because the state is concentrated in some computational basis states. In that case, we only used the sampled configurations for the QSCI calculation; second, CASCI result, i.e., the exact diagonalization result, corresponds to [125] $R = 36$; this number may seem to be comparable to $R = 27$, but it is still a nontrivial task to choose 27 configurations out of 36 possibilities.

V. DISCUSSION

In this section, we discuss various aspects of QSCI. We start from its classical and quantum computational costs in Sec. V A, and then discuss its benefits for refining VQE results in Sec. V B. In Sec. V C, several ideas for preparing input states are introduced. The aspect of QSCI as a selected CI is discussed in Sec. V D, and ideas for future directions are finally introduced.

A. Computational costs

Here, classical and quantum computational costs are examined. In QSCI, classical computing is used for generating the truncated Hamiltonian matrix $\mathbf{H}_R$ and diagonalizing it. Exploiting the Slater-Condon rules, one can generate the sparse matrix $\mathbf{H}_R$ efficiently in both R and the number of orbitals (see, e.g., Ref. [65] for details). For diagonalizing $\mathbf{H}_R$, one can employ algorithms to diagonalize a sparse matrix, such as the Lanczos method or the Davidson method. The generation and diagonalization of the Hamiltonian matrix are common procedures in the selected CI methods, and it is reported [117] that $R \simeq 5 \times 10^7$ of Slater determinants are manageable when a state-of-the-art high-performance computing resource




FIG. 8. QSCI results for the ground state of the linear hydrogen chain H_4 with eight qubits by the noisy simulator [panels (a) and (b)] and by the IonQ device [panels (c) and (d)], with and without postselection, compared with the conventional method of quantum-expectation-value estimation and classical computational methods. The resulting energies are plotted in hartree as deviations from the one obtained by the exact diagonalization (CASCI). The conventional method uses the QWC grouping in the energy estimation. Two classical methods are plotted for reference: CISD uses 27 Slater determinants, and we chose 16 Slater determinants for CIPSI, which is sufficient to achieve 0.1 mhartree accuracy. Lines specified by “VQE” show the exact energy value of the parametrized state at each iteration, and the states at four selected iterations are used as the input states for the QSCI calculations, shown by “QSCI” with four different values of R . The markers on the solid lines show the average value of some trials while each line without markers shows the result of each trial. The number of trials is ten for both QSCI and the conventional method on the noisy simulator, one for the conventional method, and five for QSCI, both on the IonQ device.

is available, even for the method that repeats the Hamiltonian generation and diagonalization. In our method, such a repetition is not needed, and thus the computational cost should be smaller. As already discussed in Sec. III C, Fig. 5 suggests that, for some challenging molecules of ~ 50 qubits, the QSCI calculation is feasible in terms of the classical cost by the current state-of-the-art classical computing, while meeting the accuracy requirement of $\epsilon \lesssim 0.001$ hartree. Note that such a system size would be beyond the reach of the exact diagonalization.

The quantum computational time is $t_Q = N_{\text{shot}} \times t_{\text{prepare}}$, where N_{shot} is the number of shots for the sampling, i.e., the repetitions of the input-state preparation and measurement, and t_{prepare} is the time needed for a single shot. Note that the total computational time can be reduced if multiple quantum computers are available, since the sampling procedures are completely parallelizable. t_{prepare} highly depends on the type of quantum device to be used and the way to prepare the input state. For example, the Sycamore processor used in the Google’s quantum supremacy experiment [1] can

achieve $N_{\text{shot}} = 1 \times 10^6$ in 200 s for a quantum circuit with 53 qubits and 20 repetitions of entangling operations, which corresponds to $N_{\text{shot}} \sim 4 \times 10^8$ in a day. Hence, Fig. 6 implies that the sampling cost is affordable for Cr_2 with several tens of qubits, while it may be challenging at the moment to achieve $\epsilon = 0.001$ hartree for a hydrogen chain with, say, 50 qubits.

We remark that the sampling cost can be significantly reduced if one can prepare a state $|\Delta\psi\rangle$ that is orthogonal to a classically tractable state $|\psi_c\rangle$ such that, for some complex numbers α and β , $|\psi_{\text{GS}}\rangle = \alpha|\psi_c\rangle + \beta|\Delta\psi\rangle$ approximates the ground state, and can sample from $|\Delta\psi\rangle$ on a quantum computer. The state $|\psi_c\rangle$ can be the Hartree-Fock state or more intricate states such as the CISD state. For example, if $|\alpha|^2 = 0.9$, then the sampling cost for a given precision can be reduced by a factor of 10. On the other hand, $|\Delta\psi\rangle$ can be prepared, e.g., by the method of Ref. [87].

In this paper, we have assumed that an input state that approximates the true eigenstate is given. Clearly, the end-to-end computational time critically depends on the method of state preparation such as VQE or other alternatives. If the

preparation method calculates the circuit parameter for the input state beforehand, e.g., by VQE, then there would be an additional cost, apart from the sampling cost for QSCI. If the state preparation method itself involves deep circuit such as in the adiabatic state preparation, then each circuit execution during the sampling task can take significantly longer time due to non-negligible gate-operation time. In this sense, the comparison with CISD is fair only in the limit where all such costs are regarded to be negligible. The comparison with CIPSI, on the other hand, is more nontrivial. CIPSI repeats the diagonalization with iteratively selected subspaces, while the diagonalization is executed only once in QSCI. Thus, in comparison to QSCI, CIPSI requires more classical computation and less (no) quantum computation. Investigating whether the total end-to-end computational time, including both quantum and classical, of a practical QSCI implementation can be reduced below that of CIPSI remains an interesting open question.

B. Use of QSCI to refine VQE results

QSCI can be viewed as a postprocessing technique for VQE and its variants when they are used to prepare the input states. Our methods have the following advantages:

Error reduction. By virtue of the classical diagonalization of a Hamiltonian matrix generated classically, the proposed methods can refine the VQE results, as demonstrated in Secs. III and IV. Although results of noiseless VQE simulations are used to prepare the input states in our numerical and experimental studies, our results suggest that QSCI is also effective to refine *dirty* VQE results subject to the statistical and physical errors. See Appendix D 6 for simulations with noisy QSCI using noisy VQE results. Figure 8 also shows the effectiveness of the pos-selection: The rate of the readout error, which is one of the major sources of physical errors, can be reduced from $O(p)$ to $O(p^2)$ with the Jordan-Wigner mapping, as discussed in Appendix A. Note that QSCI does not require extra gate operations for the measurement, unlike expectation-value estimations in VQE. As already shown in Fig. 3, our method is also effective to improve the quality of the input state even in the absence of physical and statistical errors. This feature may enable one to use ansatz with shallower circuits, or to reduce the number of optimization steps in VQE, by employing QSCI to improve the final result.

Reliability. Our method is free of errors in the sense that the resulting ground-state energy is exact within the subspace spanned by the quantum-selected configurations. This means that the obtained energy is a definite upper bound for the exact ground-state energy, which is not the case in conventional VQE because of physical and statistical errors, as discussed in Sec. II B. This is advantageous for comparing the QSCI result obtained on noisy quantum devices with the results of classical variational methods such as CISD or density matrix renormalization group (DMRG) [126–128]: The variational nature of these methods guarantees that the method that gives the lowest energy is the most accurate one. Similar variational inequalities hold for excited states in the single diagonalization scheme of QSCI, while there is no such guarantee in the sequential diagonalization scheme.

Although the latter appeared to be more accurate in our numerical simulation, the former is of great use if one is interested in giving rigorous upper bounds on excited-state energies.

Handiness. As one has the classical representations of the eigenstates as output, one can compute the expectation values of a large class of observables with no additional quantum computation. Our method becomes more valuable when more observables are to be evaluated, as exemplified in Fig. 6. Moreover, one can also analyze the classical vectors themselves, which may be useful to study the significance of each Slater determinant.

C. Use of QSCI with more general input states

As discussed in the previous sections, input states for ground state can be prepared by VQE, and those for excited states by its variants, but the proposed methods are applicable to more general input states.

Our method can in principle be applied to any kind of input states that can be prepared and sampled on a quantum computer. We give an incomplete list of possible preparation schemes for input states in the following: the adiabatic state preparation [129,130], the imaginary time evolution [131–134], classically boosted VQE [87], classically optimized shallow ansatz circuits [135], unitary coupled-cluster ansatz circuits with classically optimized parameters [110,136–138], and parametrized states classically optimized by Clifford circuits [139,140]. Note that the performance of QSCI depends on the quality of the input state and also on the form of the exact eigenstate. For example, if the exact eigenstate is the equal superposition of all the computational basis states, then our algorithm will not perform well. Note that, from a computational complexity perspective, QSCI is expected to be efficient only when the number of important configurations scales polynomially with the number of qubits. We also remark that the sampling cost required to collect important configurations from the input states must be taken into account when considering the efficiency of the proposed method, even though the input states contain information about the important configurations. The numerical results in Fig. 6 show that such a cost can be manageable in practice, at least for some of our investigated cases with practical problem sizes around 50 qubits.

The algorithm can also be useful for a Hamiltonian that has an exactly known ground state. For example, one can calculate an exact ground-state energy of a system that is solvable with the Bethe ansatz, but there are quantities (e.g., a class of correlation functions) that cannot be computed efficiently [141]. Our method provides the classical representation of an approximate eigenstate, which means that one can evaluate various physical quantities without additional quantum resource, as we already discussed for states prepared by VQE. The preparation of the Bethe ansatz states on quantum computers is addressed in Refs. [142,143].

Moreover, although we proposed the method as a hybrid quantum-classical algorithm, one can apply the method to input states that can be sampled efficiently on classical computers. This is shortly discussed in Sec. V E.

D. QSCI as selected CI

As selected CI methods, the novelty of QSCI comes simply from how to define the subspace on which we construct the subspace Hamiltonian. Quantum computers are used to sample important configurations from the input state, and there is a quantum speedup when the input state is hard to sample classically. In selected CI methods, the subspace of the Fock space for the diagonalization is either fixed by the method, e.g., CISD, or adaptively chosen according to the algorithm. We have shown experimentally that CISD performs worse even when compared to the QSCI result on the current NISQ device (Fig. 8).

One of the most advanced methods for sampling dynamically important bases is the adaptive sampling configuration interaction (ASCI) algorithm developed by Tubman and co-workers [57,65]. The idea of systematically selecting important bases based on perturbation theory was developed about 50 years ago [43–47], and a selection scheme based on Monte Carlo methods was proposed in the 1990s [51,52]. However, systematically selected CI was not widely used in quantum chemistry calculations for many years. Recently, it has undergone rapid development and is now becoming applicable to large-scale quantum chemical simulations [53–74]. Indeed, Tubman *et al.* showed that ASCI is capable of handling 34 electrons in 152 spatial orbitals [65].

ASCI has hyperparameters that define the size of the search space to adaptively select the configurations while QSCI does not have such hyperparameters and the configurations can be selected by sampling a single input state on a quantum computer with no iterative process. We will see in Appendix D 5 that, with some set of hyperparameters, QSCI can perform better than ASCI in selecting important configurations to achieve accurate ground-state energies, though we feed the exact ground state for QSCI in contrast to ASCI, which does not require any knowledge of the ground state itself.

E. Outlook

QSCI is applicable to diverse systems and has many directions for generalizations.

(1) It would be possible to consider a hybrid of the proposed method and another adaptive selected CI method, such as ASCI, by combining the configurations suggested by QSCI with those of the other method. In this way, one could improve the results of the state-of-the-art selected CI methods by using quantum computers.

(2) QSCI is essentially a selected CI where the configurations are randomly selected according to a probability distribution $p(x) = |\langle x | \psi_{\text{in}} \rangle|^2$. A classical counterpart of this approach is called Monte Carlo configuration interaction (MCCI) [51,52]. MCCI does not seem to have been extensively studied since the first proposal in 1995, and the use of a more sophisticated (classical) probability distribution for MCCI is yet to be explored. It would be an interesting future work to use a classically tractable $p(x)$ for MCCI and compare/combine it with QSCI. For example, some of the tensor network states, such as the matrix product states and the multiscale entanglement renormalization ansatz states can be efficiently sampled on classical computers [144]. We note that the random sampling of configurations in a Hilbert space

itself is popular among various classical methods in quantum chemistry such as full configuration interaction quantum Monte Carlo [145,146] and semistochastic projection Monte Carlo [147], where one can explore efficiently Hilbert spaces with enormous dimensions (e.g., $\sim 10^{35}$).

(3) Our method, compared to the conventional VQE, has an advantage that it can evaluate physical observables classically with no additional quantum computational cost. One may leverage this feature by using QSCI for the geometry optimization problem of a molecule or a molecular dynamics calculation. In those applications, one may skip the sampling for some iterations and continue to run with the same state subspace defined by the R electron configurations, thereby reducing the quantum computational cost further.

We remark that the performance of QSCI depends highly on the quality of the input state. It would be great if there is a way to start from an input state with modest quality, and then improve the quality of the input state by an iterative use of QSCI.

VI. CONCLUSION

In this work, we proposed QSCI, a class of hybrid quantum-classical algorithms, to find low-lying eigenvalues and eigenstates of a many-electron Hamiltonian. Taking approximated states of such eigenstates as input, QSCI selects important electron configurations to represent the eigenstates by sampling the input states on quantum computers, and then classically diagonalizes the Hamiltonian in the subspace(s) spanned by the selected configurations to yield better approximations for the eigenstates and their energies. QSCI is robust against noise and statistical fluctuation, as quantum computation is used only to define the subspaces. A quantum speedup potentially arises in that sampling a quantum state is, in general, classically intractable.

We verified the algorithms for ground and excited states of small molecules by numerical simulations and experiment, where the latter was conducted on the quantum device with the eight-qubit quantum circuits. We discussed potential utility of QSCI in various aspects: For instance, taking a state obtained by VQE as the input state, QSCI can be used to refine the VQE result, which may not be accurate enough due to statistical fluctuation, physical noise, and poor optimization. QSCI is particularly advantageous for estimating expectation values of multiple observables with respect to energy eigenstates, as additional observables can be evaluated without extra quantum computational cost. We also argued that QSCI is potentially feasible to tackle challenging molecules such as the chromium dimer by exploiting quantum devices with several tens of qubits, assisted by a high-performance classical computing resource for diagonalization.

ACKNOWLEDGMENTS

The authors thank Shoichiro Tsutsui for a careful reading and comments on the revised manuscript. The authors thank Amazon Web Services for supporting this work through their Amazon Braket service. A part of this work was supported by JST PRESTO JPMJPR2019 and JPMJPR191A, MEXT Quantum Leap Flagship Program (MEXTQLEAP) Grants

No. JPMXS0118067394 and No. JPMXS0120319794, and JST COI-NEXT program Grant No. JPMJPF2014.

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon reasonable request.

APPENDIX A: THE EFFECT OF POSTSELECTION

In this Appendix, we discuss the effect of postselection to mitigate errors in QSCI. To be specific, we consider the postselection technique introduced in Sec. II B, which exploits the conservation of particle number and spin, targeting the bit-flip noise. In the following, we consider to measure an N -qubit computational basis state. We assume that the state is prepared without the effects of noise but each bit of the measurement result is flipped with error probability p . It is equivalent to the situation where the bit-flip noise is introduced to each qubit independently after the input state is generated. The probability that the N -bit string describing the state is measured correctly is $(1 - p)^N$.

1. Jordan-Wigner mapping

Let us now assume that we consider an electronic Hamiltonian converted by the Jordan-Wigner mapping. In this case, the number of 1s in the N -bit string, which we denote by n_1 , corresponds to the number of electrons in the system and is sometimes known prior to the calculation for a ground state or an excited state. One can thus perform the postselection for a measurement outcome that excludes resulting bit strings with the number of 1s not equal to n_1 . Although one may still get incorrect results, the probability is reduced.

More concretely, the probability to get a result with correct n_1 is

$$(1 - p)^N + n_0 n_1 p^2 (1 - p)^{N-2} + O(p^4), \quad (\text{A1})$$

where we define $n_0 := N - n_1$. After the postselection, the probability to get the correct result is thus

$$\frac{(1 - p)^N}{(1 - p)^N + n_0 n_1 p^2 (1 - p)^{N-2} + O(p^4)} \quad (\text{A2})$$

$$= \frac{1}{1 + n_0 n_1 p^2 / (1 - p)^2 + O(p^4)} \quad (\text{A3})$$

$$= \frac{1}{1 + n_0 n_1 p^2 + O(p^3)} \quad (\text{A4})$$

$$= 1 - n_0 n_1 p^2 + O(p^3). \quad (\text{A5})$$

The error rate of getting an incorrect result is reduced from $1 - (1 - p)^N \sim pN$ to $n_0 n_1 p^2$ with the ratio being

$$\frac{n_0 n_1 p^2}{pN} = \left(\frac{n_0}{N}\right) \left(\frac{n_1}{N}\right) Np \sim O(pN), \quad (\text{A6})$$

which is less than one in a sensible situation $Np \ll 1$, where the original success probability $(1 - p)^N$ is not too small.

Although we here considered a computational basis state as the input state, we expect that the postselection similarly works for a general input state that is a superposition of computational basis states with some fixed n_1 . Note that, if

one also knows the total spin S_z of electrons prior to the calculation, one can count the number of 1s separately for up- and down-spin electrons, and make the postselection more efficient.

2. Other mappings

For most of the other fermion-qubit mappings, it is not expected that the reduction of the error probability from $O(p)$ to $O(p^2)$ happens. For example, in the parity mapping [89,148] and the Bravyi-Kitaev mapping [89], the states $|01\rangle$ and $|11\rangle$ are connected by just one bit flip, but both of them are one-electron states. This bit flip, which cannot be detected by the postselection, occurs with probability $O(p)$, and thus the error rate after the postselection is still $O(p)$. The same is true for a Hamiltonian with a reduced number of qubits using symmetries, where there is always a bit flip that does not change the total number of electrons.

APPENDIX B: DETAILS OF THE ALGORITHMS

In this Appendix, we present several detailed discussions on the QSCI algorithms.

1. Choice of β_i parameters and variational inequalities in sequential diagonalization scheme

Here, we discuss the sequential diagonalization scheme, introduced in Sec. II C 2, on how to choose the β_i parameters and a potential violation of the variational inequality, following the discussion in Ref. [97].

Suppose k low-lying eigenstates of $\hat{H}$, $|E_0\rangle, \dots, |E_{k-1}\rangle$, are known exactly. Then, the effective Hamiltonian to find the k th eigenstate can be exactly constructed as

$$\hat{H}^{(k')} = \hat{H} + \sum_{i=0}^{k-1} \beta_i |E_i\rangle \langle E_i|. \quad (\text{B1})$$

This can be formally expressed as

$$\hat{H}^{(k')} = \sum_{i=0}^{k-1} (E_i + \beta_i) |E_i\rangle \langle E_i| + \sum_{i \geq k} E_i |E_i\rangle \langle E_i|, \quad (\text{B2})$$

where E_i represents the i th eigenvalue of $\hat{H}$ in this Appendix. For $\beta_i > E_k - E_i$ ($i = 0, \dots, k-1$), the following inequality holds for an arbitrary $|\psi\rangle$ with $\langle \psi | \psi \rangle = 1$:

$$\langle \psi | \hat{H}^{(k')} | \psi \rangle \geq E_k, \quad (\text{B3})$$

where the equality holds if and only if $|\psi\rangle = |E_k\rangle$ up to a phase factor. In the language of the eigenvalue problem of Eq. (12), this implies $E_{R_k}^{(k')} \geq E_k$, where $E_{R_k}^{(k')}$ is the smallest eigenvalue of $\mathbf{H}_{R_k}^{(k')}$, the subspace matrix for $\hat{H}^{(k')}$, defined in the same way as Eq. (13).

In practice, the condition $\beta_i > E_k - E_i$ can be utilized if one has prior knowledge of the energy spectrum, e.g., based on variational quantum algorithms. However, even without such knowledge, one may still rely on the stronger condition of $\beta_i > 2 \sum_j |c_j|$ [97], which is written in terms of the coefficients c_j of the qubit Hamiltonian $\hat{H} = \sum_j c_j P_j$, expressed by the Pauli strings P_j .

In reality, the effective Hamiltonian cannot be exactly constructed as the k low-lying eigenstates would be obtained only approximately, and, hence, the inequality $E_{R_k}^{(k)} \geq E_k$ is not guaranteed. For instance, in the problem to find the first excited state, the effective Hamiltonian $\hat{H}^{(1)}$ is constructed with $|\psi_{\text{out}}^{(0)}\rangle$, the output state for the ground state obtained by the preceding step in sequential diagonalization. Unless the output state perfectly overlaps with the true ground state $|E_0\rangle$, or $|\langle\psi_{\text{out}}^{(0)}|E_0\rangle| = 1$, there is no guarantee that $\langle\psi|\hat{H}^{(1)}|\psi\rangle$ is bounded by the exact eigenvalue E_1 . Instead, $\min_{\psi}\langle\psi|\hat{H}^{(1)}|\psi\rangle$ is only bounded as [97]

$$E_1 - O((E_1 - E_0)\epsilon_0) \leq \min_{\psi}\langle\psi|\hat{H}^{(1)}|\psi\rangle \leq E_1 + \beta_0\epsilon_0, \quad (\text{B4})$$

where $\epsilon_0 = 1 - |\langle\psi_{\text{out}}^{(0)}|E_0\rangle|^2$ and $\langle\psi|\psi\rangle = 1$. A concrete example for breaching the variational inequality is given as follows. Consider a system with the unique ground state, i.e., $E_0 < E_1$. Suppose that one has a poor output state $|\psi_{\text{out}}^{(0)}\rangle$ that is orthogonal to the true ground state $|E_0\rangle$, i.e., $\epsilon_0 = 1$. Then, $\langle\psi|\hat{H}^{(1)}|\psi\rangle \geq E_0$ for any positive β_0 , where the equality holds if and only if $|\psi\rangle = |E_0\rangle$ up to a phase factor. This means that the variational inequality $E_{R_1}^{(1)} \geq E_1 (> E_0)$ is violated at least in the limit where the subspace $\mathcal{S}_{R_1}^{(1)}$ is enlarged to cover the part of the Fock space necessary to express $|E_0\rangle$. Such a subspace can be constructed, e.g., if the input state is chosen to be $|\psi_{\text{in}}^{(1)}\rangle = |E_0\rangle$ with a sufficiently large R_1 .

2. An optimal shot allocation for evaluating expectation values of multiple observables in conventional method

In this subsection, we describe the details of the numerical estimation of computational cost in Sec. III C for evaluating multiple operators. In the numerical simulation, we considered a situation where we want to calculate the expectation values of the nuclear gradient $\{\frac{\partial\hat{H}}{\partial x_i} \mid i = 1, \dots, 3N_{\text{atom}}\}$ and the nuclear Hessian $\{\frac{\partial^2\hat{H}}{\partial x_i \partial x_j} \mid i, j = 1, \dots, 3N_{\text{atom}}\}$ along with the Hamiltonian $\hat{H}(\{x_i\})$, where x_i are the nuclear coordinates and N_{atom} is the number of atoms in the molecule. The most naive way of doing it would be to calculate each expectation value completely separately. This is, though, too naive to be considered as the optimal strategy; all the observables are linear combinations of operators $a_i^\dagger a_j$ and $a_i^\dagger a_j a_k^\dagger a_l$ in the fermionic basis, and the expectation values of these operators can be reused among the operators.

Before discussing the optimal strategy for evaluating expectation values of multiple observables, let us review the one for a single observable, following the discussion in Ref. [121]. Consider a quantum state $|\psi\rangle$ and the expectation value of an operator $\hat{O}$, which can be written as a sum of operators $\hat{O}_l$:

$$\hat{O} = \sum_{l=1}^L \hat{O}_l. \quad (\text{B5})$$

Each term $\hat{O}_l$ can be either a Pauli string or a sum of Pauli strings that commute with each other, which admits the projective measurement on eigenvalues of each $\hat{O}_l$. We denote the variance of each term $\hat{O}_l$ per one shot by $\sigma_l^2 := \text{Var}(\hat{O}_l) := \langle\psi|\hat{O}_l^2|\psi\rangle - \langle\psi|\hat{O}_l|\psi\rangle^2$. By measuring each term $\hat{O}_l$ with

a number of shots M_l , the observed expectation value has the variance $\sum_l \sigma_l^2 / M_l$. Employing the method of Lagrange multiplier with the Lagrangian

$$\mathcal{L} = \sum_l M_l + \lambda \left(\sum_l \frac{\sigma_l^2}{M_l} - \epsilon^2 \right), \quad (\text{B6})$$

one can get the optimal allocation of the number of shots with the total variance of the expectation value fixed to ϵ^2 , which is

$$M_l \propto \sigma_l. \quad (\text{B7})$$

In general, σ_l is not exactly known *a priori*, so one may use σ_l for Haar random states to get a reasonable strategy. One may also try to improve the strategy by dividing the shot budget for one evaluation of an expectation value into several iterations: One can simply evaluate the expectation value with a mildly optimized strategy in the first iteration, and then, in the rest of the iterations, one can adjust the strategy by calculating σ_l by using the expectation values obtained in the previous iterations.

Generalizing the above discussion, let us consider a situation where one calculates the expectation values of a set of operators $\{\hat{O}^{(i)} \mid i = 1, \dots, n\}$. We assume that $\hat{O}^{(i)}$ is decomposed as

$$\hat{O}^{(i)} = \sum_{l=1}^L \hat{O}_l^{(i)}, \quad (\text{B8})$$

where all of $\{\hat{O}_l^{(i)} \mid i = 1, \dots, n\}$ are simultaneously measurable for each l , i.e., $[\hat{O}_l^{(i)}, \hat{O}_l^{(j)}] = 0$ for any i, j . In our numerical simulation, the grouping was done by first taking the sum of all the observables $\hat{O}^{(i)}$ with each Pauli string with negative coefficient multiplied by -1 to make it positive. Then, the greedy qubitwise grouping of Refs. [110,118] was used. Our aim here is to find a good strategy to estimate the expectation values of all the operators $\hat{O}^{(i)}$ with statistical error less than ϵ . Note that one can always rescale the observables so that the required precision is the same for all observables even when one requires different precision for different operators. To get an analytical solution, we choose the following Lagrangian with slightly modified constraint:

$$\mathcal{L} = \sum_l M_l + \lambda \left(\sum_l \sum_i \left(\frac{\sigma_l^{(i)2}}{M_l} \right) - \epsilon_{\text{tot}} \right), \quad (\text{B9})$$

where ϵ_{tot} can be $N \times \epsilon$ but it turns out that the choice of ϵ_{tot} does not affect the final result. By solving the extremal condition of this Lagrangian, one can get the best shot allocation that minimizes the total number of shots, while keeping the sum of the variances of all the operators less than ϵ_{tot} . The result implies that

$$M_l \propto \sqrt{\sum_i \sigma_l^{(i)2}}. \quad (\text{B10})$$

By estimating the variance of each operator $\hat{O}^{(i)}$ with this shot allocation, and by adjusting the total number of shots so that the statistical error of each operator is ϵ at worst, one can obtain the total number of shots $\sum_l M_l$ with desired precision for

TABLE I. Geometries of molecules. “(X, (x, y, z))” denotes three-dimensional coordinates x, y, z of an atom X in units of Å.

Molecule	Geometry
H ₂ O	(O, (0, 0, 0)), (H, (0.2774, 0.8929, 0.2544)), (H, (0.6068, -0.2383, -0.7169))
H _n (n = 4, 6, 8, 10, 12)	(H, (0, 0, 0)), (H, (0, 0, 1.0)), ..., (H, (0, 0, n × 1.0))
LiH	(Li, (0, 0, 0)), (H, (0, 0, 1.595))
N ₂	(N, (0, 0, 0)), (N, (0, 0, 1.1))
O ₂	(O, (0, 0, 0)), (O, (0, 0, 1.2))
F ₂	(F, (0, 0, 0)), (F, (0, 0, 1.4))
Cl ₂	(Cl, (0, 0, 0)), (Cl, (0, 0, 2.0))
HCl	(H, (0, 0, 0)), (Cl, (0, 0, 1.3))
CO	(C, (0, 0, 0)), (O, (0, 0, 1.1))
Cr ₂	(Cr, (0, 0, 0)), (Cr, (0, 0, 1.6))
Benzene	(C, (-1.2131, -0.6884, 0)), (C, (-1.2028, 0.7064, 0.0001)), (C, (-0.0103, -1.3948, 0)), (C, (0.0104, 1.3948, -0.0001)), (C, (1.2028, -0.7063, 0)), (C, (1.2131, 0.6884, 0)), (H, (-2.1577, -1.2244, 0)), (H, (-2.1393, 1.2564, 0.0001)), (H, (-0.0184, -2.4809, -0.0001)), (H, (0.0184, 2.4808, 0)), (H, (2.1394, -1.2563, 0.0001)), (H, (2.1577, 1.2245, 0))
Naphthalene	(C, (0, -0.7076, 0)), (C, (0, 0.7076, 0.0001)), (C, (1.225, -1.3944, 0.0001)), (C, (1.225, 1.3944, 0)), (C, (-1.225, -1.3943, 0)), (C, (-1.225, 1.3943, 0)), (C, (2.4327, -0.6958, 0)), (C, (2.4327, 0.6959, -0.0001)), (C, (-2.4327, -0.6958, -0.0001)), (C, (-2.4327, 0.6958, 0)), (H, (1.2489, -2.4822, 0.0001)), (H, (1.2489, 2.4821, -0.0001)), (H, (-1.2489, -2.4822, -0.0001)), (H, (-1.249, 2.4821, 0.0001)), (H, (3.3733, -1.239, -0.0001)), (H, (3.3732, 1.2391, -0.0001)), (H, (-3.3733, -1.239, -0.0001)), (H, (-3.3732, 1.239, 0))
Anthracene	(C, (-1.225, 0.706, 0.0001)), (C, (-1.2251, -0.7061, 0.0001)), (C, (1.2251, 0.7061, 0.0002)), (C, (1.2251, -0.7061, 0.0001)), (C, (0, 1.3937, 0.0001)), (C, (0, -1.3938, 0)), (C, (-2.4504, 1.393, -0.0001)), (C, (-2.4505, -1.393, 0)), (C, (2.4505, 1.3929, 0)), (C, (2.4505, -1.3929, 0)), (C, (-3.6587, 0.6956, -0.0001)), (C, (-3.6588, -0.6955, -0.0001)), (C, (3.6587, 0.6956, -0.0002)), (C, (3.6587, -0.6956, -0.0002)), (H, (0, 2.4838, 0)), (H, (0, -2.4839, -0.0001)), (H, (-2.4742, 2.4808, -0.0001)), (H, (-2.4744, -2.4809, 0)), (H, (2.4742, 2.4808, 0)), (H, (2.4743, -2.4808, 0)), (H, (-4.5989, 1.2394, -0.0003)), (H, (-4.5991, -1.2391, -0.0002)), (H, (4.5989, 1.2393, -0.0003)), (H, (4.5989, -1.2393, -0.0004))
Tetracene	(C, (0, 0.7045, -0.0002)), (C, (0, -0.7046, -0.0001)), (C, (-2.451, 0.7058, 0)), (C, (-2.4511, -0.7058, 0.0002)), (C, (2.4511, 0.7057, 0.0001)), (C, (2.4511, -0.7058, -0.0001)), (C, (1.2254, 1.3923, -0.0001)), (C, (1.2254, -1.3924, -0.0003)), (C, (-1.2254, 1.3923, -0.0002)), (C, (-1.2255, -1.3923, 0.0002)), (C, (-3.6764, 1.3928, -0.0001)), (C, (-3.6764, -1.3929, 0.0002)), (C, (3.6764, 1.3929, 0.0003)), (C, (3.6765, -1.3929, -0.0001)), (C, (-4.8846, 0.6957, -0.0001)), (C, (-4.8847, -0.6955, 0.0001)), (C, (4.8846, 0.6957, 0.0004)), (C, (4.8847, -0.6956, -0.0001)), (H, (1.2253, 2.4825, -0.0001)), (H, (1.2254, -2.4825, -0.0003)), (H, (-1.2254, 2.4824, -0.0003)), (H, (-1.2255, -2.4824, 0.0003)), (H, (-3.6999, 2.4807, -0.0002)), (H, (-3.7001, -2.4808, 0.0003)), (H, (3.6999, 2.4807, 0.0004)), (H, (3.7001, -2.4807, -0.0003)), (H, (-5.8248, 1.2393, -0.0002)), (H, (-5.8249, -1.2392, 0.0002)), (H, (5.8248, 1.2394, 0.0005)), (H, (5.8249, -1.2392, -0.0002))

all the operators. This may not be the optimal shot allocation to achieve the statistical error ϵ for each operator as we are minimizing the total variance rather than the maximum value of the variances, but this will give a reasonable strategy that is analytically available.

There is one comment to make on the evaluation of nuclear Hessians. In the following, the expectation value is always taken by an ansatz state $|\psi(\theta(\{x_i\}))\rangle$ parametrized by the ansatz parameters $\theta(\{x_i\})$; we assume that $\theta(\{x_i\})$ is optimized so that $|\psi(\theta(\{x_i\}))\rangle$ has the minimum energy within the ansatz for each $\{x_i\}$. We denote the energy expectation value of the state by $E(\{x_i\})$. In the case of the nuclear gradient,

$$\frac{\partial E(\{x_i\})}{\partial x_i} = \left\langle \frac{\partial \hat{H}(\{x_i\})}{\partial x_i} \right\rangle \quad (\text{B11})$$

holds thanks to the Hellmann-Feynman theorem, and it suffices to compute the right-hand side to obtain the nuclear gradient of the energy. In the case of the nuclear Hessian of the energy [149], on the other hand, it is in general necessary

to evaluate the contribution of derivatives acting on the state as well as on the Hamiltonian operator,

$$\frac{\partial E(\{x_i\})}{\partial x_i} x_j = \left\langle \frac{\partial \hat{H}(\{x_i\})}{\partial x_i} x_j \right\rangle + \dots \quad (\text{B12})$$

In our numerical simulation, we ignored the contribution of the derivatives acting on the state for simplicity. In the case of QWC, this contribution requires additional quantum resources. On the other hand, in the case of QSCI, one can generate and diagonalize the Hamiltonians at small finite distance $x_i \rightarrow x_i \pm \delta$ to get the derivatives of the state within the same selected subspace of the Fock space with no additional quantum resources. If we take this contribution into account properly, the advantage of QSCI will increase.

It should also be noted that, although we evaluate $O(N_{\text{atom}}^2)$ observables in the numerical simulations for the hydrogen chain in the main text, due to the rich geometrical symmetry of the molecules, many of the observables are zero as an operator. It is likely that, for more generic molecules, the




FIG. 9. Real-valued symmetry-preserving ansatz with n qubits and depth d .

crossing point of QSCI and QWC comes at a larger number of qubits.

APPENDIX C: DETAILS OF NUMERICAL SIMULATIONS AND EXPERIMENTS

In this Appendix, we explain details of the numerical simulations and the experiment on quantum hardware in the main text. For all the molecules examined in this study, the second-quantized electronic Hamiltonian under the Born-Oppenheimer approximation is generated by OPENFERMION [103] interfaced with PYSCF [104] using the Hartree-Fock orbitals with the STO-3G minimal basis set, unless otherwise stated. The electronic Hamiltonians are mapped to qubit ones by the Jordan-Wigner transformation. The molecular geometries used in our study are shown in Table I. Stable geometries for diatomic molecules are taken from Computational Chemistry Comparison and Benchmark DataBase (CCCBDB) database [150] and Ref. [151], while those for the other molecules are taken from PUBCHEM [152], except for the hydrogen chains that are not in their stable geometries. We list the details specific to each of simulations and experiments in the following.

1. Noiseless simulation for ground state

In Sec. III A, the H_2O molecule with six active electrons and five active orbitals, is chosen to find the ground state by QSCI. In the VQE calculation for preparing the input states, the BFGS optimizer is employed through the scientific library SCIPY [107], and the real-valued symmetry-preserving ansatz [106] is used to construct parametric quantum circuits with depth 10 (Fig. 9). The initial state of the ansatz circuits is set to be the Hartree-Fock state, and the initial parameters in the optimization are randomly chosen.

2. Noiseless simulations for excited states

In Sec. III B, QSCI is demonstrated for the same H_2O molecule but to find excited states. To prepare the input states, the VQD calculations are performed in the same setup as the previous VQE calculation, but with the penalty terms




FIG. 10. Ry ansatz with eight qubits. All the rotational gates have independent parameters. The depth is set to be 8 in our experiment.

[110–112] added to the Hamiltonian for constraining the resulting states to have $S_z = 0$ and $N_e = 6$; specifically, the following operator (in atomic units) is added to the Hamiltonian:

$$3.0(\hat{S}_z)^2 + 3.0(\hat{N}_e - 6)^2, \quad (\text{C1})$$

where $\hat{S}_z$ is the operator for the total electron spin in z direction and $\hat{N}_e$ for the particle number operator of electrons in the active space. Furthermore, the overlap terms to constrain the state to be orthogonal to lower energy eigenstates [97] are added with coefficients of unity (in hartree). For the sequential diagonalization scheme of QSCI, the coefficients β_i for ensuring orthogonality are also set to unity.

3. Noisy simulation and experiment

For the noisy simulation and experiment in Sec. IV, the input states are prepared by noiseless VQE simulations. The VQE calculations are performed with the BFGS optimizer and Ry ansatz (Fig. 10) with depth 8. Other details are described in the main text.

APPENDIX D: SUPPLEMENTAL NUMERICAL RESULTS

In this Appendix, we provide additional numerical results to supplement the contents in Sec. III.

1. Scaling of computational costs with various molecules

Figure 11 shows the scaling of the classical and quantum computational costs, discussed in Sec. III C, for different types of molecules. Here, we test three kinds of molecules: hydrogen chains, diatomic molecules, and aromatic molecules. The data for hydrogen chains are exactly the same as in the main text. Diatomic molecules are N_2 , O_2 , F_2 , Cl_2 , HCl ,







(a) R required for energy error $\epsilon = 0.001$ Hartree.(b) $1/|c_R|^2$ for energy error $\epsilon = 0.001$ Hartree.(c) R required for energy error $\epsilon = 0.01$ Hartree.(d) $1/|c_R|^2$ for energy error $\epsilon = 0.01$ Hartree.(e) R required for energy error $\epsilon = 0.1$ Hartree.(f) $1/|c_R|^2$ for energy error $\epsilon = 0.1$ Hartree.

FIG. 11. Estimated R and $1/|c_R|^2$ for various molecules with the same setup as Figs. 5 and 6. The number of qubits is varied by changing the number of atoms for hydrogen chains, and by changing the active space for the other molecules.

CO, and Cr_2 with cc-pVQZ basis, and we tested them with various active spaces just as in the main text for Cr_2 . To test with larger molecules, four aromatic molecules are chosen: benzene, naphthalene, anthracene, and tetracene. The Hamiltonian is generated by using the Hartree-Fock orbitals with

STO-3G basis. The active space of n orbitals and n electrons with varying n was employed for the diatomic and aromatic molecules. The geometries of these tested molecules are summarized in Table I.




FIG. 12. Sampling simulation with various molecules. See Fig. 7 and the main text for details.

As can be seen in Fig. 11, hydrogen chains with various numbers of atoms show the worst scalings, while Cr_2 is one of the least expensive systems among others.

2. Sampling simulations with various molecules

In this subsection, we present similar results as Fig. 7 but with various other molecules. The results, shown in Fig. 12, show the same features as H_6 in the main text, such as the

small standard deviation for QSCI and $1/|c_R|^2$ giving an accurate estimation of the number of shots for given accuracy ϵ . One can also see that the standard deviation is almost constant for hydrogen chains with various numbers of atoms, while the absolute error is highly dependent on the number of atoms. Comparing the three 12-qubit systems, it can be seen that the difference between the standard deviation and the absolute error depends on the system.

3. Accuracy of expectation values of observables other than the Hamiltonian in QSCI

Here, we examine the accuracy of the expectation values of observables other than the Hamiltonian, estimated for the output state obtained by QSCI calculation. Figure 13 shows the histograms for absolute errors of the expectation values for the gradient and Hessian, where the absolute error for an observable $\hat{O}$ is defined by

$$|\langle \psi_{\text{out}} | \hat{O} | \psi_{\text{out}} \rangle - \langle \psi_{\text{exact}} | \hat{O} | \psi_{\text{exact}} \rangle|. \quad (\text{D1})$$

Here, $|\psi_{\text{out}}\rangle$ is the output state of QSCI calculation with the idealized sampling from the exact ground state with R given in Fig. 5 for each error tolerance ϵ for energy and $|\psi_{\text{exact}}\rangle$ is the exact ground state. The observables $\hat{O}$ are set to be the nuclear gradient $\frac{\partial \hat{H}}{\partial x_i}$ ($i = 1, \dots, 3N_{\text{atom}}$) and the Hessian $\frac{\partial^2 \hat{H}}{\partial x_i \partial x_j}$ ($i, j = 1, \dots, 3N_{\text{atom}}$), where N_{atom} is the number of atoms in the molecule and x_i are coordinates of the nuclei. The absolute error is shown in the unit of hartree, hartree/ $\text{\AA}$, or hartree/ $\text{\AA}^2$, depending on the observables. Although there are some observables (i.e., components of the gradient or Hessian) whose expectation values exhibit larger absolute errors than that of the energy, the expectation values of the majority of the observables have similar accuracy as the energy.

4. Bond length dependence

The Hartree-Fock calculation is known to perform better for a stable geometry of a molecule than for the dissociation limit, so it is worth studying if QSCI also performs worse in the dissociation limit. Figure 14 shows the result of the same numerical analysis as Fig. 5, but for various bond lengths of H_2O molecules. The Hamiltonian is generated by the Hartree-Fock orbitals using STO-3G basis without specifying the active space, and is of 14-qubit after the Jordan-Wigner mapping. The bond lengths of two H–O bonds are taken to be equal, and the H–O–H angle is fixed to 104.45° . The result implies that, although there is some dependency on the bond length for larger ϵ , the dependency disappears for smaller ϵ . It can be expected from the result that the potential energy surface calculated by QSCI has a relatively constant accuracy, at least compared to the Hartree-Fock result, when the error tolerance is not very large.

5. Comparison to ASCI

Here, we investigate if there is a possibility that QSCI outperforms the state-of-the-art selected CI methods by taking ASCI for illustration. ASCI is a selected CI method solely based on classical computation, which adaptively searches for the optimal subspace of the Fock space for the diagonaliza-




FIG. 13. Histograms of absolute errors of nuclear gradients and Hessians for H_4 , H_6 , and H_8 molecules in QSCI, compared to the exact, CASCI value. Each plot corresponds to QSCI calculations with an energy error ϵ . Absolute errors are in units of hartree, hartree/ $\text{\AA}$, or hartree/ $\text{\AA}^2$, depending on the observables.

tion. In Fig. 15, we compare QSCI with ASCI, for which we follow the description in Ref. [65]. Here, we use the QSCI method with the idealized sampling from the ground state obtained by the exact diagonalization (full-CI) calculation.




FIG. 14. Estimated R and $1/|c_R|^2$ for H_2O molecule (14 qubits) with various bond lengths. Potential energy curves are also shown for reference. The estimation method of R and $1/|c_R|^2$ are the same as in Figs. 5 and 6.

The target molecule is the linear hydrogen chain H_{10} with the equal separation of $1.0 \text{ \AA}$. The basis set is STO-3G and the Hamiltonian with the Hartree-Fock orbitals is mapped to the 20 qubit one by the Jordan-Wigner mapping. For ASCI, in addition to the parameter R (called as N_{dets} in Ref. [65]), there are two additional parameters: They are denoted by ϵ and N_{cdets} in that paper, and are denoted by δ and R_{core} , respectively, in the following. The parameters δ and R_{core} determine the size of the search space for the iterative search for the new determinants, while the cost for the generation and diagonalization of the Hamiltonian, which is common for both ASCI and QSCI, is determined solely by R . We fixed $\delta = 0.05$ hartree and $r := R/R_{\text{core}} = 10$ or 20 for ASCI, and run QSCI and ASCI calculations with various R .

While in the case of $r = 10$, the two methods perform similarly, QSCI performs better for $r = 20$, where less computational cost is required for searching for a better set of




FIG. 15. Comparison of ASCI and QSCI for H_{10} molecule (20 qubits). The ASCI and QSCI energies as the difference to the exact CASCI energy are plotted against the size R of the selected subspace. The parameter r determines the size of the search space for ASCI.

configurations in ASCI. The result shows that, depending on the hyperparameters for ASCI, there is a possibility that QSCI performs better, at least in the case of the idealized sampling from the exact ground state. It should be noted that, in the present analysis, QSCI assumes knowledge of the exact ground state for sampling, whereas ASCI does not. Thus, ASCI may involve a more difficult task than QSCI in this comparison.

6. Noisy QSCI with input state preparation by noisy VQE

We present the simulation result of noisy QSCI with input states prepared by noisy VQE. More precisely, first we ex-




FIG. 16. Simulation results of noisy QSCI with noisy VQE input states for the hydrogen chain H_4 of eight qubits. Noisy VQE result is shown by the green line, and corresponding exact energy expectation value at each iteration is shown by the blue line. QSCI calculation is performed at every 100 iterations and the result of three trials is shown in a thin line, while the mean value is plotted in a solid line with markers.

cuted a simulation of VQE of the eight-qubit Hamiltonian of H_4 molecule as in Sec. IV with a noise model that envisions slightly better performance than current devices: single-qubit gate error and two-qubit error being 0.01% and 0.1%, respectively, both of which are modeled by the depolarizing noise. Measurement error is modeled by the bit-flip error of 0.1% rate at the end of the circuit. We employed the Hardware-EfficientReal ansatz of QURI Parts [102], with the number of layers (reps) being 8. We used the Nakanishi-Fujii-Todo optimizer [153] for optimization. At each estimation of the energy expectation values, we used QWC grouping [110] and 10^5 shots are distributed to each group according to Refs. [120,121].

At every 100 iterations, QSCI with postselection is performed with 10^5 shots, along with the noisy and noiseless state-vector estimation of the expectation value. The result is shown in Fig. 16. As one can see, QSCI result is better than

noisy VQE by almost an order of magnitude, going below the noiseless state-vector estimation even on a noisy simulator, at an early stage of the optimization. While it is true that the accuracy itself is not satisfactory, arguably due to the lack of accuracy of the VQE input state, it is clear that QSCI with VQE input state performs much better than a calculation only with VQE. We can expect that, with further reduction of device noise and improvements in the initial state preparation, QSCI may become a practically viable approach for accurate quantum simulations. Indeed, after the appearance of our preprint, Robledo-Moreno *et al.* [154] demonstrated that QSCI with a noise-mitigation technique called configuration recovery can run on a real quantum device as large as 77 qubits and can accurately calculate the ground-state energies of several molecules, which may be beyond the reach of classical methods. Their result illustrates the potential of noise robustness of QSCI.

-
- [1] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, R. Biswas, S. Boixo, F. G. Brandao, D. A. Buell, *et al.*, Quantum supremacy using a programmable superconducting processor, *Nature (London)* **574**, 505 (2019).
 - [2] H.-S. Zhong, H. Wang, Y.-H. Deng, M.-C. Chen, L.-C. Peng, Y.-H. Luo, J. Qin, D. Wu, X. Ding, Y. Hu, *et al.*, Quantum computational advantage using photons, *Science* **370**, 1460 (2020).
 - [3] L. S. Madsen, F. Laudenbach, M. F. Askarani, F. Rortais, T. Vincent, J. F. Bulmer, F. M. Miatto, L. Neuhaus, L. G. Helt, M. J. Collins, *et al.*, Quantum computational advantage with a programmable photonic processor, *Nature (London)* **606**, 75 (2022).
 - [4] J. Preskill, Quantum computing in the NISQ era and beyond, *Quantum* **2**, 79 (2018).
 - [5] Y. Cao, J. Romero, J. P. Olson, M. Degroote, P. D. Johnson, M. Kieferová, I. D. Kivlichan, T. Menke, B. Peropadre, N. P. Sawaya, *et al.*, Quantum chemistry in the age of quantum computing, *Chem. Rev.* **119**, 10856 (2019).
 - [6] S. McArdle, S. Endo, A. Aspuru-Guzik, S. C. Benjamin, and X. Yuan, Quantum computational chemistry, *Rev. Mod. Phys.* **92**, 015003 (2020).
 - [7] M. Cerezo, A. Arrasmith, R. Babbush, S. C. Benjamin, S. Endo, K. Fujii, J. R. McClean, K. Mitarai, X. Yuan, L. Cincio, *et al.*, Variational quantum algorithms, *Nat. Rev. Phys.* **3**, 625 (2021).
 - [8] K. Bharti, A. Cervera-Lierta, T. H. Kyaw, T. Haug, S. Alperin-Lea, A. Anand, M. Degroote, H. Heimonen, J. S. Kottmann, T. Menke, W.-K. Mok, S. Sim, L.-C. Kwek, and A. Aspuru-Guzik, Noisy intermediate-scale quantum algorithms, *Rev. Mod. Phys.* **94**, 015004 (2022).
 - [9] J. Tilly, H. Chen, S. Cao, D. Picozzi, K. Setia, Y. Li, E. Grant, L. Wossnig, I. Rungger, G. H. Booth, *et al.*, The variational quantum eigensolver: A review of methods and best practices, *Phys. Rep.* **986**, 1 (2022).
 - [10] A. Peruzzo, J. McClean, P. Shadbolt, M.-H. Yung, X.-Q. Zhou, P. J. Love, A. Aspuru-Guzik, and J. L. O'Brien, A variational eigenvalue solver on a photonic quantum processor, *Nat. Commun.* **5**, 4213 (2014).
 - [11] J. F. Gonthier, M. D. Radin, C. Buda, E. J. Duskocil, C. M. Abuan, and J. Romero, Measurements as a roadblock to near-term practical quantum advantage in chemistry: Resource analysis, *Phys. Rev. Res.* **4**, 033154 (2022).
 - [12] L. Viola, E. Knill, and S. Lloyd, Dynamical decoupling of open quantum systems, *Phys. Rev. Lett.* **82**, 2417 (1999).
 - [13] K. Temme, S. Bravyi, and J. M. Gambetta, Error mitigation for short-depth quantum circuits, *Phys. Rev. Lett.* **119**, 180509 (2017).
 - [14] Y. Li and S. C. Benjamin, Efficient variational quantum simulator incorporating active error minimization, *Phys. Rev. X* **7**, 021050 (2017).
 - [15] S. Endo, S. C. Benjamin, and Y. Li, Practical quantum error mitigation for near-future applications, *Phys. Rev. X* **8**, 031027 (2018).
 - [16] B. Koczor, Exponential error suppression for near-term quantum devices, *Phys. Rev. X* **11**, 031057 (2021).
 - [17] W. J. Huggins, S. McArdle, T. E. O'Brien, J. Lee, N. C. Rubin, S. Boixo, K. B. Whaley, R. Babbush, and J. R. McClean, Virtual distillation for quantum error mitigation, *Phys. Rev. X* **11**, 041036 (2021).
 - [18] S. McArdle, X. Yuan, and S. Benjamin, Error-mitigated digital quantum simulation, *Phys. Rev. Lett.* **122**, 180501 (2019).
 - [19] X. Bonet-Monroig, R. Sagastizabal, M. Singh, and T. E. O'Brien, Low-cost error mitigation by symmetry verification, *Phys. Rev. A* **98**, 062339 (2018).
 - [20] F. B. Maciejewski, Z. Zimborás, and M. Oszmaniec, Mitigation of readout noise in near-term quantum devices by classical post-processing based on detector tomography, *Quantum* **4**, 257 (2020).
 - [21] S. Endo, Z. Cai, S. C. Benjamin, and X. Yuan, Hybrid quantum-classical algorithms and quantum error mitigation, *J. Phys. Soc. Jpn.* **90**, 032001 (2021).
 - [22] S. Wang, P. Czarnik, A. Arrasmith, M. Cerezo, L. Cincio, and P. J. Coles, Can error mitigation improve trainability of noisy variational quantum algorithms? *Quantum* **8**, 1287 (2024).
 - [23] R. Takagi, S. Endo, S. Minagawa, and M. Gu, Fundamental limits of quantum error mitigation, *npj Quantum Inf.* **8**, 114 (2022).

- [24] K. Tsubouchi, T. Sagawa, and N. Yoshioka, Universal cost bound of quantum error mitigation based on quantum estimation theory, *Phys. Rev. Lett.* **131**, 210601 (2023).
- [25] R. Takagi, H. Tajima, and M. Gu, Universal sample lower bounds for quantum error mitigation, *Phys. Rev. Lett.* **131**, 210602 (2023).
- [26] F. Coester and H. Kümmel, Short-range correlations in nuclear wave functions, *Nucl. Phys.* **17**, 477 (1960).
- [27] J. Čížek, On the correlation problem in atomic and molecular systems. Calculation of wavefunction components in Ursell-type expansion using quantum-field theoretical methods, *J. Chem. Phys.* **45**, 4256 (1966).
- [28] J. R. McClean, S. Boixo, V. N. Smelyanskiy, R. Babbush, and H. Neven, Barren plateaus in quantum neural network training landscapes, *Nat. Commun.* **9**, 4812 (2018).
- [29] We focus on applications to quantum chemistry in this paper. However, the proposed methods can be applied to a variety of many-body Hamiltonians, including many electron and spin problems in condensed matter physics.
- [30] QSCI can also set rigorous upper bounds on the excited-state energies, depending on its algorithmic implementation.
- [31] M. Kohda, R. Imai, K. Kanno, K. Mitarai, W. Mizukami, and Y. O. Nakagawa, Quantum expectation-value estimation by computational basis sampling, *Phys. Rev. Res.* **4**, 033173 (2022).
- [32] B. M. Terhal and D. P. DiVincenzo, Adaptive quantum computation, constant depth quantum circuits and Arthur-Merlin games, *Quantum Inf. Comput.* **4**, 134 (2004).
- [33] O. Leimkuhler and K. B. Whaley, Exponential quantum speedups in quantum chemistry with linear depth, [arXiv:2503.21041](https://arxiv.org/abs/2503.21041).
- [34] A. Hafid, H. Iwakiri, K. Tsubouchi, N. Yoshioka, and M. Kohda, Hardness of classically sampling quantum chemistry circuits, [arXiv:2504.12893](https://arxiv.org/abs/2504.12893).
- [35] H.-Y. Huang, R. Kueng, and J. Preskill, Predicting many properties of a quantum system from very few measurements, *Nat. Phys.* **16**, 1050 (2020).
- [36] A. Zhao, N. C. Rubin, and A. Miyake, Fermionic partial tomography via classical shadows, *Phys. Rev. Lett.* **127**, 110504 (2021).
- [37] G. Torlai, G. Mazzola, J. Carrasquilla, M. Troyer, R. Melko, and G. Carleo, Neural-network quantum state tomography, *Nat. Phys.* **14**, 447 (2018).
- [38] M. Cramer, M. B. Plenio, S. T. Flammia, R. Somma, D. Gross, S. D. Bartlett, O. Landon-Cardinal, D. Poulin, and Y.-K. Liu, Efficient quantum state tomography, *Nat. Commun.* **1**, 149 (2010).
- [39] T. Helgaker, P. Jørgensen, and J. Olsen, *Molecular Electronic-Structure Theory* (John Wiley & Sons, Chichester, UK, 2000).
- [40] J. R. McClean, M. E. Kimchi-Schwartz, J. Carter, and W. A. de Jong, Hybrid quantum-classical hierarchy for mitigation of decoherence and determination of excited states, *Phys. Rev. A* **95**, 042308 (2017).
- [41] T. Takeshita, N. C. Rubin, Z. Jiang, E. Lee, R. Babbush, and J. R. McClean, Increasing the representation accuracy of quantum simulations of chemistry without extra quantum resources, *Phys. Rev. X* **10**, 011004 (2020).
- [42] M. Urbanek, D. Camps, R. Van Beeumen, and W. A. de Jong, Chemistry on quantum computers with virtual quantum subspace expansion, *J. Chem. Theory Comput.* **16**, 5425 (2020).
- [43] C. F. Bender and E. R. Davidson, Studies in configuration interaction: The first-row diatomic hydrides, *Phys. Rev.* **183**, 23 (1969).
- [44] J. Whitten and M. Hackmeyer, Configuration interaction studies of ground and excited states of polyatomic molecules. I. The CI formulation and studies of formaldehyde, *J. Chem. Phys.* **51**, 5584 (1969).
- [45] B. Huron, J. Malrieu, and P. Rancurel, Iterative perturbation calculations of ground and excited state energies from multiconfigurational zeroth-order wavefunctions, *J. Chem. Phys.* **58**, 5745 (1973).
- [46] R. J. Buenker and S. D. Peyerimhoff, Individualized configuration selection in CI calculations with subsequent energy extrapolation, *Theor. Chim. Acta* **35**, 33 (1974).
- [47] R. J. Buenker and S. D. Peyerimhoff, Energy extrapolation in CI calculations, *Theor. Chim. Acta* **39**, 217 (1975).
- [48] H. Nakatsuji, Cluster expansion of the wavefunction, valence and Rydberg excitations, ionizations, and inner-valence ionizations of CO₂ and N₂O studied by the SAC and SAC CI theories, *Chem. Phys.* **75**, 425 (1983).
- [49] R. Cimiraglia and M. Persico, Recent advances in multireference second order perturbation CI: The CIPSI method revisited, *J. Comput. Chem.* **8**, 39 (1987).
- [50] R. J. Harrison, Approximating full configuration interaction with selected configuration interaction and perturbation theory, *J. Chem. Phys.* **94**, 5021 (1991).
- [51] J. Greer, Estimating full configuration interaction limits from a Monte Carlo selection of the expansion space, *J. Chem. Phys.* **103**, 1821 (1995).
- [52] J. Greer, Monte Carlo configuration interaction, *J. Comput. Phys.* **146**, 181 (1998).
- [53] F. A. Evangelista, Adaptive multiconfigurational wave functions, *J. Chem. Phys.* **140**, 124114 (2014).
- [54] A. A. Holmes, H. J. Changlani, and C. Umrigar, Efficient heat-bath sampling in Fock space, *J. Chem. Theory Comput.* **12**, 1561 (2016).
- [55] J. B. Schriber and F. A. Evangelista, Communication: An adaptive configuration interaction approach for strongly correlated electrons with tunable accuracy, *J. Chem. Phys.* **144**, 161106 (2016).
- [56] A. A. Holmes, N. M. Tubman, and C. Umrigar, Heat-bath configuration interaction: An efficient selected configuration interaction algorithm inspired by heat-bath sampling, *J. Chem. Theory Comput.* **12**, 3674 (2016).
- [57] N. M. Tubman, J. Lee, T. Y. Takeshita, M. Head-Gordon, and K. B. Whaley, A deterministic alternative to the full configuration interaction quantum Monte Carlo method, *J. Chem. Phys.* **145**, 044112 (2016).
- [58] Y. Ohtsuka and J. Hasegawa, Selected configuration interaction using sampled first-order corrections to wave functions, *J. Chem. Phys.* **147**, 034102 (2017).
- [59] J. B. Schriber and F. A. Evangelista, Adaptive configuration interaction for computing challenging electronic excited states with tunable accuracy, *J. Chem. Theory Comput.* **13**, 5354 (2017).
- [60] S. Sharma, A. A. Holmes, G. Jeanmairet, A. Alavi, and C. J. Umrigar, Semistochastic heat-bath configuration interaction method: Selected configuration interaction with semistochas-

- tic perturbation theory, *J. Chem. Theory Comput.* **13**, 1595 (2017).
- [61] R. Chakraborty, P. Ghosh, and D. Ghosh, Evolutionary algorithm based configuration interaction approach, *Int. J. Quantum Chem.* **118**, e25509 (2018).
- [62] J. P. Coe, Machine learning configuration interaction, *J. Chem. Theory Comput.* **14**, 5739 (2018).
- [63] J. P. Coe, Machine learning configuration interaction for *ab initio* potential energy curves, *J. Chem. Theory Comput.* **15**, 6179 (2019).
- [64] V. Abraham and N. J. Mayhall, Selected configuration interaction in a basis of cluster state tensor products, *J. Chem. Theory Comput.* **16**, 6098 (2020).
- [65] N. M. Tubman, C. D. Freeman, D. S. Levine, D. Hait, M. Head-Gordon, and K. B. Whaley, Modern approaches to exact diagonalization and selected configuration interaction with the adaptive sampling CI method, *J. Chem. Theory Comput.* **16**, 2139 (2020).
- [66] N. Zhang, W. Liu, and M. R. Hoffmann, Iterative configuration interaction with selection, *J. Chem. Theory Comput.* **16**, 2296 (2020).
- [67] N. Zhang, W. Liu, and M. R. Hoffmann, Further development of iCIPT2 for strongly correlated electrons, *J. Chem. Theory Comput.* **17**, 949 (2021).
- [68] V. G. Chilkuri and F. Neese, Comparison of many-particle representations for selected-CI I: A tree based approach, *J. Comput. Chem.* **42**, 982 (2021).
- [69] V. G. Chilkuri and F. Neese, Comparison of many-particle representations for selected configuration interaction: II. Numerical benchmark calculations, *J. Chem. Theory Comput.* **17**, 2868 (2021).
- [70] J. J. Goings, H. Hu, C. Yang, and X. Li, Reinforcement learning configuration interaction, *J. Chem. Theory Comput.* **17**, 5482 (2021).
- [71] S. D. Pineda Flores, Chembot: A machine learning approach to selective configuration interaction, *J. Chem. Theory Comput.* **17**, 4028 (2021).
- [72] W. Jeong, C. A. Gaglioli, and L. Gagliardi, Active learning configuration interaction for excited-state calculations of polycyclic aromatic hydrocarbons, *J. Chem. Theory Comput.* **17**, 7518 (2021).
- [73] J. P. Coe, Analytic gradients for selected configuration interaction, *J. Chem. Theory Comput.* **19**, 874 (2023).
- [74] K. Seth and D. Ghosh, Active learning assisted MCCI to target spin states, *J. Chem. Theory Comput.* **19**, 524 (2023).
- [75] N. Yoshioka, T. Sato, Y. O. Nakagawa, Y.-Y. Ohnishi, and W. Mizukami, Variational quantum simulation for periodic materials, *Phys. Rev. Res.* **4**, 013052 (2022).
- [76] J. R. McClean, Z. Jiang, N. C. Rubin, R. Babbush, and H. Neven, Decoding quantum errors with subspace expansions, *Nat. Commun.* **11**, 636 (2020).
- [77] N. Yoshioka, H. Hakoshima, Y. Matsuzaki, Y. Tokunaga, Y. Suzuki, and S. Endo, Generalized quantum subspace expansion, *Phys. Rev. Lett.* **129**, 020502 (2022).
- [78] E. N. Epperly, L. Lin, and Y. Nakatsukasa, A theory of quantum subspace diagonalization, *SIAM J. Matrix Anal. Appl.* **43**, 1263 (2022).
- [79] W. J. Huggins, J. Lee, U. Baek, B. O’Gorman, and K. B. Whaley, A non-orthogonal variational quantum eigensolver, *New J. Phys.* **22**, 073009 (2020).
- [80] M. Motta, C. Sun, A. T. Tan, M. J. O’Rourke, E. Ye, A. J. Minnich, F. G. Brandão, and G. K.-L. Chan, Determining eigenstates and thermal states on a quantum computer using quantum imaginary time evolution, *Nat. Phys.* **16**, 205 (2020).
- [81] R. M. Parrish and P. L. McMahon, Quantum filter diagonalization: Quantum eigendecomposition without full quantum phase estimation, [arXiv:1909.08925](https://arxiv.org/abs/1909.08925).
- [82] N. H. Stair, R. Huang, and F. A. Evangelista, A multireference quantum Krylov algorithm for strongly correlated electrons, *J. Chem. Theory Comput.* **16**, 2236 (2020).
- [83] R. M. Parrish, E. G. Hohenstein, P. L. McMahon, and T. J. Martínez, Quantum computation of electronic transitions using a variational quantum eigensolver, *Phys. Rev. Lett.* **122**, 230401 (2019).
- [84] K. Seki and S. Yunoki, Quantum power method by a superposition of time-evolved states, *PRX Quantum* **2**, 010333 (2021).
- [85] U. Baek, D. Hait, J. Shee, O. Leimkuhler, W. J. Huggins, T. F. Stetina, M. Head-Gordon, and K. B. Whaley, Say no to optimization: A non-orthogonal quantum eigensolver, *PRX Quantum* **4**, 030307 (2023).
- [86] W. Kirby, M. Motta, and A. Mezzacapo, Exact and efficient Lanczos method on a quantum computer, *Quantum* **7**, 1018 (2023).
- [87] M. D. Radin and P. Johnson, Classically-boosted variational quantum eigensolver, [arXiv:2106.04755](https://arxiv.org/abs/2106.04755).
- [88] Instead, linear combinations of Slater determinants such as configuration state functions may be mapped to the computational basis states. QSCI can work with such a mapping if the matrix elements of the Hamiltonian in the computational basis can be efficiently computed by classical computation.
- [89] S. B. Bravyi and A. Y. Kitaev, Fermionic quantum computation, *Ann. Phys.* **298**, 210 (2002).
- [90] See, e.g., Eq. (1). There, the CI coefficients can be expressed as $\alpha_x = \langle x | \psi \rangle$.
- [91] In VQE, physical noise in the state preparation itself does not directly lead to the violation of the variational inequality, but it may be possible that an error during the measurement procedure causes it. The use of error-mitigation techniques can also lead to the breakdown of the inequality.
- [92] Note that an error that corresponds to a phase-flip error occurring at the end of a circuit does not affect the probability distribution $|\langle x | \psi_{\text{in}} \rangle|^2$ and hence the sampling outcome.
- [93] Note that the set $\mathcal{S}_R$ defined here agrees with the definition (2) in the preceding subsection when $N_{\text{in}} = 1$.
- [94] One can make each of R_i as equal as possible by the following cycle of procedures, starting from an empty set $\mathcal{S}_R$, for a given R : In the first cycle, for each of the N_{in} input states, the most frequent bit string is selected from the sampling outcome and then added to $\mathcal{S}_R$; this procedure is executed from the zeroth input state to $(N_{\text{in}} - 1)$ th input state, where one skips the state if the selected bit string already exists in $\mathcal{S}_R$; in the second cycle, the second frequent bit string is added to $\mathcal{S}_R$ for each input state according to the same rule; one repeats such a cycle until $\mathcal{S}_R$ is filled with R distinct bit strings. Suppose such procedures are finished after completing R' cycles. Then, one can ensure that at least R' most frequent bit strings for each input state are included in $\mathcal{S}_R$. This implies R' or $R' + 1$ most important configurations in each input state are included in the

- common subspace (7), in the ideal situation where statistical fluctuation and physical noise can be ignored.
- [95] E. Hylleraas and B. Undheim, Numerical calculation of the 2S terms of ortho- and para-helium, *Z. Phys.* **65**, 759 (1930).
- [96] J. MacDonald, Successive approximations by the Rayleigh-Ritz variation method, *Phys. Rev.* **43**, 830 (1933).
- [97] O. Higgott, D. Wang, and S. Brierley, Variational quantum computation of excited states, *Quantum* **3**, 156 (2019).
- [98] We implicitly assume the ground state is unique for ease of illustration. One can straightforwardly translate the description here to cases of degenerate ground (and possibly excited) states.
- [99] This is not the unique choice of the effective Hamiltonian. For instance, the orthogonality can be imposed without introducing extra parameters though the implementation would be less suitable for NISQ devices [100].
- [100] J. Lee, W. J. Huggins, M. Head-Gordon, and K. B. Whaley, Generalized unitary coupled cluster wave functions for quantum computation, *J. Chem. Theory Comput.* **15**, 311 (2019).
- [101] Y. Suzuki, Y. Kawase, Y. Masumura, Y. Hiraga, M. Nakadai, J. Chen, K. M. Nakanishi, K. Mitarai, R. Imai, S. Tamiya, *et al.*, Qulacs: A fast and versatile quantum circuit simulator for research purpose, *Quantum* **5**, 559 (2021).
- [102] QURI Parts, GitHub (2022), <https://github.com/QunaSys/quri-parts>.
- [103] J. R. McClean, N. C. Rubin, K. J. Sung, I. D. Kivlichan, X. Bonet-Monroig, Y. Cao, C. Dai, E. S. Fried, C. Gidney, B. Gimby, *et al.*, OpenFermion: The electronic structure package for quantum computers, *Quantum Sci. Technol.* **5**, 034014 (2020).
- [104] Q. Sun, T. C. Berkelbach, N. S. Blunt, G. H. Booth, S. Guo, Z. Li, J. Liu, J. D. McClain, E. R. Sayfutyarova, S. Sharma, *et al.*, PySCF: The Python-based simulations of chemistry framework, *WIREs Comput. Mol. Sci.* **8**, e1340 (2018).
- [105] B. T. Gard, L. Zhu, G. S. Barron, N. J. Mayhall, S. E. Economou, and E. Barnes, Efficient symmetry-preserving state preparation circuits for the variational quantum eigensolver algorithm, *npj Quantum Inf.* **6**, 10 (2020).
- [106] Y. Ibe, Y. O. Nakagawa, N. Earnest, T. Yamamoto, K. Mitarai, Q. Gao, and T. Kobayashi, Calculating transition amplitudes by variational quantum deflation, *Phys. Rev. Res.* **4**, 013173 (2022).
- [107] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, *et al.*, SciPy 1.0: Fundamental algorithms for scientific computing in Python, *Nat. Methods* **17**, 261 (2020).
- [108] In this paper, we define the chemical accuracy by 1 kcal/mol $\simeq 1.6 \times 10^{-3}$ hartree for the deviation of the calculated energy from the one obtained by the exact diagonalization of the Hamiltonian.
- [109] For the active space restriction of five active orbitals and six active electrons with $S_z = 0$, the number of the Slater determinants is $\binom{5}{3} \times \binom{5}{3} = 100$. If one does not know the number of electrons and S_z of the ground state before the calculation, then one would need to deal with the full Hamiltonian in the Fock space of $2^{10} = 1024$ dimensions.
- [110] J. R. McClean, J. Romero, R. Babbush, and A. Aspuru-Guzik, The theory of variational hybrid quantum-classical algorithms, *New J. Phys.* **18**, 023023 (2016).
- [111] I. G. Ryabinkin, S. N. Genin, and A. F. Izmaylov, Constrained variational quantum eigensolver: Quantum computer search engine in the Fock space, *J. Chem. Theory Comput.* **15**, 249 (2019).
- [112] K. Kuroiwa and Y. O. Nakagawa, Penalty methods for a variational quantum eigensolver, *Phys. Rev. Res.* **3**, 013197 (2021).
- [113] To show this explicitly, assume $N_s = 2$ for simplicity. The single diagonalization with the subspace $\mathcal{S}_R = \mathcal{S}_{R_0}^{(0)} \cup \mathcal{S}_{R_1}^{(1)}$, where the subspaces on the right-hand side denote those of the sequential diagonalization, has at least the same representability as the sequential diagonalization calculation with $\mathcal{S}_{R_1}^{(1)}$.
- [114] H. R. Larsson, H. Zhai, C. J. Umrigar, and G. K.-L. Chan, The chromium dimer: Closing a chapter of quantum chemistry, *J. Am. Chem. Soc.* **144**, 15932 (2022).
- [115] M. Motta, C. Genovese, F. Ma, Z.-H. Cui, R. Sawaya, G. K.-L. Chan, N. Chepiga, P. Helms, C. Jiménez-Hoyos, A. J. Millis, *et al.*, Ground-state properties of the hydrogen chain: Dimerization, insulator-to-metal transition, and magnetic phases, *Phys. Rev. X* **10**, 031058 (2020).
- [116] P. Stampfuß, M. Vogel, and W. Wenzel, Improved implementation and application of the individually selecting configuration interaction method, in *High Performance Computing in Science and Engineering '02*, edited by E. Krause and W. Jäger (Springer, Berlin, Heidelberg, 2003), pp. 215–229.
- [117] Y. Garniron, T. Applencourt, K. Gasperich, A. Benali, A. Ferté, J. Paquier, B. Pradines, R. Assaraf, P. Reinhardt, J. Toulouse, *et al.*, Quantum package 2.0: An open-source determinant-driven suite of programs, *J. Chem. Theory Comput.* **15**, 3591 (2019).
- [118] O. Crawford, B. van Straaten, D. Wang, T. Parks, E. Campbell, and S. Brierley, Efficient quantum measurement of Pauli operators in the presence of finite sampling error, *Quantum* **5**, 385 (2021).
- [119] This shot allocation may not be possible in practice without prior knowledge of the exact ground state, but this estimation gives the lower bound on the required number of total shots among possible shot allocation strategies, for a given error tolerance with the given grouping method and the state.
- [120] D. Wecker, M. B. Hastings, and M. Troyer, Progress towards practical quantum variational algorithms, *Phys. Rev. A* **92**, 042303 (2015).
- [121] N. C. Rubin, R. Babbush, and J. McClean, Application of fermionic marginal constraints to hybrid quantum algorithms, *New J. Phys.* **20**, 053020 (2018).
- [122] More precisely, the fit was performed by a function
- $$\ln(N_{\text{shot}}(N_q)) = B \ln(N_q) + A,$$
- where A and B are the free parameters. Similarly, in Fig. 5, a linear function $cN_q + d$ was used to fit the data for $\ln(R)$, rather than $D \times 2^{CN_q}$ for R .
- [123] It is numerically shown in Appendix D 3 that the accuracy of the gradients and Hessians in QSCI is of the same order as ϵ when expressed in the units of hartree/Å and hartree/Å², respectively.
- [124] More precisely, the error rate of the single-qubit depolarizing noise for each single-qubit gate is set to p_1 , where p_1 is the single-qubit gate infidelity. For the two-qubit gates, single-qubit depolarizing noise is applied to each of the two qubits with probability $1 - \sqrt{1 - p_2}$ for a two-qubit gate infidelity

- p_2 . The bit-flip noise is applied to each qubit with probability p_{ro} , the measurement infidelity.
- [125] The number of Slater determinants that have the required particle number and S_z is $\binom{4}{2} \times \binom{4}{2} = 36$.
 - [126] S. R. White, Density matrix formulation for quantum renormalization groups, *Phys. Rev. Lett.* **69**, 2863 (1992).
 - [127] S. R. White, Density-matrix algorithms for quantum renormalization groups, *Phys. Rev. B* **48**, 10345 (1993).
 - [128] U. Schollwöck, The density-matrix renormalization group, *Rev. Mod. Phys.* **77**, 259 (2005).
 - [129] E. Farhi, J. Goldstone, S. Gutmann, and M. Sipser, Quantum computation by adiabatic evolution, [arXiv:quant-ph/0001106](https://arxiv.org/abs/quant-ph/0001106).
 - [130] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, Simulated quantum computation of molecular energies, *Science* **309**, 1704 (2005).
 - [131] C. P. Williams, Probabilistic nonunitary quantum computing, in *Quantum Information and Computation II*, edited by E. Donkor, A. R. Pirich, and H. E. Brandt (International Society for Optics and Photonics, Bellingham, WA, 2004), Vol. 5436, pp. 297–306.
 - [132] H. Terashima and M. Ueda, Nonunitary quantum circuit, *Int. J. Quantum Inf.* **03**, 633 (2005).
 - [133] S. McArdle, T. Jones, S. Endo, Y. Li, S. C. Benjamin, and X. Yuan, Variational ansatz-based quantum simulation of imaginary time evolution, *npj Quantum Inf.* **5**, 75 (2019).
 - [134] Y. Mao, M. Chaudhary, M. Kondappan, J. Shi, E. O. Ilo-Okeke, V. Ivannikov, and T. Byrnes, Measurement-based deterministic imaginary time evolution, *Phys. Rev. Lett.* **131**, 110602 (2023).
 - [135] K. N. Okada, K. Osaki, K. Mitarai, and K. Fujii, Identification of topological phases using classically-optimized variational quantum eigensolver, *Phys. Rev. Res.* **5**, 043217 (2023).
 - [136] J. Romero, R. Babbush, J. R. McClean, C. Hempel, P. J. Love, and A. Aspuru-Guzik, Strategies for quantum computing molecular energies using the unitary coupled cluster ansatz, *Quantum Sci. Technol.* **4**, 014008 (2019).
 - [137] K. Kuroiwa and Y. O. Nakagawa, Averaging gate approximation error and performance of unitary coupled cluster ansatz in pre-FTQC era, *Quantum* **9**, 1800 (2025).
 - [138] M. R. Hirsbrunner, D. Chamaki, J. W. Mullinax, and N. M. Tubman, Beyond MP2 initialization for unitary coupled cluster quantum circuits, *Quantum* **8**, 1538 (2024).
 - [139] K. Mitarai, Y. Suzuki, W. Mizukami, Y. O. Nakagawa, and K. Fujii, Quadratic Clifford expansion for efficient benchmarking and initialization of variational quantum algorithms, *Phys. Rev. Res.* **4**, 033012 (2022).
 - [140] G. S. Ravi, P. Gokhale, Y. Ding, W. Kirby, K. Smith, J. M. Baker, P. J. Love, H. Hoffmann, K. R. Brown, and F. T. Chong, CAFQA: A classical simulation bootstrap for variational quantum algorithms, in *Proceedings of the 28th ACM International Conference on Architectural Support for Programming Languages and Operating Systems* (Association for Computing Machinery, New York, NY, 2022), Vol. 1, pp. 15–29.
 - [141] F. Verstraete, J. I. Cirac, and J. I. Latorre, Quantum circuits for strongly correlated quantum systems, *Phys. Rev. A* **79**, 032316 (2009).
 - [142] J. S. Van Dyke, G. S. Barron, N. J. Mayhall, E. Barnes, and S. E. Economou, Preparing Bethe ansatz eigenstates on a quantum computer, *PRX Quantum* **2**, 040329 (2021).
 - [143] A. Sopena, M. H. Gordon, D. García-Martín, G. Sierra, and E. López, Algebraic Bethe circuits, *Quantum* **6**, 796 (2022).
 - [144] A. J. Ferris and G. Vidal, Perfect sampling with unitary tensor networks, *Phys. Rev. B* **85**, 165146 (2012).
 - [145] G. H. Booth, A. J. W. Thom, and A. Alavi, Fermion Monte Carlo without fixed nodes: A game of life, death, and annihilation in Slater determinant space, *J. Chem. Phys.* **131**, 054106 (2009).
 - [146] D. Cleland, G. H. Booth, and A. Alavi, Communications: Survival of the fittest: Accelerating convergence in full configuration-interaction quantum Monte Carlo, *J. Chem. Phys.* **132**, 041103 (2010).
 - [147] F. R. Petruzielo, A. A. Holmes, H. J. Changlani, M. P. Nightingale, and C. J. Umrigar, Semistochastic projector Monte Carlo method, *Phys. Rev. Lett.* **109**, 230201 (2012).
 - [148] J. T. Seeley, M. J. Richard, and P. J. Love, The Bravyi-Kitaev transformation for quantum computation of electronic structure, *J. Chem. Phys.* **137**, 224109 (2012).
 - [149] K. Mitarai, Y. O. Nakagawa, and W. Mizukami, Theory of analytical energy derivatives for the variational quantum eigensolver, *Phys. Rev. Res.* **2**, 013129 (2020).
 - [150] R. D. Johnson III, *et al.*, NIST computational chemistry comparison and benchmark database, NIST Standard Reference Database (2022), <https://cccbdb.nist.gov/>.
 - [151] Y.-L. Wang, H.-S. Hu, W.-L. Li, F. Wei, and J. Li, Relativistic effects break periodicity in group 6 diatomic molecules, *J. Am. Chem. Soc.* **138**, 1126 (2016).
 - [152] S. Kim, J. Chen, T. Cheng, A. Gindulyte, J. He, S. He, Q. Li, B. A. Shoemaker, P. A. Thiessen, B. Yu, *et al.*, PubChem 2023 update, *Nucl. Acids Res.* **51**, D1373 (2023).
 - [153] K. M. Nakanishi, K. Fujii, and S. Todo, Sequential minimal optimization for quantum-classical hybrid algorithms, *Phys. Rev. Res.* **2**, 043158 (2020).
 - [154] J. Robledo-Moreno, M. Motta, H. Haas, A. Javadi-Abhari, P. Jurcevic, W. Kirby, S. Martiel, K. Sharma, S. Sharma, T. Shirakawa, I. Sitdikov, R.-Y. Sun, K. J. Sung, M. Takita, M. C. Tran, S. Yunoki, and A. Mezzacapo, Chemistry beyond the scale of exact diagonalization on a quantum-centric supercomputer, *Sci. Adv.* **11**, eadu9991 (2025).