

Polarization symmetry as a signature of Bell states in Compton scattering

Peter Caradonna^{1,*} and Kenji Shimazoe^{1,2,†}

¹Department of Nuclear Engineering and Management, *The University of Tokyo*, Bunkyo-ku, Tokyo, Japan

²Department of Bioengineering, *The University of Tokyo*, Bunkyo-ku, Tokyo, Japan



(Received 30 December 2025; accepted 27 May 2026; published 12 June 2026)

We analyze bipartite photon states in coincidence Compton polarimetry at energies where polarization must be inferred from Compton scattering. Polarization symmetry, defined as equality of signal and idler degree-of-polarization responses for all scattering angles, holds if and only if the initial two-photon state is pure and Bell equivalent. This identifies polarization symmetry as a state-level invariant that uniquely witnesses Bell-orbit structure under coincidence Compton scattering.

DOI: [10.1103/prj4-9vk7](https://doi.org/10.1103/prj4-9vk7)

I. INTRODUCTION

Entanglement certification in photonic systems is typically based on polarization measurements using optical elements. In the photon energy range from a few hundred keV up to the MeV scale, this strategy has not found practical implementation. Instead, polarization is accessed indirectly through Compton scattering signatures inferred from detector measurements [1]. In this regime, coincidence Compton polarimetry has provided a standard framework for studying polarization correlations of back-to-back 511 keV annihilation photon pairs, both theoretically [2–8] and experimentally [9–16].

A separate line of theoretical work developed a Compton scattering map for 511 keV annihilation photons and used it to demonstrate polarization symmetry, in which the degree of polarization of the idler beam matched that of the scattered signal beam for every signal-photon scattering direction considered in coincidence [17]. That analysis was intentionally restricted to this specific physical system. The unresolved question is therefore a classification problem: Which bipartite photon states exhibit the same polarization symmetry under coincidence Compton scattering, and to what extent does this symmetry identify a distinctive entanglement structure at high photon energies? This question is pertinent to ongoing efforts to understand which polarization-correlation signatures in coincidence Compton polarimetry may be useful in entanglement-based extensions of positron emission tomography in the MeV regime [18]. More broadly, polarization-based signatures can function as operational probes of coherence and purity, although they need not remain intact once polarization becomes coupled to additional degrees of freedom [19].

II. POLARIZATION-SYMMETRY SOLUTIONS

We consider a general two-photon state produced by a photon-pair source, in which the signal photon undergoes a single Compton-scattering event from a stationary electron, as shown in Fig. 1. The electron spin is treated as unpolarized at the ensemble level, while the idler photon does not interact directly. The scattering direction of the signal photon is specified by the polar angle θ and azimuthal angle ϕ . The analysis is performed in coincidence, so the degree of polarization of the idler beam is evaluated conditionally on the subset of selected signal-photon scattering events with scattering direction (θ, ϕ) . The goal is to determine which initial two-photon states satisfy polarization symmetry, defined here as equality between the degrees of polarization of the scattered signal and coincidence-conditioned idler beams for every scattering direction (θ, ϕ) . This is the symmetry identified in Ref. [17] for 511 keV annihilation photons, and here we determine the full class of two-photon states for which it holds.

Because polarization information in Compton scattering is inferred statistically rather than through projective measurement, the analysis is formulated in the Stokes representation established in Refs. [6,7,17,20]. The strategy is to represent a general two-photon density matrix in Stokes space, apply the Compton scattering map to the signal subsystem, and then impose the polarization-symmetry condition algebraically for all (θ, ϕ) . Full intermediate algebra is given in Appendixes A–C.

Let ρ denote a general two-photon state in the circular-polarization basis $\{|R, R\rangle, |R, L\rangle, |L, R\rangle, |L, L\rangle\}$, where $|R\rangle$ and $|L\rangle$ denote the right- and left-circular polarization states, respectively, following the convention used in Ref. [6]. Its matrix elements are ρ_{mn} with $m, n = 1, \dots, 4$, and it satisfies $\rho = \rho^\dagger$ and $\text{Tr}[\rho] = 1$.

Express ρ in terms of the 4×4 Stokes correlation matrix $S(\rho)$, following Ref. [17], whose matrix elements are given by [21]

$$S_{\mu\nu} = \text{Tr}[(\sigma_\mu \otimes \sigma_\nu)\rho], \quad \mu, \nu = 0, 1, 2, 3.$$

For an initial state ρ with Stokes matrix $S(\rho)$, the evolved matrix in the scattering geometry of Fig. 1 is

*Contact author: petercaradonna65@gmail.com

†Contact author: shimazoe@g.ecc.u-tokyo.ac.jp




FIG. 1. Single Compton scattering of the signal photon with incident energy E_0 from a general photon-pair source. The signal photon scatters from a stationary electron by a polar angle θ and azimuthal angle ϕ , emerging with energy $E(\theta)$, while the idler photon is selected through coincidence conditioning. Kinematic conventions follow Ref. [17].

described by

$$S'(\rho) = S(\rho) M(-\phi) T_{\text{un}}(\theta), \quad (1)$$

where $M(\phi)$ rotates the signal Stokes vector by 2ϕ about its circular-polarization axis and $T_{\text{un}}(\theta)$ is the symmetric unpolarized Compton-scattering Mueller matrix with entries $t_{11}, t_{12}, t_{33}, t_{44}$ [7]. The quantity $S'(\rho)$ is an unnormalized, intensity-weighted Stokes matrix associated with the selected Compton-scattering channel and geometry rather than the Stokes matrix of a trace-normalized output density matrix. A normalized output density matrix could be recovered through the inverse Stokes-to-density-matrix map [20], which includes the required renormalization.

From $S'(\rho)$, the degrees of polarization of the coincidence-conditioned idler beam and the scattered signal beam, denoted by p_i and p_s , respectively, are obtained from the matrix elements in the first column and first row. Because $S'(\rho)$ depends on the scattering direction through Eq. (1), both p_i and p_s are implicit functions of (θ, ϕ) :

$$\begin{aligned} p_i &= \frac{1}{S'_{00}} \sqrt{(S'_{10})^2 + (S'_{20})^2 + (S'_{30})^2}, \\ p_s &= \frac{1}{S'_{00}} \sqrt{(S'_{01})^2 + (S'_{02})^2 + (S'_{03})^2}. \end{aligned} \quad (2)$$

Polarization symmetry is defined by the condition that the signal and idler degrees of polarization are equal for all scattering directions,

$$p_i(\theta, \phi) = p_s(\theta, \phi). \quad (3)$$

Using Eq. (2), and noting that both expressions share the same normalization factor S'_{00} , the condition in Eq. (3) is equivalent, after canceling the common factor and squaring both sides, to

$$(S'_{10})^2 + (S'_{20})^2 + (S'_{30})^2 - (S'_{01})^2 - (S'_{02})^2 - (S'_{03})^2 = 0. \quad (4)$$

Using Eq. (4), we therefore define the Stokes-space quantity

$$\begin{aligned} \mathcal{S}_{\text{pol}}(\theta, \phi) &\equiv (S'_{10})^2 + (S'_{20})^2 + (S'_{30})^2 \\ &\quad - (S'_{01})^2 - (S'_{02})^2 - (S'_{03})^2, \end{aligned} \quad (5)$$

so that polarization symmetry is equivalent to the requirement

$$\mathcal{S}_{\text{pol}}(\theta, \phi) = 0, \quad (\theta, \phi) \in [0, \pi] \times [0, 2\pi]. \quad (6)$$

Evaluating Eq. (5) using the results obtained from Eq. (1) yields the decomposition

$$\mathcal{S}_{\text{pol}}(\theta, \phi) = \sum_{j=1}^6 c_j(\rho, \phi) h_j(\theta). \quad (7)$$

The six angular functions $h_j(\theta)$ are built from $t_{11}, t_{12}, t_{33}, t_{44}$. A key structural point is that the set $\{h_j(\theta)\}_{j=1}^6$ is linearly independent for any fixed incident energy $E_0 > 0$ on the physical domain. Appendix A establishes this by clearing the Klein-Nishina denominator, expanding the resulting polynomials in $x = \cos \theta$, and exhibiting a nonzero 6×6 minor with determinant proportional to $E_0^{11}(E_0 + 1)^3$. Therefore, Eq. (7) can vanish for all θ only if

$$c_j(\rho, \phi) = 0, \quad j = 1, \dots, 6,$$

for all ϕ .

The conditions $c_j(\rho, \phi) = 0$ simplify by grouping independent ϕ functions (e.g., $\cos 2\phi$ and $\sin 2\phi$, or $1, e^{4i\phi}, e^{8i\phi}$), which forces separate vanishing of the corresponding coefficients. Appendixes B and C show that the coupled system can be solved sequentially and reduces to compact structural constraints.

From c_1 , together with the normalization condition $\text{Tr}[\rho] = 1$, the diagonal entries of ρ are constrained to form pairs whose sums equal $\frac{1}{2}$. From c_2 , diagonal weights collapse to a one-parameter form and one coherence is eliminated. Collecting these relations gives

$$\begin{aligned} (\rho_{11}, \rho_{22}, \rho_{33}, \rho_{44}) &= (\rho_{11}, \tfrac{1}{2} - \rho_{11}, \tfrac{1}{2} - \rho_{11}, \rho_{11}), \\ \rho_{24} &= -\rho_{13}. \end{aligned} \quad (8)$$

From c_3 , linear independence of $\cos 2\phi$ and $\sin 2\phi$ enforces

$$\rho_{34} = -\rho_{12}. \quad (9)$$

From c_4 (equivalently c_5), grouping $\{1, e^{4i\phi}, e^{8i\phi}\}$ yields the pair

$$\rho_{12}^2 + \rho_{14}\rho_{23}^* = 0, \quad 2|\rho_{12}|^2 + |\rho_{14}|^2 + |\rho_{23}|^2 = \tfrac{1}{4}. \quad (10)$$

The remaining constraint c_6 becomes an identity once Eqs. (8)–(10) hold.

Positive semidefiniteness imposes principal-minor bounds on the (1, 4) and (2, 3) sub-blocks of ρ , and when these are combined with Eq. (10), the bounds are forced to saturation. This yields

$$|\rho_{14}| = \rho_{11}, \quad |\rho_{23}| = \tfrac{1}{2} - \rho_{11}, \quad |\rho_{12}| = A(\rho_{11}), \quad (11)$$

where

$$A(\rho_{11}) = \sqrt{\rho_{11}(\tfrac{1}{2} - \rho_{11})}, \quad 0 \leq \rho_{11} \leq \tfrac{1}{2}. \quad (12)$$

The function $A(\rho_{11})$ defines the upper semicircle $(\rho_{11} - \frac{1}{4})^2 + A^2 = (\frac{1}{4})^2$ in the (ρ_{11}, A) plane and satisfies $0 \leq A \leq \frac{1}{4}$ (see Fig. 4 in Appendix C). Using Eq. (11), write $\rho_{14} = \rho_{11}e^{i\alpha_{14}}$ and $\rho_{23} = (\frac{1}{2} - \rho_{11})e^{i\alpha_{23}}$ with phases $\alpha_{14}, \alpha_{23} \in [0, 2\pi)$. Equation (10) then fixes ρ_{12} up to a global parity branch. With $\Delta\alpha = \alpha_{14} - \alpha_{23}$ and $\Sigma\alpha = \alpha_{14} + \alpha_{23}$, the complete family is

$$\rho \rightarrow \rho^{(\pm)} = \begin{pmatrix} \rho_{11} & \pm iAe^{i\Delta\alpha/2} & \pm iAe^{i\Sigma\alpha/2} & \rho_{11}e^{i\alpha_{14}} \\ \mp iAe^{-i\Delta\alpha/2} & \frac{1}{2} - \rho_{11} & (\frac{1}{2} - \rho_{11})e^{i\alpha_{23}} & \mp iAe^{i\Sigma\alpha/2} \\ \mp iAe^{-i\Sigma\alpha/2} & (\frac{1}{2} - \rho_{11})e^{-i\alpha_{23}} & \frac{1}{2} - \rho_{11} & \mp iAe^{i\Delta\alpha/2} \\ \rho_{11}e^{-i\alpha_{14}} & \pm iAe^{-i\Sigma\alpha/2} & \pm iAe^{-i\Delta\alpha/2} & \rho_{11} \end{pmatrix}, \quad \begin{aligned} A &= \sqrt{\rho_{11}(\frac{1}{2} - \rho_{11})}, \\ \rho_{11} &\in [0, 1/2], \\ \Delta\alpha &= \alpha_{14} - \alpha_{23}, \\ \Sigma\alpha &= \alpha_{14} + \alpha_{23}, \\ \alpha_{14}, \alpha_{23} &\in [0, 2\pi). \end{aligned} \quad (13)$$

Imposing polarization symmetry for all azimuthal angles restricts the general two-qubit density matrix ρ to the family $\rho^{(\pm)}$ given in Eq. (13). The solutions are parametrized by a discrete $\pm$ branch and the continuous variables $(\rho_{11}, \alpha_{14}, \alpha_{23})$. The derivation hinges on two reusable mathematical structures: (1) the separation $\mathcal{S}_{\text{pol}} = \sum_j c_j h_j$ together with linear independence of $\{h_j\}$ and (2) elimination of ϕ dependence by enforcing vanishing of coefficients multiplying independent ϕ functions. The intermediate expressions for $h_j(\theta)$ and the rank-minor calculation are given in Appendix A, and the intermediate expressions for $c_j(\rho, \phi)$ and the solution set are given in Appendixes B and C.

The two branches are not physically distinct. They are related by a local-unitary transformation acting independently on each subsystem,

$$\rho^{(\pm)} = (\sigma_z \otimes \sigma_z) \rho^{(\mp)} (\sigma_z \otimes \sigma_z).$$

Direct evaluation shows $\text{Tr}[(\rho^{(\pm)})^2] = 1$, so the symmetry conditions restrict the solutions to pure states. Each branch may thus be written as a rank-1 projector, $\rho^{(\pm)} = |\psi_{\pm}\rangle\langle\psi_{\pm}|$.

Introducing complex probability amplitudes

$$\begin{aligned} a &= \sqrt{\rho_{11}}, \quad b = -i\sqrt{\frac{1}{2} - \rho_{11}} e^{-i\Delta\alpha/2}, \\ c &= -i\sqrt{\frac{1}{2} - \rho_{11}} e^{-i\Sigma\alpha/2}, \quad d = \sqrt{\rho_{11}} e^{-i\alpha_{14}}, \end{aligned} \quad (14)$$

the corresponding pure states take the form

$$|\psi_{\pm}\rangle = a|RR\rangle \pm b|RL\rangle \pm c|LR\rangle + d|LL\rangle. \quad (15)$$

Applying the spin-flip operation [22], the concurrence evaluates directly to $C(\psi_{\pm}) = 2|ad - bc| = 1$, establishing that the global polarization-symmetry condition uniquely selects maximally entangled pure two-photon states.

III. EQUIVALENCE TO THE BELL ORBIT

The previous section established that both branches $|\psi_{\pm}\rangle$ are pure states with concurrence $C = 1$. This section makes the associated Bell-state structure explicit by exhibiting the Schmidt decomposition and the corresponding local-unitary equivalence.

For each branch, introduce the 2×2 coefficient matrix

$$\Lambda_{\pm} = \begin{pmatrix} a & \pm b \\ \pm c & d \end{pmatrix}.$$

Using the probability amplitude definitions in Eq. (14), direct evaluation gives

$$\Lambda_{\pm} \Lambda_{\pm}^{\dagger} = \frac{1}{2} \mathbb{I}_2.$$

The two singular values are therefore equal to $1/\sqrt{2}$, so the Schmidt coefficients of $|\psi_{\pm}\rangle$ are

$$(1/\sqrt{2}, 1/\sqrt{2}).$$

This is consistent with the concurrence result obtained in Sec. II and makes the Bell-state structure explicit.

Since $\Lambda_{\pm}$ is proportional to a unitary matrix, define

$$U_{\pm} = \sqrt{2} \Lambda_{\pm}.$$

The corresponding states given in Eq. (15) may then be written as

$$|\psi_{\pm}\rangle = (U_{\pm} \otimes \mathbb{I}_2)|\Phi^{+}\rangle,$$

where $|\Phi^{+}\rangle = (|RR\rangle + |LL\rangle)/\sqrt{2}$ is a standard Bell state. Equivalently, Eq. (15) can be written as

$$|\psi_{\pm}\rangle = \frac{1}{\sqrt{2}}|u_r^{\pm}\rangle \otimes |R\rangle + \frac{1}{\sqrt{2}}|u_l^{\pm}\rangle \otimes |L\rangle,$$

with

$$|u_r^{\pm}\rangle = U_{\pm}|R\rangle = \sqrt{2}a|R\rangle \pm \sqrt{2}c|L\rangle,$$

$$|u_l^{\pm}\rangle = U_{\pm}|L\rangle = \pm\sqrt{2}b|R\rangle + \sqrt{2}d|L\rangle.$$

Unitarity of $U_{\pm}$ ensures that $|u_r^{\pm}\rangle$ and $|u_l^{\pm}\rangle$ form an orthonormal basis. Both branches therefore lie on the Bell orbit and differ only by local-unitary transformations acting on one subsystem.

An equivalent representation follows from the identity

$$(U \otimes \mathbb{I}_2)|\Phi^{+}\rangle = (\mathbb{I}_2 \otimes U^{\top})|\Phi^{+}\rangle,$$

which shows that the same Bell orbit can be generated by a local unitary acting on the second qubit instead.

Each Schmidt vector defines a pure single-qubit state and may be represented by a Bloch vector. As illustrated in Fig. 2, the states $|u_r^{\pm}\rangle$ and $|u_l^{\pm}\rangle$ correspond to antipodal points on the Bloch sphere, reflecting orthogonality and maximal entanglement.

The Schmidt basis is related to the circular polarization basis by a local-unitary rotation, and variation of the free phase parameters causes the associated Bloch axis to sweep the entire sphere. This behavior expresses the continuous




FIG. 2. Branch-dependent Schmidt basis states for the idler subsystem and fixed polarization basis states for the signal subsystem. In panel (a), the blue idler states $|u_r^\pm\rangle$ are paired with the blue signal state $|R\rangle$ in panel (b), while the magenta idler states $|u_l^\pm\rangle$ are paired with the magenta signal state $|L\rangle$ in panel (b). For each branch, $|u_r^\pm\rangle$ and $|u_l^\pm\rangle$ form an antipodal pair on the Bloch sphere, generated from $U_\pm = \sqrt{2}\Lambda_\pm$, so the + and - branches define two branch-dependent diameters. The two diameters are not generally parallel; the same-color cross-branch overlaps satisfy $\langle u_r^- | u_l^+ \rangle = \langle u_l^- | u_r^+ \rangle = 4\rho_{11} - 1$, so the relative angular separation is controlled by ρ_{11} .

local-unitary freedom characteristic of maximally entangled two-qubit states. This explicit construction clarifies why the polarization-symmetry condition derived earlier is satisfied only for these states.

Transforming $\rho^{(\pm)}$ to the Stokes representation and evolving it according to Eq. (1), the degrees of polarization of the idler and signal beams, as defined in Eq. (2), are given by

$$p_i = p_s = \frac{t_{12}}{t_{11}} = \frac{\sin^2 \theta}{1 + \cos^2 \theta + (E_0 - E(\theta))(1 - \cos \theta)},$$

independent of the azimuthal angle. The incident energy E_0 is the signal-photon energy in units of mc^2 , and $E(\theta)$ is the scattered photon energy given by the Compton relation

$$E(\theta) = \frac{E_0}{1 + E_0(1 - \cos \theta)}. \quad (16)$$

Figure 3 illustrates this equality for several incident photon energies.

IV. CONCLUSION

Under the single-Compton-scattering conditions considered here, polarization symmetry between idler and signal beams arises exclusively for maximally entangled Bell states. The result established in this work therefore applies to the symmetry condition $p_i = p_s$ within the model studied here.

From an operational standpoint, the symmetry can be tested through coincidence-conditioned reconstruction of the signal and idler degrees of polarization as functions of scattering direction. Work is ongoing to formulate this reconstruction as an explicit measurement protocol for the symmetry. Verification for an unknown state therefore requires angularly resolved coincidence data over the relevant domain.

More broadly, the result shows that entanglement allows polarization information associated with the coincidence-conditioned idler ensemble to be reconstructed from coin-




FIG. 3. p_i and p_s shown to be equal for states on the Bell orbit, plotted vs the scattering angle θ for selected incident energies, $E_0 = 0.511, 0.2, 1.5$, and 5 MeV. The black solid curve traces the peak value from Stokes vectors reconstructed for a fixed (θ, ϕ) direction.

cidence data without direct interaction with the idler beam, thereby preserving the intrinsic energy resolution of that beam.

ACKNOWLEDGMENT

This work was supported by the Japan Society for the Promotion of Science through JSPS KAKENHI Grant No. JP22H05022.

DATA AVAILABILITY

The data that support the findings of this article are not publicly available. The data are available from the authors upon reasonable request.

APPENDIX A: LINEAR INDEPENDENCE OF THE ANGULAR FUNCTIONS $h_j(\theta)$

This Appendix shows that the six functions $h_j(\theta)$ in Eq. (7) entering the polarization-symmetry analysis form a linearly independent set for any fixed incident energy $E_0 > 0$ in the physical range. The argument proceeds by rewriting $h_j(\theta)$ as polynomials in $\cos \theta$ after clearing the Klein-Nishina denominator.

Introduce the shorthand

$$x \equiv \cos \theta, \quad D(x) \equiv 1 + E_0(1 - x), \quad (A1)$$

so that the scattered energy given in Eq. (16) is expressed as

$$E(x) = \frac{E_0}{D(x)}.$$

For $x \in [-1, 1]$ and $E_0 > 0$, one has $D(x) \geq 1 > 0$, so $D(x)$ never vanishes on the physical domain. Using the expressions of Table I in Ref. [6], the Compton matrix elements are written

as follows:

$$\begin{aligned} t_{11}(x) &= \frac{1}{D(x)} [(1+x^2)D(x) + E_0^2(1-x)^2], \\ t_{12}(x) &= 1-x^2, \quad t_{33}(x) = 2x, \\ t_{44}(x) &= \frac{1}{D(x)} [2xD(x) + E_0^2(1-x)^2x]. \end{aligned} \quad (\text{A2})$$

In terms of the matrix elements given in Eq. (A2), the six angular functions are

$$\begin{aligned} h_1(\theta) &= t_{44}^2, \quad h_2(\theta) = -t_{11}^2, \\ h_3(\theta) &= t_{11}t_{12}, \quad h_4(\theta) = (2-t_{12})^2, \\ h_5(\theta) &= t_{33}^2, \quad h_6(\theta) = t_{12}(2-t_{12}), \end{aligned} \quad (\text{A3})$$

which, in terms of x , are functions of x and $D(x)$.

The only denominators arising in Eq. (A2) are from the t_{11} and t_{44} elements, and always as powers of $D(x)$. Multiplying by $D(x)^2$ removes all denominators. Using Eqs. (A1) and (A3), define

$$f_j(x) \equiv h_j(\theta) D(x)^2, \quad j = 1, \dots, 6. \quad (\text{A4})$$

Each $f_j(x)$ is a polynomial in x of degree at most six with coefficients that are polynomials in E_0 . Using Eq. (A2) to evaluate Eq. (A3), one observes, for example, that the purely polynomial functions h_j are

$$h_4(\theta) = (1+x^2)^2, \quad h_5(\theta) = 4x^2, \quad h_6(\theta) = 1-x^4, \quad (\text{A5})$$

so that evaluating Eq. (A4) given Eq. (A5) for $f_4(x)$, $f_5(x)$, and $f_6(x)$ gives

$$\begin{aligned} f_4(x) &= (1+x^2)^2 D(x)^2, \quad f_5(x) = 4x^2 D(x)^2, \\ f_6(x) &= (1-x^4) D(x)^2. \end{aligned}$$

Each $f_j(x)$ in Eq. (A4) can be expanded in the monomial basis $\{1, x, x^2, x^3, x^4, x^5, x^6\}$ as

$$f_j(x) = \sum_{k=0}^6 a_{jk}(E_0) x^k, \quad j = 1, \dots, 6, \quad (\text{A6})$$

where the coefficients $a_{jk}(E_0)$ are explicit polynomials in E_0 . Ordering the columns by the monomials $\{1, x, x^2, x^3, x^4, x^5, x^6\}$ then gives the explicit 6×7 coefficient matrix denoted as $A(E_0)$ such that

$$A(E_0) = \begin{pmatrix} a_{10} & a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} \\ a_{20} & a_{21} & a_{22} & a_{23} & a_{24} & a_{25} & a_{26} \\ a_{30} & a_{31} & a_{32} & a_{33} & a_{34} & a_{35} & a_{36} \\ a_{40} & a_{41} & a_{42} & a_{43} & a_{44} & a_{45} & a_{46} \\ a_{50} & a_{51} & a_{52} & a_{53} & a_{54} & a_{55} & a_{56} \\ a_{60} & a_{61} & a_{62} & a_{63} & a_{64} & a_{65} & a_{66} \end{pmatrix}, \quad (\text{A7})$$

with entries of Eq. (A7) given by

$$\begin{aligned} a_{10} &= 0, & a_{20} &= -E_0^4 - 2E_0^3 - 3E_0^2 - 2E_0 - 1, & a_{30} &= E_0^3 + 2E_0^2 + 2E_0 + 1, \\ a_{11} &= 0, & a_{21} &= 4E_0^4 + 6E_0^3 + 6E_0^2 + 2E_0, & a_{31} &= -3E_0^3 - 4E_0^2 - 2E_0, \\ a_{12} &= E_0^4 + 4E_0^3 + 8E_0^2 + 8E_0 + 4, & a_{22} &= -6E_0^4 - 8E_0^3 - 7E_0^2 - 4E_0 - 2, & a_{32} &= 2E_0^3 + E_0^2, \\ a_{13} &= -4E_0^4 - 12E_0^3 - 16E_0^2 - 8E_0, & a_{23} &= 4E_0^4 + 8E_0^3 + 8E_0^2 + 4E_0, & a_{33} &= 2E_0^3 + 2E_0^2, \\ a_{14} &= 6E_0^4 + 12E_0^3 + 8E_0^2, & a_{24} &= -E_0^4 - 6E_0^3 - 5E_0^2 - 2E_0 - 1, & a_{34} &= -3E_0^3 - 2E_0^2 - 2E_0 - 1, \\ a_{15} &= -4E_0^4 - 4E_0^3, & a_{25} &= 2E_0^3 + 2E_0^2 + 2E_0, & a_{35} &= E_0^3 + 2E_0^2 + 2E_0, \\ a_{16} &= E_0^4, & a_{26} &= -E_0^2, & a_{36} &= -E_0^2, \\ a_{40} &= E_0^2 + 2E_0 + 1, & a_{50} &= 0, & a_{60} &= E_0^2 + 2E_0 + 1, \\ a_{41} &= -2E_0^2 - 2E_0, & a_{51} &= 0, & a_{61} &= -2E_0^2 - 2E_0, \\ a_{42} &= 3E_0^2 + 4E_0 + 2, & a_{52} &= 4E_0^2 + 8E_0 + 4, & a_{62} &= E_0^2, \\ a_{43} &= -4E_0^2 - 4E_0, & a_{53} &= -8E_0^2 - 8E_0, & a_{63} &= 0, \\ a_{44} &= 3E_0^2 + 2E_0 + 1, & a_{54} &= 4E_0^2, & a_{64} &= -E_0^2 - 2E_0 - 1, \\ a_{45} &= -2E_0^2 - 2E_0, & a_{55} &= 0, & a_{65} &= 2E_0^2 + 2E_0, \\ a_{46} &= E_0^2, & a_{56} &= 0, & a_{66} &= -E_0^2. \end{aligned}$$

The vector $F(x)$ of the polynomials $f_j(x)$ given in Eq. (A4) can be written compactly as

$$F(x) \equiv A(E_0)X(x) = \begin{pmatrix} f_1(x) \\ f_2(x) \\ f_3(x) \\ f_4(x) \\ f_5(x) \\ f_6(x) \end{pmatrix} = D(x)^2 \begin{pmatrix} h_1(x) \\ h_2(x) \\ h_3(x) \\ h_4(x) \\ h_5(x) \\ h_6(x) \end{pmatrix},$$

where $D(x) > 0$ and $X(x) = (1, x, x^2, x^3, x^4, x^5, x^6)^\top$.

Linear dependence of the set $\{f_j\}_{j=1}^6$ in Eq. (A4) means the existence of constants $c_1, \dots, c_6$, not all zero, such that

$$\sum_{j=1}^6 c_j f_j(x) = 0, \quad \text{for all } x. \quad (\text{A8})$$

Substituting the expression for the polynomials $f_j(x)$ given in Eq. (A6) into Eq. (A8) gives

$$\sum_{j=1}^6 c_j \left(\sum_{k=0}^6 a_{jk}(E_0) x^k \right) = \sum_{k=0}^6 \left(\sum_{j=1}^6 c_j a_{jk}(E_0) \right) x^k = 0.$$

For this polynomial to vanish for all x , each coefficient must vanish,

$$\sum_{j=1}^6 c_j a_{jk}(E_0) = 0, \quad k = 0, \dots, 6.$$

In matrix form, this condition is

$$A(E_0)^\top c = 0,$$

where $c = (c_1, c_2, c_3, c_4, c_5, c_6)^\top$.

If $A(E_0)$ has row rank 6, then $A(E_0)^\top$ has column rank 6 and the nullspace of $A(E_0)^\top$ is trivial. This forces $c = 0$ and excludes any nontrivial linear dependence among the f_j . If the row rank were smaller than 6, a nonzero c would exist and the set $\{f_j\}$ would be linearly dependent.

The matrix $A(E_0)$ is 6×7 , so $\det A(E_0)$ is not defined. Rank is instead determined by the largest nonzero square minor. If any 6×6 submatrix of $A(E_0)$ has nonzero determinant, then

$$\text{rank } A(E_0) \geq 6.$$

Since $A(E_0)$ has only six rows, one always has $\text{rank } A(E_0) \leq 6$, and a single nonzero 6×6 minor forces $\text{rank } A(E_0) = 6$.

One convenient choice is the 6×6 submatrix $B(E_0)$ obtained by dropping the first column of the matrix $A(E_0)$ in Eq. (A7),

$$B(E_0) \equiv \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} & a_{15} & a_{16} \\ a_{21} & a_{22} & a_{23} & a_{24} & a_{25} & a_{26} \\ a_{31} & a_{32} & a_{33} & a_{34} & a_{35} & a_{36} \\ a_{41} & a_{42} & a_{43} & a_{44} & a_{45} & a_{46} \\ a_{51} & a_{52} & a_{53} & a_{54} & a_{55} & a_{56} \\ a_{61} & a_{62} & a_{63} & a_{64} & a_{65} & a_{66} \end{pmatrix},$$

whose determinant evaluates to

$$\det B(E_0) = 64 E_0^{11} (E_0 + 1)^3.$$

For any physical $E_0 > 0$, this determinant is nonzero, so $B(E_0)$ has rank 6 and hence $A(E_0)$ has rank 6.

For completeness, another choice of 6×6 minor also has nonvanishing determinant. Taking the columns corresponding to the monomials $\{1, x, x^2, x^3, x^4, x^5\}$ gives a 6×6 matrix $A_{\text{sub}}(E_0)$ whose determinant evaluates to

$$\det A_{\text{sub}}(E_0) = 64 E_0^8 (E_0 + 1)^3 (E_0 + 2)^3,$$

which is strictly nonzero for any $E_0 > 0$. Either $B(E_0)$ or $A_{\text{sub}}(E_0)$ is sufficient to establish $\text{rank } A(E_0) = 6$.

Full row rank of $A(E_0)$ implies that $A(E_0)^\top c = 0$ has only the trivial solution $c = 0$, so no nontrivial linear combination of the f_j vanishes. The polynomials f_j are therefore linearly independent, and the original angular functions h_j inherit this property.

Since $D(x)$ is strictly positive for all scattering angles $\theta \in [0, \pi]$ and $E_0 > 0$, multiplication by $D(x)^2$ does not introduce or remove any zeros of linear combinations. Linear independence of $\{f_j(x)\}_{j=1}^6$ as polynomials in x is therefore equivalent to linear independence of the original functions $\{h_j(\theta)\}_{j=1}^6$ as functions of the scattering angle θ , because

$$\begin{aligned} \sum_{j=1}^6 c_j f_j(x) &= \sum_{j=1}^6 c_j D(x)^2 h_j(x) \\ &= D(x)^2 \sum_{j=1}^6 c_j h_j(x) = 0, \quad \text{for all } x, \end{aligned}$$

implies, after division by $D(x)^2 > 0$,

$$\sum_{j=1}^6 c_j h_j(x) = 0, \quad \text{for all } x.$$

In summary, the determinant conditions above show that for any fixed $E_0 > 0$ the set $\{h_j(\theta)\}_{j=1}^6$ is linearly independent as a family of functions of θ .

APPENDIX B: CONSTRAINTS OF THE PREFACTORS c_1 – c_6

This Appendix derives the 4×4 density matrix ρ that yields polarization symmetry, defined by the condition $p_i = p_s$ for all azimuthal angles ϕ of the scattered signal photon, where p_i and p_s denote the degrees of polarization of the idler and signal beams, respectively, as defined in Eq. (2). The starting point is a general two-photon state ρ in the circular-polarization basis. For completeness, the relevant definitions and constraints are restated here.

The density matrix takes the symbolic form

$$\rho = \begin{pmatrix} \rho_{11} & \rho_{12} & \rho_{13} & \rho_{14} \\ \rho_{12}^* & \rho_{22} & \rho_{23} & \rho_{24} \\ \rho_{13}^* & \rho_{23}^* & \rho_{33} & \rho_{34} \\ \rho_{14}^* & \rho_{24}^* & \rho_{34}^* & \rho_{44} \end{pmatrix}, \quad (\text{B1})$$

expressed in the circular-polarization basis $\{|R, R\rangle, |R, L\rangle, |L, R\rangle, |L, L\rangle\}$, ordered from top to bottom and left to right, where $|R\rangle$ and $|L\rangle$ denote the right- and left-circular polarization states, respectively, following the convention used in Ref. [6]. Hermiticity enforces $\rho = \rho^\dagger$, normalization imposes $\text{Tr}[\rho] = 1$, and the diagonal entries satisfy $\rho_{mm} \in \mathbb{R}$.

Following the method of Abouraddy *et al.* [21], the density matrix ρ is mapped to Stokes parameters through

$$S_{\mu\nu} = \text{Tr}[(\sigma_\mu \otimes \sigma_\nu)\rho], \quad \mu, \nu = 0, 1, 2, 3,$$

where σ_0 is the 2×2 identity matrix and $(\sigma_1, \sigma_2, \sigma_3)$ are the Pauli matrices. The resulting matrix S , referred to here as the Stokes correlation matrix, is

$$\rho \mapsto S(\rho) = \begin{pmatrix} S_{00} & S_{01} & S_{02} & S_{03} \\ S_{10} & S_{11} & S_{12} & S_{13} \\ S_{20} & S_{21} & S_{22} & S_{23} \\ S_{30} & S_{31} & S_{32} & S_{33} \end{pmatrix}. \quad (\text{B2})$$

Here and throughout, the indices (0, 1, 2, 3) label the operator basis $(\sigma_0, \sigma_1, \sigma_2, \sigma_3)$ in this exact order.

The marginal Stokes vectors of subsystem 1 and subsystem 2 are obtained directly from the correlation matrix $S(\rho)$ of Eq. (B2). For a normalized input state such as ρ , the zeroth Stokes component satisfies

$$S_{00} = \text{Tr}[(\sigma_0 \otimes \sigma_0)\rho] = \text{Tr}[\rho] = 1.$$

The first column of $S(\rho)$ gives the Stokes vector of subsystem 1, and the first row gives that of subsystem 2,

$$|S_1\rangle = \begin{pmatrix} S_{00} \\ S_{10} \\ S_{20} \\ S_{30} \end{pmatrix}, \quad |S_2\rangle = \begin{pmatrix} S_{00} \\ S_{01} \\ S_{02} \\ S_{03} \end{pmatrix}. \quad (\text{B3})$$

The evolved Stokes matrix $S'(\rho)$ for the interaction geometry in Fig. 1 is [Eq. (31) of Ref. [17]]

$$S'(\rho) = S(\rho)M^\top(\phi)T_{\text{un}}^\top(\theta) = S(\rho)M(-\phi)T_{\text{un}}(\theta), \quad (\text{B4})$$

where the rotation matrix $M(\phi)$ represents a rotation by an angle 2ϕ about the S_{03} axis of the signal-photon Stokes vector in Eq. (B3) and is given by [23]

$$M(\phi) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos 2\phi & \sin 2\phi & 0 \\ 0 & -\sin 2\phi & \cos 2\phi & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

which implies $M^\top(\phi) = M(-\phi)$.

The unpolarized Compton-scattering Mueller matrix $T_{\text{un}}(\theta)$ is symmetric, $T_{\text{un}}(\theta) = T_{\text{un}}^\top(\theta)$, and takes the form

[Eq. (28) of Ref. [6]]

$$T_{\text{un}}(\theta) \propto \begin{pmatrix} t_{11} & t_{12} & 0 & 0 \\ t_{12} & 2 - t_{12} & 0 & 0 \\ 0 & 0 & t_{33} & 0 \\ 0 & 0 & 0 & t_{44} \end{pmatrix}.$$

In Fig. 1, θ denotes the Compton-scattering angle and ϕ denotes the azimuthal angle that specifies the scattering plane. The quantity $S'(\rho)$ is an unnormalized, intensity-weighted Stokes matrix associated with the selected Compton-scattering channel and geometry rather than the Stokes matrix of a trace-normalized output density matrix. A normalized output density matrix could be recovered through the inverse Stokes-to-density-matrix map [20,21], which includes the required renormalization.

Polarization symmetry is imposed by requiring that the signal and idler degrees of polarization be equal for all scattering directions,

$$p_i(\theta, \phi) = p_s(\theta, \phi).$$

Since the same factor S'_{00} appears in both expressions in Eq. (2), it cancels in the symmetry condition. After canceling the common factor and squaring both sides, define the Stokes-space function

$$\mathcal{S}_{\text{pol}}(\theta, \phi) \equiv (S'_{10})^2 + (S'_{20})^2 + (S'_{30})^2 - (S'_{01})^2 - (S'_{02})^2 - (S'_{03})^2. \quad (\text{B5})$$

Polarization symmetry is therefore equivalent to the requirement

$$\mathcal{S}_{\text{pol}}(\theta, \phi) = 0,$$

which is the condition stated in Eq. (6). This form keeps the constraint in the Stokes representation; a direct expression in terms of the density-matrix elements ρ_{mn} would be unwieldy. Once Eq. (B4) is evaluated, the matrix elements $S'_{\mu\nu}$ become explicit functions of the density-matrix elements in Eq. (B1), the scattering angle θ , and the azimuthal angle ϕ .

Substituting the explicit expressions for the Stokes parameters into Eq. (B5), and then rearranging and grouping terms according to the Compton-scattering matrix elements of $T_{\text{un}}(\theta)$ [Eq. (7) of Ref. [17]], the entries of the rotation matrix $M(\phi)$ [Eq. (4) of Ref. [17]], and the amplitudes ρ_{mn} [Eq. (B1)], gives six linearly independent functions $h_j(\theta)$ and corresponding prefactors $c_j(\rho, \phi)$. This produces the decomposition

$$\mathcal{S}_{\text{pol}}(\theta, \phi) = \sum_{j=1}^6 c_j(\rho, \phi) h_j(\theta), \quad (\text{B6})$$

where the $h_j(\theta)$ are analyzed in Appendix A and $c_j(\rho, \phi)$ are explicit functions of the density-matrix entries and the azimuthal angle. Polarization symmetry therefore requires Eq. (B6) to vanish for all $(\theta, \phi) \in [0, \pi] \times [0, 2\pi]$.

Requiring polarization symmetry for every azimuthal angle ϕ leads to the six conditions:

$$c_1 : 2(\rho_{22} + \rho_{44}) - 1 = 0, \quad (\text{B7})$$

$$c_2 : (2\rho_{33} + 2\rho_{44} - 1)^2 + 4|\rho_{13} + \rho_{24}|^2 = 0, \quad (\text{B8})$$

$$c_3 : \text{Re}(\rho_{12} + \rho_{34}) \cos(2\phi) - \text{Im}(\rho_{12} + \rho_{34}) \sin(2\phi) = 0, \quad (\text{B9})$$

$$c_4 : 1 - 4(|\rho_{14}|^2 + |\rho_{23}|^2) + 8 \text{Re}(\rho_{12}\rho_{34}^*) + 8 \text{Re}[e^{4i\phi}(\rho_{12}\rho_{34} - \rho_{14}\rho_{23}^*)] = 0, \quad (\text{B10})$$

$$c_5 : -1 + 4(|\rho_{12}|^2 + |\rho_{14}|^2 + |\rho_{23}|^2 + |\rho_{34}|^2) + 8 \text{Re}[e^{4i\phi}(\rho_{14}\rho_{23}^* - \rho_{12}\rho_{34})] = 0, \quad (\text{B11})$$

$$c_6 : e^{4i\phi} \{ (2\rho_{33} + 2\rho_{44} - 1)(\rho_{12} - \rho_{34}) - 2[(\rho_{13} + \rho_{24})\rho_{23}^* + \rho_{14}(\rho_{13} + \rho_{24})^*] \} \\ + [(2\rho_{33} + 2\rho_{44} - 1)(\rho_{12} - \rho_{34})^* - 2((\rho_{13} + \rho_{24})\rho_{14}^* + \rho_{23}(\rho_{13} + \rho_{24})^*)] = 0. \quad (\text{B12})$$

The prefactors $c_1, \dots, c_6$ are treated as a coupled system of constraints and are solved sequentially to determine the class of density matrices compatible with polarization symmetry.

APPENDIX C: SOLUTION SETS FOR c_1 – c_6

1. Diagonal relations from c_1 and the trace

Constraint c_1 in Eq. (B7) involves only the diagonal elements:

$$2(\rho_{22} + \rho_{44}) = 1.$$

Solving gives

$$\rho_{44} = \frac{1}{2} - \rho_{22}.$$

Combining this with the normalization condition $\text{Tr}[\rho] = 1$ gives

$$\rho_{11} + \rho_{22} + \rho_{33} + \rho_{44} = \rho_{11} + \rho_{22} + \rho_{33} + \left(\frac{1}{2} - \rho_{22}\right) \\ = \rho_{11} + \rho_{33} + \frac{1}{2} = 1.$$

This implies

$$\rho_{11} + \rho_{33} = \frac{1}{2}.$$

Therefore, c_1 and normalization yield

$$\rho_{33} = \frac{1}{2} - \rho_{11} \quad (\text{C1})$$

and

$$\rho_{44} = \frac{1}{2} - \rho_{22}. \quad (\text{C2})$$

In words, c_1 enforces that (ρ_{11}, ρ_{33}) and (ρ_{22}, ρ_{44}) each form a pair that sums to $\frac{1}{2}$.

a. Balancing diagonal weight and eliminating one coherence via c_2

Constraint c_2 in Eq. (B8) imposes

$$(2\rho_{33} + 2\rho_{44} - 1)^2 + 4|\rho_{13} + \rho_{24}|^2 = 0. \quad (\text{C3})$$

Both terms are non-negative, so each must vanish separately. Using Eq. (C1) for ρ_{33} and Eq. (C2) for ρ_{44} , the first term becomes

$$2\rho_{33} + 2\rho_{44} - 1 = 2\left(\frac{1}{2} - \rho_{11}\right) + 2\left(\frac{1}{2} - \rho_{22}\right) - 1 = 0,$$

which gives

$$\rho_{11} + \rho_{22} = \frac{1}{2}.$$

This implies

$$\rho_{22} = \frac{1}{2} - \rho_{11}. \quad (\text{C4})$$

With ρ_{22} expressed in terms of ρ_{11} , substitution of Eq. (C4) into Eq. (C2) yields

$$\rho_{44} = \frac{1}{2} - \rho_{22} = \frac{1}{2} - \left(\frac{1}{2} - \rho_{11}\right),$$

that is,

$$\rho_{44} = \rho_{11}. \quad (\text{C5})$$

The second term in Eq. (C3) gives

$$4|\rho_{13} + \rho_{24}|^2 = 0,$$

which implies

$$\rho_{24} = -\rho_{13}. \quad (\text{C6})$$

At this stage, the diagonal elements of ρ in Eq. (B1) can all be expressed in terms of ρ_{11} , so the diagonal has the structure

$$\{\rho_{11}, \rho_{22}, \rho_{33}, \rho_{44}\} \mapsto \{\rho_{11}, \frac{1}{2} - \rho_{11}, \frac{1}{2} - \rho_{11}, \rho_{11}\} \quad (\text{C7})$$

and one off-diagonal relation is fixed as

$$\rho_{24} = -\rho_{13}. \quad (\text{C8})$$

This relation is derived from c_2 and used in the following discussions for c_3, c_4, c_5 , and c_6 .

b. Constraint on c_3

Constraint c_3 in Eq. (B9) has the structure

$$\text{Re}(\rho_{12} + \rho_{34}) \cos(2\phi) - \text{Im}(\rho_{12} + \rho_{34}) \sin(2\phi) = 0.$$

Because $\cos(2\phi)$ and $\sin(2\phi)$ are linearly independent as functions of ϕ , the only way for c_3 to vanish for every ϕ is for both prefactors to vanish separately. This enforces

$$\text{Re}(\rho_{12} + \rho_{34}) = 0, \quad \text{Im}(\rho_{12} + \rho_{34}) = 0,$$

and hence

$$\rho_{34} = -\rho_{12}. \quad (\text{C9})$$

This relation is derived from c_3 and used in the following discussions for c_4 , c_5 , and c_6 .

c. Constraint on c_4

Using Eq. (C9) in Eq. (B10), and then multiplying by $e^{4i\phi}$ to collect powers of $e^{4i\phi}$, gives

$$4e^{8i\phi}(-\rho_{12}^2 - \rho_{14}\rho_{23}^*) + e^{4i\phi}(1 - 8|\rho_{12}|^2 - 4|\rho_{14}|^2 - 4|\rho_{23}|^2) - 4(\rho_{12}^{*2} + \rho_{23}\rho_{14}^*) = 0. \quad (C10)$$

This is a linear combination of $\{e^{8i\phi}, e^{4i\phi}, 1\}$, which are linearly independent as functions of ϕ . For Eq. (C10) to hold for all ϕ , each coefficient must vanish:

$$\begin{aligned} e^{8i\phi} : 4(-\rho_{12}^2 - \rho_{14}\rho_{23}^*) &= 0, \\ e^{4i\phi} : 1 - 8|\rho_{12}|^2 - 4|\rho_{14}|^2 - 4|\rho_{23}|^2 &= 0, \\ 1 : -4(\rho_{12}^{*2} + \rho_{23}\rho_{14}^*) &= 0. \end{aligned}$$

Dividing out common factors gives

$$\rho_{12}^2 + \rho_{14}\rho_{23}^* = 0, \quad (C11a)$$

$$2|\rho_{12}|^2 + |\rho_{14}|^2 + |\rho_{23}|^2 = \frac{1}{4}, \quad (C11b)$$

Condition (C11b) is the complex conjugate of Eq. (C11a). Independence of ϕ therefore enforces the pair

$$\rho_{12}^2 + \rho_{14}\rho_{23}^* = 0 \quad (C12)$$

and

$$2|\rho_{12}|^2 + |\rho_{14}|^2 + |\rho_{23}|^2 = \frac{1}{4}. \quad (C13)$$

2. Amplitude-phase relations and positive semidefinite saturation

Write the remaining coherences in magnitude-phase form where phases are needed:

$$\begin{aligned} \rho_{12} &= |\rho_{12}|e^{i\alpha_{12}}, \\ \rho_{14} &= |\rho_{14}|e^{i\alpha_{14}}, \\ \rho_{23} &= |\rho_{23}|e^{i\alpha_{23}}. \end{aligned}$$

From Eq. (C12),

$$\rho_{12}^2 + \rho_{14}\rho_{23}^* = 0 \implies \rho_{12}^2 = -\rho_{14}\rho_{23}^*. \quad (C14)$$

Taking magnitudes in Eq. (C14) gives

$$|\rho_{12}|^2 = |\rho_{14}||\rho_{23}|. \quad (C15)$$

Substituting Eq. (C15) into Eq. (C13) and rearranging yields

$$(|\rho_{14}| + |\rho_{23}|)^2 = \frac{1}{4} \implies |\rho_{14}| + |\rho_{23}| = \frac{1}{2}, \quad (C16)$$

since magnitudes are non-negative. Equation (C16) is equivalently

$$|\rho_{23}| = \frac{1}{2} - |\rho_{14}|. \quad (C17)$$

A phase relation follows from Eq. (C14) and the polar definitions:

$$|\rho_{12}|^2 e^{i2\alpha_{12}} = -|\rho_{14}||\rho_{23}|e^{i(\alpha_{14}-\alpha_{23})}. \quad (C18)$$

Taking square roots (keeping both branches) gives

$$\rho_{12} = \pm i \sqrt{|\rho_{14}||\rho_{23}|} e^{i(\alpha_{14}-\alpha_{23})/2}.$$

Using $|\rho_{23}| = \frac{1}{2} - |\rho_{14}|$ from Eq. (C17) leads to

$$\rho_{12} = \pm i \sqrt{|\rho_{14}|\left(\frac{1}{2} - |\rho_{14}|\right)} e^{i(\alpha_{14}-\alpha_{23})/2}. \quad (C19)$$

Taking arguments in Eq. (C18) (for $|\rho_{12}| > 0$) gives

$$\alpha_{12} = \frac{\pi}{2} + \frac{\alpha_{14} - \alpha_{23}}{2} \pmod{\pi}. \quad (C20)$$

To enforce positive semidefiniteness, consider principal 2×2 minors of ρ in Eq. (B1) together with Eq. (C5) and the diagonal structure in Eq. (C7). For the (1, 4) block

$$\begin{pmatrix} \rho_{11} & \rho_{14} \\ \rho_{14}^* & \rho_{11} \end{pmatrix},$$

the determinant condition implies

$$|\rho_{14}| \leq \rho_{11}. \quad (C21)$$

For the (2, 3) block

$$\begin{pmatrix} \frac{1}{2} - \rho_{11} & \rho_{23} \\ \rho_{23}^* & \frac{1}{2} - \rho_{11} \end{pmatrix},$$

the determinant condition implies

$$|\rho_{23}| \leq \frac{1}{2} - \rho_{11}. \quad (C22)$$

Combining Eq. (C22) with Eq. (C17) yields

$$|\rho_{14}| \geq \rho_{11}.$$

Together with Eq. (C21), this forces saturation:

$$|\rho_{14}| = \rho_{11}. \quad (C23)$$

Thus, ρ_{14} has the explicit polar form

$$\rho_{14} = \rho_{11}e^{i\alpha_{14}}.$$

Substituting Eq. (C23) into Eq. (C19) gives

$$\rho_{12} = \pm i \sqrt{\rho_{11}\left(\frac{1}{2} - \rho_{11}\right)} e^{i(\alpha_{14}-\alpha_{23})/2},$$

which is consistent with the phase relation in Eq. (C20). From Eq. (C17), one also finds

$$\rho_{23} = \left(\frac{1}{2} - \rho_{11}\right)e^{i\alpha_{23}}.$$

Here, α_{23} remains a free phase, while $|\rho_{23}|$ is fixed.

Collecting the amplitudes,

$$\begin{aligned} |\rho_{14}| &= \rho_{11}, \\ |\rho_{23}| &= \frac{1}{2} - \rho_{11}, \\ |\rho_{12}| &= \sqrt{\rho_{11}\left(\frac{1}{2} - \rho_{11}\right)}, \end{aligned} \quad (C24)$$

together with the phase relation in Eq. (C20) (for $|\rho_{12}| > 0$). For later use, define

$$A(\rho_{11}) \equiv \sqrt{\rho_{11}\left(\frac{1}{2} - \rho_{11}\right)}, \quad 0 \leq \rho_{11} \leq \frac{1}{2}. \quad (C25)$$

Squaring and completing the square in ρ_{11} gives

$$A^2 = \rho_{11}\left(\frac{1}{2} - \rho_{11}\right) = -\rho_{11}^2 + \frac{1}{2}\rho_{11} = \frac{1}{16} - \left(\rho_{11} - \frac{1}{4}\right)^2,$$




FIG. 4. Plot of $A(\rho_{11}) = \sqrt{\rho_{11}(1/2 - \rho_{11})}$ on the physical domain $0 \leq \rho_{11} \leq 1/2$. The point at $(1/4, 0)$ marks the center of the circle $(\rho_{11} - 1/4)^2 + A^2 = (1/4)^2$. The curve $A(\rho_{11})$ parametrizes the upper semicircle. The radial arrow indicates the maximum value $A_{\max} = 1/4$ at $\rho_{11} = 1/4$, which acts as the extremal coherence admitted by the density-matrix parametrization.

so

$$(\rho_{11} - \frac{1}{4})^2 + A^2 = (\frac{1}{4})^2,$$

which is the equation of a circle in the (ρ_{11}, A) plane with center $(1/4, 0)$ and radius $1/4$. On the physical domain $0 \leq \rho_{11} \leq 1/2$ and $A \geq 0$, the graph of $A(\rho_{11})$ is the upper semicircle as shown in Fig. 4.

At the end points, $A(0) = A(1/2) = 0$. The interior maximum is obtained by maximizing $A^2 = \rho_{11}(1/2 - \rho_{11})$:

$$\frac{d}{d\rho_{11}} A^2 = \frac{1}{2} - 2\rho_{11} = 0 \implies \rho_{11} = \frac{1}{4},$$

$$A_{\max} = A\left(\frac{1}{4}\right) = \sqrt{\frac{1}{4}\left(\frac{1}{2} - \frac{1}{4}\right)} = \frac{1}{4}.$$

The range is therefore

$$A(\rho_{11}) \in [0, \frac{1}{4}].$$

Because the positive-semidefinite bounds in Eqs. (C21) and (C22) are saturated, the corresponding 2×2 principal minors vanish. For a positive semidefinite matrix, this forces the associated rows and columns to be linearly dependent. The saturated $(1, 4)$ block gives

$$\rho_{24} = e^{i\alpha_{14}} \rho_{12}^*, \quad \rho_{34} = e^{i\alpha_{14}} \rho_{13}^*,$$

while the saturated $(2, 3)$ block gives

$$\rho_{13} = e^{i\alpha_{23}} \rho_{12}, \quad \rho_{34} = e^{-i\alpha_{23}} \rho_{24}.$$

Using

$$\rho_{12} = \pm i A e^{i(\alpha_{14} - \alpha_{23})/2},$$

one therefore obtains

$$\rho_{13} = \pm i A e^{i(\alpha_{14} + \alpha_{23})/2}, \quad \rho_{24} = \mp i A e^{i(\alpha_{14} + \alpha_{23})/2}.$$

The remaining coherences are then fixed up to a global parity branch. Adopting the convention of retaining the $\pm$

on ρ_{12} ,

$$\begin{aligned} \rho_{12} &= \pm i A e^{i\Delta\alpha/2}, & \rho_{34} &= \mp i A e^{i\Delta\alpha/2}, \\ \rho_{13} &= \pm i A e^{i\Sigma\alpha/2}, & \rho_{24} &= \mp i A e^{i\Sigma\alpha/2}, \end{aligned} \quad (\text{C26})$$

where

$$\Delta\alpha = \alpha_{14} - \alpha_{23}, \quad \Sigma\alpha = \alpha_{14} + \alpha_{23}. \quad (\text{C27})$$

The continuous freedom is captured by three real parameters $(\rho_{11}, \alpha_{14}, \alpha_{23})$ with $\rho_{11} \in [0, 1/2]$ and $\alpha_{14}, \alpha_{23} \in [0, 2\pi)$.

a. Constraint on c_5

Using Eq. (C9) in Eq. (B11), and grouping terms by their dependence on $e^{4i\phi}$, gives

$$\begin{aligned} e^{4i\phi} (\rho_{12}^2 + \rho_{14}\rho_{23}^*) + e^{-4i\phi} (\rho_{12}^{*2} + \rho_{23}\rho_{14}^*) \\ + (2|\rho_{12}|^2 + |\rho_{14}|^2 + |\rho_{23}|^2 - \frac{1}{4}) = 0. \end{aligned} \quad (\text{C28})$$

This is a linear combination of $\{e^{4i\phi}, e^{-4i\phi}, 1\}$, which are independent as functions of ϕ . For Eq. (C28) to hold for all ϕ , each coefficient must vanish:

$$e^{4i\phi} : \rho_{12}^2 + \rho_{14}\rho_{23}^* = 0,$$

$$e^{-4i\phi} : \rho_{12}^{*2} + \rho_{23}\rho_{14}^* = 0,$$

$$1 : 2|\rho_{12}|^2 + |\rho_{14}|^2 + |\rho_{23}|^2 - \frac{1}{4} = 0.$$

Thus, ϕ independence of c_5 enforces

$$\rho_{12}^2 + \rho_{14}\rho_{23}^* = 0,$$

$$\rho_{12}^{*2} + \rho_{23}\rho_{14}^* = 0,$$

$$2|\rho_{12}|^2 + |\rho_{14}|^2 + |\rho_{23}|^2 = \frac{1}{4}. \quad (\text{C29})$$

The relations in Eq. (C29) are the complex conjugate counterparts of the relations in Eq. (C11) that arise from the analysis of c_4 , so c_5 does not introduce any additional independent constraint.

b. Constraint on c_6

Finally, consider Eq. (B12). Using Eqs. (C1) and (C5), one obtains

$$2\rho_{33} + 2\rho_{44} - 1 = 2(\frac{1}{2} - \rho_{11}) + 2\rho_{11} - 1 = 0.$$

Using Eq. (C6), one also has

$$\rho_{13} + \rho_{24} = 0.$$

Every term in Eq. (B12) is therefore proportional either to $2\rho_{33} + 2\rho_{44} - 1$ or to $\rho_{13} + \rho_{24}$, and hence vanishes identically. Constraint c_6 supplies no further independent restriction within this solution sector.

The rearrangement of Eq. (B5) into the linearly independent functions $h_j(\theta)$, given in Eq. (B6), implies that $S_{\text{pol}}(\theta, \phi)$ can vanish for all scattering angles only if each prefactor $c_j(\rho, \phi)$ multiplying $h_j(\theta)$ vanishes separately. The analysis above shows that, for any fixed $E_0 > 0$, the resulting solution set is parametrized by ρ_{11} and two phases $(\alpha_{14}, \alpha_{23})$, together with a global parity branch, and that no further constraints arise from c_5 and c_6 beyond those already enforced by c_1 – c_4 .

Collecting all constraints from c_1 through c_6 , together with the amplitude and phase relations in Eqs. (C24)–(C27), shows that polarization symmetry restricts ρ to a two-branch family

parametrized by the real triple $(\rho_{11}, \alpha_{14}, \alpha_{23})$ with $0 \leq \rho_{11} \leq \frac{1}{2}$ and $\alpha_{14}, \alpha_{23} \in [0, 2\pi)$. Therefore, the density matrix takes

one of two forms distinguished by the global sign choice for ρ_{12} .

Branch + (the + sign is assigned to ρ_{12}):

$$\rho^{(+)} = \begin{pmatrix} \rho_{11} & +iAe^{i\Delta\alpha/2} & +iAe^{i\Sigma\alpha/2} & \rho_{11}e^{i\alpha_{14}} \\ -iAe^{-i\Delta\alpha/2} & \frac{1}{2} - \rho_{11} & (\frac{1}{2} - \rho_{11})e^{i\alpha_{23}} & -iAe^{i\Sigma\alpha/2} \\ -iAe^{-i\Sigma\alpha/2} & (\frac{1}{2} - \rho_{11})e^{-i\alpha_{23}} & \frac{1}{2} - \rho_{11} & -iAe^{i\Delta\alpha/2} \\ \rho_{11}e^{-i\alpha_{14}} & +iAe^{-i\Sigma\alpha/2} & +iAe^{-i\Delta\alpha/2} & \rho_{11} \end{pmatrix}.$$

Branch - (the - sign is assigned to ρ_{12}):

$$\rho^{(-)} = \begin{pmatrix} \rho_{11} & -iAe^{i\Delta\alpha/2} & -iAe^{i\Sigma\alpha/2} & \rho_{11}e^{i\alpha_{14}} \\ +iAe^{-i\Delta\alpha/2} & \frac{1}{2} - \rho_{11} & (\frac{1}{2} - \rho_{11})e^{i\alpha_{23}} & +iAe^{i\Sigma\alpha/2} \\ +iAe^{-i\Sigma\alpha/2} & (\frac{1}{2} - \rho_{11})e^{-i\alpha_{23}} & \frac{1}{2} - \rho_{11} & +iAe^{i\Delta\alpha/2} \\ \rho_{11}e^{-i\alpha_{14}} & -iAe^{-i\Sigma\alpha/2} & -iAe^{-i\Delta\alpha/2} & \rho_{11} \end{pmatrix}.$$

-
- [1] F. Lei, A. J. Dean, and G. L. Hills, Compton polarimetry in gamma-ray astronomy, *Space Sci. Rev.* **82**, 309 (1997).
- [2] M. H. L. Pryce and J. C. Ward, Angular correlation effects with annihilation radiation, *Nature (London)* **160**, 435 (1947).
- [3] B. C. Hiesmayr and P. Moskal, Witnessing entanglement in Compton scattering processes via mutually unbiased bases, *Sci. Rep.* **9**, 8166 (2019).
- [4] P. Caradonna, D. Reutens, T. Takahashi, S. Takeda, and V. Vegh, Probing entanglement in Compton interactions, *J. Phys. Commun.* **3**, 105005 (2019).
- [5] S. Shivashankara, Entanglement entropy of Compton scattering with a witness, *Can. J. Phys.* **101**, 757 (2023).
- [6] P. Caradonna, I. D'Amico, D. G. Jenkins, and D. P. Watts, Stokes-parameter representation for Compton scattering of entangled and classically correlated two-photon systems, *Phys. Rev. A* **109**, 033719 (2024).
- [7] P. Caradonna, Kinematic analysis of multiple Compton scattering in quantum-entangled two-photon systems, *Ann. Phys. (NY)* **470**, 169779 (2024).
- [8] M. Bała, W. Krzemień, B. C. Hiesmayr, J. Baran, K. Dulski, K. Klimaszewski, L. Raczyński, R. Y. Shopa, and W. Wiślicki, Probing arbitrary polarized photon pairs undergoing double Compton scatterings by a dedicated MC simulator validated with experimental data, *Eur. Phys. J. C* **85**, 1115 (2025).
- [9] C. S. Wu and I. Shakhov, The angular correlation of scattered annihilation radiation, *Phys. Rev.* **77**, 136 (1950).
- [10] H. Langhoff, Die Linearpolarisation der Vernichtungsstrahlung von Positronen, *Z. Phys.* **160**, 186 (1960).
- [11] L. R. Kasday, J. D. Ullman, and C. S. Wu, Angular correlation of Compton-scattered annihilation photons and hidden variables, *Nuovo Cimento B* **25**, 633 (1975).
- [12] D. P. Watts, J. Bordes, J. R. Brown, A. Cherlin, R. Newton, J. Allison, M. Bashkanov, N. Efthimiou, and N. A. Zachariou, Photon quantum entanglement in MeV regime and its application in PET imaging, *Nat. Commun.* **12**, 2646 (2021).
- [13] D. Abdurashitov, A. Baranov, D. Borisenko, F. Guber, A. Ivashkin, S. Morozov, S. Musin, A. Strizhak, I. Tkachev, V. Volkov, and B. Zhuikov, Setup of Compton polarimeters for measuring entangled annihilation photons, *J. Instrum.* **17**, P03010 (2022).
- [14] D. Kim, A. N. Rachman, U. Taisei, M. Uenomachi, K. Shimazoe, and H. Takahashi, Background reduction in PET by double Compton scattering of quantum entangled annihilation photons, *J. Instrum.* **18**, P07007 (2023).
- [15] J. Bordes, J. R. Brown, D. P. Watts, M. Bashkanov, K. Gibson, R. Newton, and N. Zachariou, First detailed study of the quantum decoherence of entangled gamma photons, *Phys. Rev. Lett.* **133**, 132502 (2024).
- [16] I. Tkachev, S. Musin, D. Abdurashitov, A. Baranov, F. Guber, A. Ivashkin, and A. Strizhak, Measuring the evolution of entanglement in Compton scattering, *Sci. Rep.* **15**, 6064 (2025).
- [17] P. Caradonna, Compton scattering mediated by quantum entanglement, *Phys. Rev. A* **111**, 053708 (2025).
- [18] K. Shimazoe and M. Uenomachi, Quantum sensing and imaging in the MeV regime of nuclear medicine, *Appl. Phys. Express* **19**, 040104 (2026).
- [19] Gyaprasad and R. Joshi, Tunable decoherence of quantum polarization states via birefringence-frequency coupling using liquid crystal, *Opt. Commun.* **608**, 132984 (2026).
- [20] P. Caradonna and K. Shimazoe, Entanglement, steering, and separability in Compton-scattered annihilation photons, *Phys. Rev. A* **112**, 032413 (2025).
- [21] A. F. Abouraddy, A. V. Sergienko, B. E. Saleh, and M. C. Teich, Quantum entanglement and the two-photon Stokes parameters, *Opt. Commun.* **201**, 93 (2002).
- [22] W. K. Wootters, Entanglement of formation of an arbitrary state of two qubits, *Phys. Rev. Lett.* **80**, 2245 (1998).
- [23] W. H. McMaster, Matrix representation of polarization, *Rev. Mod. Phys.* **33**, 8 (1961).