

Dissecting Rate-Limiting Processes in Biomolecular Condensate Exchange Dynamics

Ross Kliegman^{1,*}, Eli Kengmana², Rebecca Schulman^{2,3,4} and Yaojun Zhang^{1,5,†}

¹Department of Physics & Astronomy, *Johns Hopkins University*, Baltimore, Maryland 21218, USA

²Department of Chemistry, *Johns Hopkins University*, Baltimore, Maryland 21218, USA

³Department of Chemical & Biomolecular Engineering, *Johns Hopkins University*, Baltimore, Maryland 21218, USA

⁴Department of Computer Science, *Johns Hopkins University*, Baltimore, Maryland 21218, USA

⁵Department of Biophysics, *Johns Hopkins University*, Baltimore, Maryland 21218, USA



(Received 25 August 2025; accepted 10 June 2026; published 20 July 2026)

An increasing number of biomolecules have been shown to phase-separate into biomolecular condensates—membraneless subcellular compartments capable of regulating distinct biochemical processes within living cells. The speed with which they exchange components with the cellular environment can influence how fast biochemical reactions occur inside condensates and how fast condensates respond to environmental changes, thereby directly impacting condensate function. While fluorescence recovery after photobleaching (FRAP) experiments are routinely performed to measure this exchange timescale, it remains a challenge to distinguish the various physical processes limiting fluorescence recovery and identify each associated timescale. Here, we present a reaction-diffusion model for condensate exchange dynamics and show that such exchange can differ significantly from that of conventional liquid droplets due to the presence of a percolated molecular network, which gives rise to different mobility species in the dense phase. In this model, exchange can be limited by diffusion of either the high- or low-mobility species in the dense phase, diffusion in the dilute phase, or the attachment (detachment) of molecules to (from) the network at the surface or throughout the bulk of the condensate. Combining analytic derivations and numerical simulations, we quantify the contributions of these distinct physical processes to the overall exchange timescale and predict an experimentally testable scaling relationship between the exchange timescale and condensate size. Our theory suggests that the exchange dynamics can be accelerated via a pore-mediated pathway in which molecules pass through the pores of the meshwork and attach directly in the condensate interior. Notably, this pathway permits a parameter regime in which the exchange timescale becomes independent of condensate size, a scaling behavior that is supported by our FRAP measurements on a biosynthetic DNA nanostar system. Our work offers insight into the rate-limiting physical processes that can control condensate material exchange, with implications for natural and engineered systems.

DOI: [10.1103/h9cl-j5x5](https://doi.org/10.1103/h9cl-j5x5)

I. INTRODUCTION

Living cells organize their contents into distinct functional compartments. Beyond traditional membrane-bound organelles, subcellular structures can take the form of dynamic, liquidlike networks of molecules called “biomolecular condensates” [1,2]. These condensates are dense assemblies of distinct proteins and nucleic acids that are driven by multivalent interactions to segregate out of the intracellular milieu. They enable functions vital for life, including gene regulation [3–5], signal transduction [6–8], and stress response [9–11], and, when misregulated, they have been implicated in various

diseases, most notably neurodegeneration [12–14] and cancer [15–18]. Understanding how condensates form and evolve over time in cells can therefore deepen our physical understanding of emergent self-organization in biological systems and potentially inform human health.

The liquidlike nature of condensates is often critical for their biological function, as it enables exchange of material with the surrounding dilute phase. For instance, metabolic condensates, such as purinosomes [19,20], are enriched in enzymes, substrates, and other biomolecules involved in specific metabolic pathways [21]. Regulating metabolic activity in condensates requires not only that reactants can partition into them, but also that products can later escape. However, with viscosities orders of magnitude larger than conventional oil droplets [22], condensates are thought to experience slow internal diffusion, limiting the exchange dynamics. More broadly, the speed of material exchange can influence condensate responses to environmental changes, as well as the number, size, and spatial distribution of condensates via Ostwald ripening [23,24]. Collectively, these effects can impact condensate function, motivating a need

*Contact author: rkriegm1@jhu.edu

†Contact author: yaojunz@jhu.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

for tools to accurately measure and interpret the exchange dynamics.

The timescales of molecular exchange are commonly measured with an experimental technique known as fluorescence recovery after photobleaching (FRAP) [25–28]. In a typical FRAP experiment, fluorescently labeled molecules are photobleached within a region of interest (ROI) upon irradiation with a high-intensity laser. The fluorescence intensity in the ROI then recovers over time due to molecular exchange with the surroundings until constant intensity is eventually restored. Photobleaching can be performed on a subregion within a droplet, known as partial FRAP, or on an entire droplet, known as full FRAP. Whereas partial FRAP experiments predominantly measure internal mixing within the dense phase, exchange between dense and dilute phases can be probed using full FRAP. Exchange dynamics have been studied in a range of experimental condensate systems [6,10,29–32], and complementary theories have been developed to extract meaningful physical quantities from measured fluorescence recovery curves [33–36]. While it is typically assumed that the exchange dynamics are limited by molecular diffusion, recent studies suggest that condensate material exchange can also be limited by other physical processes owing to the complexity of molecular interactions [37–41], e.g., interface resistance [40,41].

The exchange dynamics of condensates are ultimately determined by the constituent biomolecules and their microscopic structures and interactions. These interactions often include a complex combination of electrostatic, π - π , cation- π , hydrophobic, hydrogen-bonding, and complementary domain-domain interactions [42,43]. It has been proposed that a range of molecules, from intrinsically disordered proteins with low-complexity sequences to associative proteins with multiple interaction domains, generally conform to a “sticker-spacer” architecture [44–46], in which “stickers” are residues, sequence motifs, or larger folded domains that form reversible physical cross-links driving phase separation, and “spacers” exclude volume and connect the stickers to form polymers.

In the sticker-spacer framework, it follows that phase-separating molecules often form dynamically restructuring networks that go beyond traditional liquid-liquid phase separation [Fig. 1(a)], sometimes referred to as “phase separation coupled to percolation” [47,48]. In the modified physical picture [Fig. 1(b)], attachment and detachment of molecules to and from the percolated network naturally give rise to different mobility populations within the condensate for the same type of molecule. The low-mobility population (referred to as “species 1”) represents molecules bound to the network, and the high-mobility population (referred to as “species 2”) represents freely diffusing molecules detached from the network.

Indeed, multiple mobility populations have been reported in the dense phase of an *in vitro* reconstituted postsynaptic density system [49] as well as in single-component A1-LCD condensates [50]. Specifically, in the postsynaptic density system, adaptive single-molecule tracking has revealed that molecules within condensates stochastically switch between confined and freely diffusing states, consistent with motion on and off a percolated molecular network [49]. In A1-LCD condensates, single-fluorogen imaging has revealed




FIG. 1. Schematics of a condensate in (a) the conventional model, which assumes uniform molecular mobility inside the condensate (depicted in gray), and (b) our proposed model, in which transient attachment to and detachment from the molecular network can give rise to multiple mobilities for the same molecule inside the condensate. Connected blue molecules are bound to the network, whereas individual pink molecules are freely diffusing. By attaching and detaching, the two mobility species can convert between one another with rates $k_{2 \rightarrow 1}$ and $k_{1 \rightarrow 2}$, respectively.

that slow fluorogens localize to nanoscale hydrophobic hubs, whereas faster fluorogens explore more weakly interacting regions of the condensate, supporting the existence of a spatially heterogeneous landscape for condensate dynamics [50]. Together, these studies demonstrate that even nominally single-component condensates can harbor rich internal dynamical heterogeneity, yet a theory to interpret such dynamic complexity has been lacking.

Here, we develop a minimal reaction-diffusion model that incorporates two mobility species in the dense phase. We show that exchange can be accelerated through a pore-mediated pathway, where molecules pass through meshwork pores and attach directly within the condensate interior. Using theory and simulations, we identify distinct rate-limiting processes with characteristic scaling relationships, and make measurements consistent with these predictions using FRAP experiments on DNA nanostar condensates.

II. RESULTS

A. A reaction-diffusion model for condensate exchange dynamics

To explore how the presence of two mobility species impacts the condensate exchange timescale, we first develop a reaction-diffusion model for a phase-separated system at equilibrium. Assuming spherical symmetry, we describe the recovery dynamics of a fully bleached condensate (equivalent to the exchange dynamics) by the following coupled reaction-diffusion equations:

$$\frac{\partial c_1}{\partial t} = \nabla \cdot \left[D_1 \left(\nabla c_1 - c_1 \frac{\nabla c_1^{\text{eq}}}{c_1^{\text{eq}}} \right) \right] + k(c_1^{\text{eq}} c_2 - c_2^{\text{eq}} c_1), \quad (1a)$$

$$\frac{\partial c_2}{\partial t} = \nabla \cdot \left[D_2 \left(\nabla c_2 - c_2 \frac{\nabla c_2^{\text{eq}}}{c_2^{\text{eq}}} \right) \right] - k(c_1^{\text{eq}} c_2 - c_2^{\text{eq}} c_1), \quad (1b)$$

where $c_1(r, t)$ and $c_2(r, t)$ are the bleached concentrations of species 1 and 2, respectively; $D_1(r)$ and $D_2(r)$ are their position-dependent diffusion coefficients; $c_1^{\text{eq}}(r)$ and $c_2^{\text{eq}}(r)$ are their equilibrium concentration profiles; and k is a parameter that encodes how fast molecules convert between species. The coordinate r is the distance from the center of the spherical condensate, and t denotes the time. We note

that no percolated network is present in the dilute phase, so $c_1(r, t) = c_1^{\text{eq}}(r) = 0$ there.

In Eqs. (1a) and (1b), the first terms on the right-hand side represent conventional Fickian diffusion in a concentration gradient, and the second terms represent excess chemical potentials that drive molecules toward nonuniform equilibrium concentration profiles [34]. The third and fourth terms account for mobility switching due to binding and unbinding with the network. Molecules can attach to the percolated network and lower their mobility with a rate $k_{2 \rightarrow 1}(r)$, and detach from the network and regain higher mobility with a rate $k_{1 \rightarrow 2}(r)$. Detailed balance requires that the fluxes of association ($c_2 k_{2 \rightarrow 1}$) and dissociation ($c_1 k_{1 \rightarrow 2}$) are equal at equilibrium, which allows us to characterize these rates in terms of a single parameter, $k(r) \equiv k_{2 \rightarrow 1}(r)/c_1^{\text{eq}}(r) = k_{1 \rightarrow 2}(r)/c_2^{\text{eq}}(r)$. For simplicity, we assume k to be a constant, independent of the distance from the center of the condensate. The association and dissociation fluxes are then $k c_1^{\text{eq}} c_2$ and $k c_2^{\text{eq}} c_1$, respectively. Since $c_1(r, t) = c_1^{\text{eq}}(r) = 0$ outside the condensate, there is no association or dissociation flux in this region, and the dynamics of free scaffold molecules are described by a single Eq. (1b) without the third and fourth terms on the right.

Analogous to a full FRAP experiment, we set the initial condition to be

$$c_i(r, 0) = \begin{cases} c_i^{\text{eq}}(r), & r \leq R \\ 0, & r > R \end{cases} \quad (2)$$

for a fully bleached droplet, where $i = 1, 2$, and R is the droplet radius. We impose no-flux boundary conditions to conserve total particle number in the system:

$$\left. \frac{\partial c_i(r, t)}{\partial r} \right|_{r=0} = \left. \frac{\partial c_i(r, t)}{\partial r} \right|_{r=R} = 0. \quad (3)$$

Upon solving for $c_1(r, t)$ and $c_2(r, t)$, we can obtain a normalized brightness curve $I(t)$ for the fraction of molecules inside the droplet that are unbleached at a time t :

$$I(t) = 1 - \frac{\int_0^R [c_1(r, t) + c_2(r, t)] r^2 dr}{\int_0^R [c_1^{\text{eq}}(r) + c_2^{\text{eq}}(r)] r^2 dr}. \quad (4)$$

Finally, the characteristic timescale τ of the exchange dynamics is identified by fitting $I(t)$ to an exponential function of the form $1 - e^{-t/\tau}$. We note that the system reaches equilibrium when $c_1(r, t)/c_1^{\text{eq}}(r) = c_2(r, t)/c_2^{\text{eq}}(r) = f_b$, where the constant f_b is the fraction of total molecules that are bleached. In most cases, $f_b \ll 1$, so that $I(t \rightarrow \infty) = 1$.

B. Quantifying the timescales of rate-limiting processes

1. Analytical derivations

A proxy for condensate material exchange, fluorescence recovery in a bleached droplet is a multistep process involving dilute-phase diffusion, network attachment and detachment, and dense-phase diffusion. We outline the rate-limiting steps of FRAP recovery in Fig. 2. First, unbleached molecules must diffuse through the dilute phase to reach the droplet surface. In the limit of low dilute-phase concentration, we derive the dilute-phase diffusion-limited timescale τ_{dil} shown in Eq. (5a). Next, in the absence of species 2 in the dense phase (i.e.,




FIG. 2. Schematic of rate-limiting processes in the exchange dynamics of biomolecular condensates. In order for fluorescence to recover in a bleached droplet, unbleached molecules have to diffuse in the dilute phase with a timescale τ_{dil} until they encounter the droplet, attach to the network at the interface with a timescale τ_{int} or within the droplet bulk with a timescale τ_{con} , and diffuse within the network as species 1 (blue) with a timescale $\tau_{1,\text{den}}$ or through the network mesh inside the droplet as species 2 (pink) with a timescale $\tau_{2,\text{den}}$. In the slow-attachment limit, the faster of the two attachment mechanisms (τ_{int} vs τ_{con}) controls the exchange speed, whereas in the slow dense-phase diffusion limit, the faster of the two dense-phase diffusive mechanisms ($\tau_{1,\text{den}}$ vs $\tau_{2,\text{den}}$) controls the exchange speed.

$c_{2,\text{den}} = 0$), unbleached molecules enter the droplet by attaching to the network at the droplet interface and subsequently diffuse into the bulk dense phase as species 1. In this case, if the interfacial attachment and detachment process is rate limiting, we derive the interface-limited timescale τ_{int} shown in Eq. (5b), whereas if dense-phase diffusion of species 1 is rate limiting, we derive the timescale $\tau_{1,\text{den}}$ shown in Eq. (5c):

$$\tau_{\text{dil}} = \frac{c_{1,\text{den}} R^2}{3 c_{2,\text{dil}} D_{2,\text{dil}}}, \quad (5a)$$

$$\tau_{\text{int}} = \frac{R}{3 k c_{2,\text{dil}} \delta_{\text{eff}}}, \quad (5b)$$

$$\tau_{1,\text{den}} = \frac{R^2}{\pi^2 D_{1,\text{den}}}. \quad (5c)$$

Here, δ_{eff} is the effective width of the droplet interface, and $c_{i,\text{dil}}$ and $D_{i,\text{dil}}$ ($c_{i,\text{den}}$ and $D_{i,\text{den}}$) are the equilibrium concentration and diffusion coefficient of species i in the dilute (dense) phase, respectively, with $i = 1, 2$. These timescales largely follow our previous results in Ref. [40]. Detailed derivations of each timescale are provided in the Supplemental Material [51].

In the presence of species 2 in the dense phase (i.e., $c_{2,\text{den}} > 0$), a pore-mediated mode of recovery emerges as unbleached molecules can now alternatively enter the droplet by passing through the pores of the network and diffusing into the dense phase as species 2, which then attach to and detach from the network throughout the bulk of the droplet (Fig. 2). If the conversion between species 1 and 2 is fast relative to the diffusion timescales (i.e., k is sufficiently large), the bleached concentrations c_1 and c_2 can be treated as having reached local equilibrium: $c_1^{\text{eq}}(r) c_2(r, t) = c_2^{\text{eq}}(r) c_1(r, t)$. Equations (1a) and (1b) then reduce to a single “effective diffusion” equation:

$$\frac{\partial c(r, t)}{\partial t} = \nabla \cdot \left[D(r) \left(\nabla c(r, t) - c(r, t) \frac{\nabla c^{\text{eq}}(r)}{c^{\text{eq}}(r)} \right) \right], \quad (6)$$

where $c(r, t) = c_1(r, t) + c_2(r, t)$ is the total bleached concentration profile, $c^{\text{eq}}(r) = c_1^{\text{eq}}(r) + c_2^{\text{eq}}(r)$ is the total equilibrium concentration profile, and $D(r) = [D_1(r)c_1^{\text{eq}}(r) + D_2(r)c_2^{\text{eq}}(r)]/c^{\text{eq}}(r)$ is the effective diffusivity profile. In this fast-conversion regime, if the dilute-phase diffusion is rate limiting, the recovery timescale reverts to τ_{dil} in Eq. (5a) with $c_{1,\text{den}}$ replaced by $c_{1,\text{den}} + c_{2,\text{den}}$. However, if the dense-phase diffusion is rate limiting, we derive a new recovery timescale,

$$\tau = \frac{R^2}{\pi^2 D_{\text{eff}}}, \quad (7)$$

where D_{eff} is the effective dense-phase diffusion coefficient [52,53]:

$$D_{\text{eff}} = \frac{c_{1,\text{den}}D_{1,\text{den}} + c_{2,\text{den}}D_{2,\text{den}}}{c_{1,\text{den}} + c_{2,\text{den}}}. \quad (8)$$

Since molecules can either attach and diffuse as species 1 or detach and diffuse as species 2, D_{eff} is a weighted sum of the two diffusion processes, with the weights given by their respective occupancies. By definition, species 2 molecules diffuse more rapidly than species 1 (i.e., $D_{2,\text{den}} \gg D_{1,\text{den}}$). Thus, the presence of species 2 molecules in the dense phase can speed up the exchange dynamics (i.e., $D_{\text{eff}} > D_{1,\text{den}}$). However, because molecules may spend much less time as species 2 than as species 1, the degree of acceleration depends on the specific parameter values. When diffusion as species 1 dominates (i.e., $D_{1,\text{den}} \gg D_{2,\text{den}}c_{2,\text{den}}/c_{1,\text{den}}$), the recovery timescale is comparable to $\tau_{1,\text{den}}$ in Eq. (5c). Conversely, when species 2 diffusion dominates, the acceleration is significant and the timescale becomes

$$\tau_{2,\text{den}} = \frac{c_{1,\text{den}}R^2}{\pi^2 c_{2,\text{den}}D_{2,\text{den}}}, \quad (9)$$

where we have assumed $c_{1,\text{den}} \gg c_{2,\text{den}}$ (justified below). Detailed derivations are provided in the Supplemental Material [51].

In the opposite limit, if the diffusive processes are much faster than the conversion, then the bleached molecules of species 2 inside the droplet will be quickly replaced by unbleached molecules diffusing in from the dilute phase. This leads to a fast and partial recovery, with the recovery amplitude determined by the fraction of species 2 molecules in the dense phase [$c_{2,\text{den}}/(c_{1,\text{den}} + c_{2,\text{den}})$]. In contrast, the bleached molecules of species 1 can only be replaced via a slow attachment and detachment process, where they exchange with unbleached molecules of species 2 at the droplet interface and throughout the droplet bulk. This leads to a slow mode of recovery following the initial fast mode until the droplet is fully recovered. We show an example of such two-step recovery at the end of the following section (Numerical Simulations). Here, we argue that, in many cases, species 1 is expected to be much more abundant than species 2 in the dense phase (i.e., $c_{1,\text{den}} \gg c_{2,\text{den}}$) because species 1 is bound to the network and therefore energetically favored over species 2. In this situation, the contribution of the first fast mode is negligible so that the recovery curve mainly reflects the second slow process. We will thus focus on the recovery timescale of this second conversion-limited mode in the following.

Since conversion is slower than diffusion, the bleached molecules of each species can be considered equilibrated with

respect to their equilibrium concentration profiles:

$$c_i(r, t) \approx \alpha_i(t)c_i^{\text{eq}}(r). \quad (10)$$

The time-dependent component of $c_i(r, t)$ is given by $\alpha_i(t)$, which asymptotically approaches the overall bleached fraction, f_b , for $i = 1, 2$. To derive the timescale of recovery, we first apply the divergence theorem to Eqs. (1a) and (1b), yielding

$$\frac{d}{dt} \int_0^\infty c_1 r^2 dr = \int_0^\infty k(c_1^{\text{eq}}c_2 - c_2^{\text{eq}}c_1)r^2 dr, \quad (11a)$$

$$\frac{d}{dt} \int_0^\infty c_2 r^2 dr = \int_0^\infty k(c_2^{\text{eq}}c_1 - c_1^{\text{eq}}c_2)r^2 dr. \quad (11b)$$

We then substitute the approximate forms of $c_1(r, t)$ and $c_2(r, t)$ [Eq. (10)] into Eq. (11), leading to a relaxation time for $\alpha_1(t)$ and $\alpha_2(t)$:

$$\tau = \frac{\int_0^\infty c_1^{\text{eq}}(r)r^2 dr}{\int_0^\infty k c_1^{\text{eq}}(r)c_2^{\text{eq}}(r)r^2 dr}, \quad (12)$$

which is the conversion-limited recovery timescale.

There are two contributions to the integral in the denominator of Eq. (12): the integration over the interface region and the integration throughout the bulk of the condensate. By separating them into two terms, we derive in the Supplemental Material [51] an explicit form of the conversion-limited timescale,

$$\tau = \frac{R}{3kc_{2,\text{dil}}\delta_{\text{eff}} + kc_{2,\text{den}}R}, \quad (13)$$

which combines the effects of network attachment both at the interface and in the bulk of the droplet. We note that we have assumed a constant conversion rate k at the interface and in the condensate bulk. If the conversion rates differed in these two regions, the region-specific rates k_{int} and k_{con} would replace k in the interface- and conversion-limited contributions to the denominator, respectively. Once again, the presence of species 2 molecules in the dense phase can speed up the exchange dynamics, with the degree of acceleration determined by the specific parameter values. When conversion at the interface dominates the recovery time (i.e., $c_{2,\text{dil}}\delta_{\text{eff}} \gg c_{2,\text{den}}R$), the recovery timescale is comparable to τ_{int} in Eq. (5b). Conversely, when conversion throughout the condensate bulk dominates, the acceleration becomes significant and we obtain the bulk conversion-limited timescale:

$$\tau_{\text{con}} = \frac{1}{kc_{2,\text{den}}}. \quad (14)$$

At this point, we have identified five rate-limiting processes for the exchange dynamics of droplets with two mobility species and derived analytical expressions for their corresponding timescales in Eqs. (5), (9), and (14). Each physical process has a distinct timescale that scales with droplet size differently. Intuitively, the diffusion-limited processes in both dense and dilute phases are associated with timescales that scale as R^2/D [Eqs. (5a), (5c), and (9)]. The prefactors $c_{1,\text{den}}/c_{2,\text{dil}}$ and $c_{1,\text{den}}/c_{2,\text{den}}$ in Eqs. (5a) and (9) account for replacing bleached molecules of concentration $c_{1,\text{den}}$ with unbleached molecules of concentrations $c_{2,\text{dil}}$ and $c_{2,\text{den}}$, respectively. The interfacial timescale [Eq. (5b)] accounts for

exchange of a volume of molecules ($\sim R^3$) over a surface ($\sim R^2$) and is therefore linear in R . Lastly, the bulk conversion-limited timescale [Eq. (14)] is independent of R , which arises due to slow detachment of bleached molecules throughout the bulk of the dense phase. Once detached, these molecules can quickly escape the droplet, allowing unbleached molecules to attach to the network. The recovery timescale is therefore limited by the lifetime of a molecule in the network: $1/k_{1 \rightarrow 2} = 1/(kc_{2,\text{den}})$.

Putting together the rate-limiting steps, we propose the following expression for the overall timescale of fluorescence recovery:

$$\tau = \frac{c_{1,\text{den}}R^2}{3c_{2,\text{dil}}D_{2,\text{dil}}} + \frac{R}{3kc_{2,\text{dil}}\delta_{\text{eff}} + kc_{2,\text{den}}R} + \frac{R^2}{\pi^2 D_{\text{eff}}} \\ = \tau_{\text{dil}} + (\tau_{\text{int}}^{-1} + \tau_{\text{con}}^{-1})^{-1} + (\tau_{1,\text{den}}^{-1} + \tau_{2,\text{den}}^{-1})^{-1}, \quad (15)$$

where, following diffusion in the dilute phase, two additional processes, network attachment and detachment and dense-phase diffusion, are required for total fluorescence recovery. Again, we have invoked the assumption that $c_{1,\text{den}} \gg c_{2,\text{den}}$, which is motivated by the lower energy attained when molecules are attached to the droplet network. Both network attachment and detachment and dense-phase diffusion involve a combination of two competing processes in parallel: molecules can attach and detach at the interface as well as in the bulk of the condensate, and they can diffuse as species 1 or species 2, the slower of which is neglected in favor of the faster. It is worth noting that, by setting $c_{2,\text{den}} = 0$ in Eq. (15), i.e., assuming a single mobility species inside the droplet, we recover results of our previous study [40]: $\tau = \tau_{\text{dil}} + \tau_{\text{int}} + \tau_{1,\text{den}}$. In particular, τ_{int} arises due to the droplet's "interface resistance," which was previously modeled with a phenomenological parameter κ but now acquires a clear physical meaning: τ_{int} is governed by the molecular attachment and detachment at the droplet interface. For $c_{2,\text{den}} > 0$, the emergence of a pore-mediated mode of recovery in Fig. 2 can speed up the exchange dynamics and leads to two previously unrecognized terms in the recovery time, τ_{con} and $\tau_{2,\text{den}}$, resulting in a complex dependence of τ on the droplet radius R . We demonstrate this complex dependence via numerical simulations and FRAP experiments on DNA nanostar droplets below.

2. Numerical simulations

In the previous section, we derived the timescale of fluorescence recovery by analytically solving the reaction-diffusion system described by Eqs. (1)–(4) in various limits. Here, we numerically verify these timescales and visualize the different FRAP signatures in each rate-limiting case. Consistent with equilibrium solutions of the Cahn-Hilliard equation [54], we first specify the functional form of the equilibrium concentration profiles with a sharp but smooth transition at the droplet interface ($r = R$) over a finite width $\sim l$ [Fig. 3(a)],

$$c_i^{\text{eq}}(r) = \frac{c_{i,\text{dil}} + c_{i,\text{den}}}{2} + \frac{c_{i,\text{dil}} - c_{i,\text{den}}}{2} \tanh \frac{r - R}{l}, \quad (16)$$




FIG. 3. Representative simulation in a dilute-phase diffusion-limited scenario (parameters from top row in Table I). (a) Equilibrium concentration profiles and (b) diffusivity profiles for species 1 and 2. (c) Simulated radial concentration profiles of the bleached molecules for a few illustrative times. (d) Simulated brightness curve with exponential fit using nonlinear least squares [56]. Simulations were performed with radial step size $dr = 20$ nm over a system size of $r_{\text{max}} = 30$ μm for 1000 recorded time points (the solver dynamically selects both the time step and order for time integration).

with $i = 1, 2$, and we assume a similar form for the diffusivity profiles [Fig. 3(b)],

$$D_i(r) = \frac{D_{i,\text{dil}} + D_{i,\text{den}}}{2} + \frac{D_{i,\text{dil}} - D_{i,\text{den}}}{2} \tanh \frac{r - R}{l}. \quad (17)$$

We then solve Eqs. (1a) and (1b) numerically under spherical symmetry, using the initial and boundary conditions given by Eqs. (2) and (3), respectively, except that the boundary at $r = +\infty$ is replaced by $r = r_{\text{max}}$ for the finite size of the system. These coupled partial differential equations were solved using the `pdepe` function in MATLAB, which employs finite-difference spatial discretization with a variable-step, variable-order solver for time integration [55]. The numerical solutions for $c_1(r, t)$ and $c_2(r, t)$ [Fig. 3(c)] are used to compute a brightness curve in accordance with Eq. (4), which is subsequently fitted to extract the recovery timescale [Fig. 3(d)].

We show an example where the FRAP recovery is limited by dilute-phase diffusion in Fig. 3. Guided by Eq. (15), we choose physiological parameters of condensate systems [22,57,58] that lead to $\tau \approx \tau_{\text{dil}}$ (Table I). The numerically extracted relaxation time $\tau = 15.1$ s of an $R = 1$ μm droplet is indeed close to the theoretical prediction of $\tau_{\text{dil}} = 13.3$ s. We repeat a similar procedure for various parameter sets in which the timescale of fluorescence recovery is limited by interfacial attachment and detachment, dense-phase diffusion of the low-mobility species, dense-phase diffusion of the high-mobility species, and attachment and detachment throughout the bulk of the condensate, totaling five cases. Simulated spatial fluorescence recovery profiles for each of these cases are shown in Fig. 4 with parameters listed in Table I. The two cases of dense-phase diffusion-limited recovery [Figs. 4(c) and 4(d)] can readily be distinguished by the pronounced gradient present due to unbleached molecules gradually diffusing

TABLE I. Parameters for numerical simulations of five rate-limiting cases.

Case	$c_{1,\text{den}}$ (μM)	$c_{1,\text{dil}}$ (μM)	$c_{2,\text{den}}$ (μM)	$c_{2,\text{dil}}$ (μM)	$D_{1,\text{den}}$ ($\mu\text{m}^2/\text{s}$)	$D_{1,\text{dil}}^a$ ($\mu\text{m}^2/\text{s}$)	$D_{2,\text{den}}$ ($\mu\text{m}^2/\text{s}$)	$D_{2,\text{dil}}$ ($\mu\text{m}^2/\text{s}$)	l (μm)	k ($\mu\text{M}^{-1} \text{s}^{-1}$)
$\tau \approx \tau_{\text{dil}}$	2000	0	0	5	0.1	0.1	1	10	0.1	1
$\tau \approx \tau_{\text{int}}$	1000	0	0	10	0.02	0.02	1	50	0.1	0.005
$\tau \approx \tau_{1,\text{den}}$	1000	0	0	10	0.02	0.02	1	50	0.1	1
$\tau \approx \tau_{2,\text{den}}$	1995	0	5	10	5×10^{-5}	5×10^{-5}	0.1	50	0.01	0.005
$\tau \approx \tau_{\text{con}}$	980	0	20	10	0.02	0.02	1	50	0.05	1×10^{-4}

^aSince there are no species 1 molecules in the dilute phase, we choose $D_{1,\text{dil}} = D_{1,\text{den}}$ to parametrize the diffusivity profile. This choice sets the diffusion coefficient of species 1 to be constant both inside the droplet and at its interface.

into the condensate and bleached ones diffusing out, whereas the remaining three cases all display a uniform recovery as the fast diffusion of molecules within the condensate quickly homogenizes the intensity profile. Details of the numerical implementation and fitting are provided in the Supplemental Material [51].

Based on Eqs. (5), (9), and (14), condensate recovery timescales are expected to follow different scaling laws in different rate-limiting cases. As shown in Fig. 5, the different timescales indeed scale differently with droplet size *in silico* as well. When diffusive processes are rate limiting, the scaling law is quadratic; when the interfacial flux is rate limiting, the scaling law is linear; and when the network attachment and detachment throughout the droplet is rate limiting, the scaling law is independent of droplet size. We note that the theoretical curves in Fig. 5 are predictions from Eqs. (5), (9), and (14)

with parameters in Table I, without any fitting to the simulated results. The deviations between theory and simulations in Figs. 5(c) and 5(d) reflect the presence of fast-decaying modes with non-negligible amplitudes in the diffusive processes [see Eq. (S40) in the Supplemental Material [51]]. Such deviations can be mitigated by fitting the simulated $I(t)$ curves in the time range $t \geq 0.5\tau$ to a function $1 - A \exp(-t/\tau)$, depicted as gray diamonds in Figs. 5(c) and 5(d).

Lastly, to provide a more complete picture, we present an example of a two-step recovery for a condensate with a significant high-mobility fraction (i.e., $c_{2,\text{den}} \sim c_{1,\text{den}}$). As discussed in the Analytical Derivations section, when diffusion is faster than conversion, the droplet recovery consists of an initial fast mode due to the rapid replacement of bleached species 2 molecules by unbleached molecules diffusing in from the dilute phase, followed by a second slower mode due to the gradual detachment of bleached species 1 molecules from the network. As shown in Fig. S1(b) [51], the exchange dynamics proceeds in stages, giving rise to a double-exponential recovery.

C. Application to a DNA nanostar system

Finally, we seek to employ our theory in an experimental system composed of DNA nanostars—a model system for investigating biomolecular condensation. Thanks to advances in DNA synthesis techniques, DNA nanostars offer highly programmable interactions: binding specificity and affinity can be tuned via the sequence and length of single-stranded overhangs, and valence via the number of arms. These features collectively enable a diverse range of phase behaviors [59–62]. Our DNA nanostars are composed of three arms of double-stranded DNA, each with a short tail of single-stranded DNA known as a “sticky end” due to its propensity to Watson-Crick base-pair with complementary strands [Fig. 6(a)] [63]. The sticky ends make these nanostars readily phase separable, and micrometer-sized droplets can be seen with confocal microscopy [Figs. 6(b) and S2]. Z-stack imaging also confirms that these droplets are mostly spherical (Fig. S2 [51]). Experiments were performed in a 40 mM Tris-Cl buffer solution at pH 7.9 containing 2.5 μM DNA nanostars, 1 mM dithiothreitol (DTT), 27.5 mM magnesium acetate, and 2.5 mM nucleoside triphosphates (NTPs) at 37 °C. Additional details about sequences and sample preparation are given in the Supplemental Material [51].

DNA nanostars form porous networks inside their condensates, with the volume fraction occupied by the nanostars on




FIG. 4. Simulated FRAP recovery profiles when fluorescence recovery is limited by (a) dilute-phase diffusion, (b) interfacial attachment and detachment, (c) dense-phase diffusion of species 1, (d) dense-phase diffusion of species 2, and (e) attachment and detachment throughout the bulk of the condensate. Green indicates fluorescent molecules and black indicates bleached molecules. Simulations were performed with radial step sizes $1/5$ of the interface width l , and the system size was $r_{\text{max}} = 30 \mu\text{m}$. 1000 time points were recorded over 4τ -long runtimes.




FIG. 5. Theoretical and simulated scaling laws show good agreement in each rate-limiting case: (a) dilute-phase diffusion-limited timescale scales with R^2 , (b) interface-limited timescale scales with R , (c) dense-phase diffusion-limited timescale (species 1) scales with R^2 , (d) dense-phase diffusion-limited timescale (species 2) scales with R^2 , and (e) bulk conversion-limited timescale is independent of R . Red curves, theoretical predictions from Eqs. (5), (9), and (14) using parameters from Table I; black crosses, simulation results for droplets of radii $R = 0.5, 1$, and $2 \mu\text{m}$. Gray diamonds in (c) and (d) are from the same simulation results fit to the modified function $1 - A \exp(-t/\tau)$ over a time range $t \geq 0.5\tau$, which disentangles the leading mode from the fast-decaying modes in diffusive processes.

the order of 1% or less [61,64]. This property makes them a prime system for observing the pore-mediated mode of recovery discussed above: nearby molecules may penetrate




FIG. 6. Experimental characterization of exchange dynamics in DNA nanostar condensates. (a) Nanostar schematic. The Y-shaped nanostar has 16-base pair (bp) double-stranded arms (shown in blue), sticky ends with the palindromic sequence 5'-GCTAGC-3' (shown in red), and 5'-TT-3' domains between each of the three arms at the nanostar center (shown in gray) that confer angular flexibility between arms. The nanostars were first annealed in a low-salt solution, then added to a higher-salt solution at 37 °C to incubate condensates, following the same protocol as in Ref. [63]. (b) Representative snapshots from a FRAP experiment of an $R = 1 \mu\text{m}$ nanostar droplet. (c) FRAP recovery curves for droplets of small sizes ($R \lesssim 1.5 \mu\text{m}$). (d) Exchange timescale versus droplet radius for all droplets, fitted with a shifted quadratic function of the form $\tau = a + bR^2$, where $a = 145.0 \pm 5.6 \text{ s}$ and $b = 14.1 \pm 0.7 \mu\text{m}^{-2} \text{ s}$. Error bars on the data points represent 3σ . A replicate under the same experimental conditions is shown in Fig. S3 [51], which confirms the same scaling behavior.

the droplet surface and diffuse freely within the droplet before attaching to the network. If this were the dominant recovery mechanism, we expect to observe conversion-limited recovery for small droplets, transitioning to diffusion-limited recovery for large droplets. Upon performing FRAP on nanostar droplets of varying sizes [Fig. 6(b)], we noticed that the recovery curves and hence the recovery timescales were nearly identical for droplets of small sizes ($R \lesssim 1.5 \mu\text{m}$), despite spanning nearly a twofold size range [Figs. 6(c) and 6(d)]. This observation aligns with the scaling behavior of bulk conversion-limited recovery (τ_{con}). If the recovery of these droplets were diffusion limited, whether by dilute-phase diffusion or dense-phase diffusion (of either species), we would expect nearly a fourfold difference in exchange timescale. For larger droplets, we see a quadratic scaling that arises above the lower-limit plateau set by the bulk conversion-limited timescale [Fig. 6(d)]. Upon fitting the data with the shifted quadratic function $\tau = a + bR^2$, we find the constants a and b are well constrained: $a = 145.0 \pm 5.6 \text{ s}$ and $b = 14.1 \pm 0.7 \mu\text{m}^{-2} \text{ s}$. The observed scalings were independently validated by a replicate experiment (Fig. S3 [51]).

Our theory suggests a plausible explanation for the origin of the observed plateau regime: the recovery is bulk conversion limited, i.e., $\tau_{\text{con}} = 1/(kc_{2,\text{den}}) = 145 \text{ s}$. In contrast to the detachment timescale of a nanostar from the network inferred here, the dissociation time of a single sticky end (6 bp) is typically much shorter, in the range $\tau_b = 0.01\text{--}10 \text{ s}$ [65–67]. The difference is likely explained by multivalency, which enhances the binding affinity [30,68] and thereby extends the effective lifetime of a nanostar far beyond that of a single sticky end. Interestingly, the dissociation time of a DNA origami dimer held together by three 6-bp stickers has been measured to be on the order of 100 s [69]. Given that physical parameters such as temperature and ionic strength can strongly impact the detachment timescales of both a single sticker and an entire nanostar [70], quantitative verification of the relevant timescales in our system requires future experiments, for example, via single-molecule tracking.

Nanostar droplets are porous, with a measured pore size comparable to the arm length. It has been reported that the partition coefficient for 70 kDa dextran (hydrodynamic radius 6 nm) is about 0.3–0.6 in such systems [64,71]. Given that our nanostars have a hydrodynamic radius (5–7 nm) similar to that of dextran, we expect the unbound species of

nanostars to partition in these nanostar droplets to a similar extent as the dextran. Assuming a partition coefficient of 0.5 and dilute-phase concentration of 1 μM [63,72], $c_{2,\text{den}} \approx 0.5 c_{2,\text{dil}} = 0.5 \mu\text{M}$, then the rate of nanostar attachment inside the condensate can be estimated as $k = 1/(ac_{2,\text{den}}) \approx 0.014 \mu\text{M}^{-1} \text{s}^{-1}$. This rate appears to be much lower than the reported on-rate for nanostars in dilute solution, which ranges from 0.1 to 1 $\mu\text{M}^{-1} \text{s}^{-1}$ [65,73,74]. The discrepancy likely arises because we have modeled the attachment flux as $kc_1^{\text{eq}}c_2$ in Eq. (1), implicitly assuming that every nanostar in the percolated network (species 1) can bind to freely diffusing species 2 nanostars. However, many nanostars in the network may already be in a fully bound state or spatially occluded and thus unavailable for binding, leading to an apparent smaller value of the attachment rate k . If nanostar attachment in the dense phase occurs with the same on-rate as in dilute solution, we can infer that more than 90% of nanostar sticky ends are not available for binding in these droplets.

The plateau in the τ versus R curve for small droplets also suggests that the interface-limited timescale τ_{int} is much longer than the bulk conversion-limited timescale τ_{con} , i.e., $\tau_{\text{int}}/\tau_{\text{con}} = Rc_{2,\text{den}}/(3\delta_{\text{eff}}c_{2,\text{dil}}) \gg 1$. Nanostar condensates have a surface tension of $\sigma \approx 1 \mu\text{N/m}$ [61], which can be used to estimate the interfacial width as $l \sim (k_{\text{B}}T/\sigma)^{1/2} = 60 \text{ nm}$ [64] and a corresponding effective width of $\delta_{\text{eff}} \approx l/2 = 30 \text{ nm}$ (see the Supplemental Material [51]). Thus, even for the smallest measured droplet ($R = 0.84 \mu\text{m}$), we have $\tau_{\text{int}}/\tau_{\text{con}} \approx 5$, indicating that τ_{int} is indeed much longer than τ_{con} for all droplets studied in these experiments. This also suggests that if no high-mobility species were present inside the nanostar droplets, the exchange timescale would be at least five times longer for these measured droplets, as predicted by Eq. (15).

Beyond the plateau regime, the negligible contribution of τ_{int} to the overall recovery time in Eq. (15) implies a reduced approximate expression for the recovery time:

$$\tau = \frac{1}{kc_{2,\text{den}}} + \left(\frac{c_{1,\text{den}}}{3c_{2,\text{dil}}D_{2,\text{dil}}} + \frac{1}{\pi^2 D_{\text{eff}}} \right) R^2. \quad (18)$$

While the first term originated from τ_{con} sets the plateau value a , the second term from τ_{dil} , $\tau_{1,\text{den}}$, and $\tau_{2,\text{den}}$ scales with R^2 and sets the prefactor $b = c_{1,\text{den}}/(3c_{2,\text{dil}}D_{2,\text{dil}}) + 1/(\pi^2 D_{\text{eff}})$. Taking $c_{1,\text{den}} = 200 \mu\text{M}$, $c_{2,\text{dil}} = 1 \mu\text{M}$, and $D_{2,\text{dil}} = 50 \mu\text{m}^2/\text{s}$ (from the Stokes-Einstein relation) [63,72], we find that $\tau_{\text{dil}} \ll \tau$ for all droplets, suggesting that recovery is not limited by dilute-phase diffusion but by dense-phase diffusion. We then estimate $D_{\text{eff}} \approx 8 \times 10^{-3} \mu\text{m}^2/\text{s}$ from the fitted value of b , which sets upper bounds for $D_{1,\text{den}}$ and $D_{2,\text{den}}$: $D_{1,\text{den}} \leq 8 \times 10^{-3} \mu\text{m}^2/\text{s}$ and $D_{2,\text{den}} \leq 3 \mu\text{m}^2/\text{s}$ based on Eq. (8), assuming $c_{2,\text{den}} = 0.5 \mu\text{M}$.

III. DISCUSSION

A hallmark of biomolecular condensates is their dynamic exchange of material with their surroundings, a feature often crucial to their function. In this work, we developed a minimal reaction-diffusion model to describe condensate exchange dynamics and explored how such dynamics can deviate from those of conventional liquid droplets due to the formation of a percolated network in the condensate. Our theory suggests

that, in the presence of two mobility states, material exchange can be accelerated via a pore-mediated pathway in which molecules pass through the pores of the meshwork and attach and detach directly in the condensate interior. Notably, this pathway leads to a parameter regime in which the exchange timescale becomes independent of condensate size, a prediction consistent with our FRAP measurements on DNA nanostar condensates.

In this study, we focused on the exchange dynamics of *in vitro* single-component condensates and approximated a condensate as a two-state system at equilibrium. However, molecules inside biological condensates, even in single-component systems, are likely to form a variable number of bonds rather than being simply fully bound or fully unbound. As a result, molecules inside condensates may exhibit a broad range of mobilities due to the multivalency of underlying interactions [49,50]. Moreover, the relative abundances of different mobility states can evolve over time, for example, during condensate aging [75]. Reducing such systems to a two-state description can, in principle, give rise to non-Markovian effects that our minimal model does not capture. A natural next step would be to incorporate a distribution of mobility states into the model. Nevertheless, a concise two-state model can help explore the physical principles underlying condensate exchange dynamics while remaining analytically tractable. Thus, our work serves as a first step toward addressing more complex dynamical problems in phase-separated systems.

The developed reaction-diffusion model can also be readily extended to describe multicomponent systems. Condensates in living cells are complex assemblies of distinct proteins and nucleic acids, and much theoretical effort has been dedicated to understanding thermodynamic principles in multicomponent systems with many coexisting phases [76–79]. However, less effort has been directed toward understanding the dynamics of such systems. Multicomponent condensates generally employ a scaffold-client framework [30,80]. While our model focused on the exchange dynamics of scaffold molecules, it can be generalized to describe client dynamics, where client molecules bind to and unbind from an equilibrated scaffold network, thus switching between low and high mobilities [81]. Since clients may bind not only to the percolated network (species 1) but also to unbound scaffold molecules (species 2), modeling client dynamics in this paradigm may require introducing a third species. In a similar spirit, the developed model can also be extended to predict the dynamics of condensates in the cell nucleus, such as transcriptional condensates and double-strand-break repair condensates, in which molecules diffusing within the condensate are also capable of binding to and unbinding from DNA [82–85].

The pore-mediated pathway involves molecules passing through the pores of the meshwork and attaching and detaching directly within the condensate. In which systems, and for which molecules, would this pathway be favored and experimentally measurable? Because diffusion within the connected network for bound molecules is generally much slower than for unbound molecules, this pathway is likely to be most relevant when the mesh size of the percolated network is comparable to or larger than the size of the molecule of interest, allowing a sufficient concentration of high-mobility species.

It has been reported that the mesh size (or correlation length) for *in vitro* reconstituted condensates of purified LAF-1 protein is about 6 nm [86], while for reconstituted coacervates composed of two oppositely charged intrinsically disordered proteins, histone H1 and prothymosin- α , it is 2.4–4.3 nm [87]. More generally, in a recent work, we established a quantitative relationship between mesh size and concentration, which suggests that condensates at physiologically relevant concentrations of 100–400 mg/ml exhibit mesh sizes decreasing from 8 to 3 nm [88]. Since protein sizes span the reported mesh-size range, the pore-mediated pathway may apply to the exchange dynamics of scaffolds in condensates with sufficiently low dense-phase concentrations and to the exchange dynamics of small clients more generally.

What are the possible biological implications of the pore-mediated pathway of condensate exchange dynamics? Although the bulk viscosity of condensates can be orders of magnitude higher than that of conventional oil droplets, small molecules may sneak through pores in the meshwork and enter the condensate interior via this pore-mediated pathway, potentially bypassing slow diffusion caused by such high viscosity. The accelerated material exchange and diffusion enabled by this pathway may therefore speed up biochemical reactions in metabolic condensates, as well as facilitate the response of condensates to changes in the intracellular environment. If so, could the porosity of condensates be evolutionarily selected to promote more efficient dynamics?

Fascinating soft matter systems in their own right, biomolecular condensates are also increasingly implicated in cellular physiology and disease [89,90]. We hope that our

work will motivate further theoretical and experimental investigations into the complex dynamics in multicomponent, multistate condensates, shedding light on their functional roles and paving the way for applications in condensate bio-engineering.

ACKNOWLEDGMENTS

We are grateful to Ned Wingreen, Frank Jülicher, Alexander Grosberg, Jeremy Schmit, Daniel Reich, and Omar Saleh for insightful discussions and constructive feedback on the manuscript. R.K. and Y.Z. were supported by a startup fund at Johns Hopkins University, NIH Award No. R35GM162296, and the Alfred P. Sloan Foundation through a Sloan Research Fellowship to Y.Z. (FG-2025-25076). E.K. acknowledges support from Sloan Foundation Grant No. 147170 to R.S. and NIH Grant No. T32GM080189. R.S. acknowledges support from Sloan Foundation Grant No. 147170, DOE BES Award No. DE-SC0010426, NSF Grant No. FET-2107246, ONR Award No. N00014-23-1-2868, and the Kent Gordon Croft Investment Management Faculty Scholar fund. This research was also supported in part by NSF Grant No. PHY-2309135 and the Gordon and Betty Moore Foundation Grant No. 2919.02 to the Kavli Institute for Theoretical Physics (KITP).

DATA AVAILABILITY

The data that support the findings of this article are openly available [91].

- [1] S. F. Banani, H. O. Lee, A. A. Hyman, and M. K. Rosen, Biomolecular condensates: Organizers of cellular biochemistry, *Nat. Rev. Mol. Cell Biol.* **18**, 285 (2017).
- [2] Y. Shin and C. P. Brangwynne, Liquid phase condensation in cell physiology and disease, *Science* **357**, eaaf4382 (2017).
- [3] I. I. Cisse, I. Izeddin, S. Z. Causse, L. Boudarene, A. Senecal, L. Muresan, C. Dugast-Darzacq, B. Hajj, M. Dahan, and X. Darzacq, Real-time dynamics of RNA polymerase II clustering in live human cells, *Science* **341**, 664 (2013).
- [4] B. R. Sabari, A. Dall'Agnese, A. Boija, I. A. Klein, E. L. Coffey, K. Shrinivas, B. J. Abraham, N. M. Hannett, A. V. Zamudio, J. C. Manteiga, *et al.*, Coactivator condensation at super-enhancers links phase separation and gene control, *Science* **361**, eaar3958 (2018).
- [5] T. Hirose, K. Ninomiya, S. Nakagawa, and T. Yamazaki, A guide to membraneless organelles and their various roles in gene regulation, *Nat. Rev. Mol. Cell Biol.* **24**, 288 (2023).
- [6] P. Li, S. Banjade, H.-C. Cheng, S. Kim, B. Chen, L. Guo, M. Llaguno, J. V. Hollingsworth, D. S. King, S. F. Banani, *et al.*, Phase transitions in the assembly of multivalent signalling proteins, *Nature (London)* **483**, 336 (2012).
- [7] H. Wu, Higher-order assemblies in a new paradigm of signal transduction, *Cell* **153**, 287 (2013).
- [8] X. Su, J. A. Ditlev, E. Hui, W. Xing, S. Banjade, J. Okrut, D. S. King, J. Taunton, M. K. Rosen, and R. D. Vale, Phase separation of signaling molecules promotes T cell receptor signal transduction, *Science* **352**, 595 (2016).
- [9] A. Molliex, J. Temirov, J. Lee, M. Coughlin, A. P. Kanagaraj, H. J. Kim, T. Mittag, and J. P. Taylor, Phase separation by low complexity domains promotes stress granule assembly and drives pathological fibrillization, *Cell* **163**, 123 (2015).
- [10] A. Patel, H. O. Lee, L. Jawerth, S. Maharana, M. Jahnel, M. Y. Hein, S. Stoyanov, J. Mahamid, S. Saha, T. M. Franzmann, *et al.*, A liquid-to-solid phase transition of the ALS protein FUS accelerated by disease mutation, *Cell* **162**, 1066 (2015).
- [11] G. Gaglia, R. Rashid, C. Yapp, G. N. Joshi, C. G. Li, S. L. Lindquist, K. A. Sarosiek, L. Whitesell, P. K. Sorger, and S. Santagata, HSF1 phase transition mediates stress adaptation and cell fate decisions, *Nat. Cell Biol.* **22**, 151 (2020).
- [12] S. Wegmann, B. Eftekharzadeh, K. Tepper, K. M. Zoltowska, R. E. Bennett, S. Dujardin, P. R. Laskowski, D. MacKenzie, T. Kamath, C. Commins, *et al.*, Tau protein liquid-liquid phase separation can initiate tau aggregation, *EMBO J* **37**, EMBJ201798049 (2018).
- [13] S. Alberti and A. A. Hyman, Biomolecular condensates at the nexus of cellular stress, protein aggregation disease and ageing, *Nat. Rev. Mol. Cell Biol.* **22**, 196 (2021).
- [14] J. Nam and Y. Gwon, Neuronal biomolecular condensates and their implications in neurodegenerative diseases, *Front. Aging Neurosci.* **15**, 1145420 (2023).
- [15] S. Spann, M. Tereshchenko, G. J. Mastromarco, S. J. Ihn, and H. O. Lee, Biomolecular condensates in neurodegeneration and cancer, *Traffic* **20**, 890 (2019).

- [16] A. Boija, I. A. Klein, and R. A. Young, Biomolecular condensates and cancer, *Cancer Cell* **39**, 174 (2021).
- [17] K. Taniue and N. Akimitsu, Aberrant phase separation and cancer, *FEBS J.* **289**, 17 (2022).
- [18] X. Tong, R. Tang, J. Xu, W. Wang, Y. Zhao, X. Yu, and S. Shi, Liquid–liquid phase separation in tumor biology, *Signal Transduction Targeted Ther.* **7**, 221 (2022).
- [19] S. An, R. Kumar, E. D. Sheets, and S. J. Benkovic, Reversible compartmentalization of de novo purine biosynthetic complexes in living cells, *Science* **320**, 103 (2008).
- [20] A. M. Pedley, J. P. Boylan, C. Y. Chan, E. L. Kennedy, M. Kyoung, and S. J. Benkovic, Purine biosynthetic enzymes assemble into liquid-like condensates dependent on the activity of chaperone protein HSP90, *J. Biol. Chem.* **298**, 101845 (2022).
- [21] M. Prouteau and R. Loewith, Regulation of cellular metabolism through phase separation of enzymes, *Biomolecules* **8**, 160 (2018).
- [22] H. Wang, F. M. Kelley, D. Milovanovic, B. S. Schuster, and Z. Shi, Surface tension and viscosity of protein condensates quantified by micropipette aspiration, *Biophys. Rep.* **1**, 100011 (2021).
- [23] I. Lifshitz and V. Slyozov, The kinetics of precipitation from supersaturated solid solutions, *J. Phys. Chem. Solids* **19**, 35 (1961).
- [24] C. Wagner, Theorie der Alterung von Niederschlägen durch Umlösen (Ostwald-Reifung), *Z. Elektrochem.* **65**, 581 (1961).
- [25] M.-M. Poo and R. A. Cone, Lateral diffusion of rhodopsin in *Necturus* rods, *Exp. Eye Res.* **17**, 503 (1973).
- [26] P. Liebman and G. Entine, Lateral diffusion of visual pigment in photoreceptor disk membranes, *Science* **185**, 457 (1974).
- [27] J. Lippincott-Schwartz, E. Snapp, and A. Kenworthy, Studying protein dynamics in living cells, *Nat. Rev. Mol. Cell Biol.* **2**, 444 (2001).
- [28] F. Pincet, V. Adrien, R. Yang, J. Delacotte, J. E. Rothman, W. Urbach, and D. Tareste, FRAP to characterize molecular diffusion and interaction in various membrane environments, *PLoS One* **11**, e0158457 (2016).
- [29] K. A. Burke, A. M. Janke, C. L. Rhine, and N. L. Fawzi, Residue-by-residue view of *in vitro* FUS granules that bind the C-terminal domain of RNA polymerase II, *Mol. Cell* **60**, 231 (2015).
- [30] S. F. Banani, A. M. Rice, W. B. Peeples, Y. Lin, S. Jain, R. Parker, and M. K. Rosen, Compositional control of phase-separated cellular bodies, *Cell* **166**, 651 (2016).
- [31] S. Jain, J. R. Wheeler, R. W. Walters, A. Agrawal, A. Barsic, and R. Parker, ATPase-modulated stress granules contain a diverse proteome and substructure, *Cell* **164**, 487 (2016).
- [32] W. M. J. Aumiller, F. Pir Cakmak, B. W. Davis, and C. D. Keating, RNA-based coacervates as a model for membraneless organelles: Formation, properties, and interfacial liposome assembly, *Langmuir* **32**, 10042 (2016).
- [33] N. O. Taylor, M.-T. Wei, H. A. Stone, and C. P. Brangwynne, Quantifying dynamics in phase-separated condensates using fluorescence recovery after photobleaching, *Biophys. J.* **117**, 1285 (2019).
- [34] L. Hubatsch, L. M. Jawerth, C. Love, J. Bauermann, T. D. Tang, S. Bo, A. A. Hyman, and C. A. Weber, Quantitative theory for the diffusive dynamics of liquid condensates, *eLife* **10**, e68620 (2021).
- [35] S. Bo, L. Hubatsch, J. Bauermann, C. A. Weber, and F. Jülicher, Stochastic dynamics of single molecules across phase boundaries, *Phys. Rev. Res.* **3**, 043150 (2021).
- [36] F. Muzzopappa, J. Hummert, M. Anfossi, S. A. Tashev, D.-P. Herten, and F. Erdel, Detecting and quantifying liquid–liquid phase separation in living cells by model-free calibrated half-bleaching, *Nat. Commun.* **13**, 7787 (2022).
- [37] A. W. Folkmann, A. Putnam, C. F. Lee, and G. Seydoux, Regulation of biomolecular condensates by interfacial protein clusters, *Science* **373**, 1218 (2021).
- [38] C. F. Lee, Scaling law and universal drop size distribution of coarsening in conversion-limited phase separation, *Phys. Rev. Res.* **3**, 043081 (2021).
- [39] H. T. Nguyen, N. Hori, and D. Thirumalai, Condensates in RNA repeat sequences are heterogeneously organized and exhibit reptation dynamics, *Nat. Chem.* **14**, 775 (2022).
- [40] Y. Zhang, A. G. Pyo, R. Kliegman, Y. Jiang, C. P. Brangwynne, H. A. Stone, and N. S. Wingreen, The exchange dynamics of biomolecular condensates, *eLife* **12**, RP91680 (2024).
- [41] L. Hubatsch, S. Bo, T. S. Harmon, A. A. Hyman, C. A. Weber, and F. Jülicher, Transport kinetics across interfaces between coexisting liquid phases, *eLife* **14**, RP106842 (2025).
- [42] G. L. Dignon, R. B. Best, and J. Mittal, Biomolecular phase separation: From molecular driving forces to macroscopic properties, *Annu. Rev. Phys. Chem.* **71**, 53 (2020).
- [43] J. A. Joseph, A. Reinhardt, A. Aguirre, P. Y. Chew, K. O. Russell, J. R. Espinosa, A. Garaizar, and R. Collepardo-Guevara, Physics-driven coarse-grained model for biomolecular phase separation with near-quantitative accuracy, *Nat. Comput. Sci.* **1**, 732 (2021).
- [44] A. N. Semenov and M. Rubinstein, Thermoreversible gelation in solutions of associative polymers. 1. Statics, *Macromolecules* **31**, 1373 (1998).
- [45] J. Wang, J.-M. Choi, A. S. Holehouse, H. O. Lee, X. Zhang, M. Jahnel, S. Maharana, R. Lemaitre, A. Pozniakovsky, D. Drechsel, *et al.*, A molecular grammar governing the driving forces for phase separation of prion-like RNA binding proteins, *Cell* **174**, 688 (2018).
- [46] G. M. Ginell and A. S. Holehouse, An introduction to the stickers-and-spacers framework as applied to biomolecular condensates, in *Phase-Separated Biomolecular Condensates: Methods and Protocols*, edited by H.-X. Zhou, J.-H. Spille, and P. R. Banerjee (Springer US, New York, 2023), pp. 95–116.
- [47] J.-M. Choi, A. S. Holehouse, and R. V. Pappu, Physical principles underlying the complex biology of intracellular phase transitions, *Annu. Rev. Biophys.* **49**, 107 (2020).
- [48] T. Mittag and R. V. Pappu, A conceptual framework for understanding phase separation and addressing open questions and challenges, *Mol. Cell* **82**, 2201 (2022).
- [49] Z. Shen, B. Jia, Y. Xu, J. Wessén, T. Pal, H. S. Chan, S. Du, and M. Zhang, Biological condensates form percolated networks with molecular motion properties distinctly different from dilute solutions, *eLife* **12**, e81907 (2023).
- [50] T. Wu, M. R. King, Y. Qiu, M. Farag, R. V. Pappu, and M. D. Lew, Single-fluorogen imaging reveals distinct environmental and structural features of biomolecular condensates, *Nat. Phys.* **21**, 778 (2025).
- [51] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/h9cl-j5x5> for additional information about the theory,

- numerical simulations, and experiments, which includes Refs. [52,53,56,60,63,92].
- [52] J. Crank, *The Mathematics of Diffusion*, Oxford Science Publications (Clarendon Press, Oxford, UK, 1979).
 - [53] B. L. Sprague, R. L. Pego, D. A. Stavreva, and J. G. McNally, Analysis of binding reactions by fluorescence recovery after photobleaching, *Biophys. J.* **86**, 3473 (2004).
 - [54] J. W. Cahn and J. E. Hilliard, Free energy of a nonuniform system. I. Interfacial free energy, *J. Chem. Phys.* **28**, 258 (1958).
 - [55] R. Skeel and M. Berzins, A method for the spatial discretization of parabolic equations in one space variable, *SIAM J. Sci. Stat. Comput.* **11**, 1 (1990).
 - [56] M. Branch, T. Coleman, and Y. Li, A subspace, interior, and conjugate gradient method for large-scale bound-constrained minimization problems, *SIAM J. Sci. Comput.* **21**, 1 (1999).
 - [57] P. M. McCall, K. Kim, A. Shevchenko, M. Ruer-Gruß, J. Peychl, J. Guck, A. Shevchenko, A. A. Hyman, and J. Brugués, A label-free method for measuring the composition of multicomponent biomolecular condensates, *Nat. Chem.* **17**, 1891 (2025).
 - [58] L. Jawerth, E. Fischer-Friedrich, S. Saha, J. Wang, T. Franzmann, X. Zhang, J. Sachweh, M. Ruer, M. Ijavi, S. Saha, *et al.*, Protein condensates as aging Maxwell fluids, *Science* **370**, 1317 (2020).
 - [59] S. Biffi, R. Cerbino, F. Bomboi, E. M. Paraboschi, R. Asselta, F. Sciortino, and T. Bellini, Phase behavior and critical activated dynamics of limited-valence DNA nanostars, *Proc. Natl. Acad. Sci. USA* **110**, 15633 (2013).
 - [60] Y. Sato, T. Sakamoto, and M. Takinoue, Sequence-based engineering of dynamic functions of micrometer-sized DNA droplets, *Sci. Adv.* **6**, eaba3471 (2020).
 - [61] B.-j. Jeon, D. T. Nguyen, G. R. Abraham, N. Conrad, D. K. Fygenson, and O. A. Saleh, Salt-dependent properties of a coacervate-like, self-assembled DNA liquid, *Soft Matter* **14**, 7009 (2018).
 - [62] S. Agarwal, D. Osmanovic, M. Dizani, M. A. Klocke, and E. Franco, Dynamic control of DNA condensation, *Nat. Commun.* **15**, 1915 (2024).
 - [63] E. Kengmana, E. Ornelas-Gatdula, K.-L. Chen, and R. Schulman, Spatial control over reactions via localized transcription within membraneless DNA nanostar droplets, *J. Am. Chem. Soc.* **146**, 32942 (2024).
 - [64] D. T. Nguyen, B.-j. Jeon, G. R. Abraham, and O. A. Saleh, Length-dependence and spatial structure of DNA partitioning into a DNA liquid, *Langmuir* **35**, 14849 (2019).
 - [65] N. F. Dupuis, E. D. Holmstrom, and D. J. Nesbitt, Single-molecule kinetics reveal cation-promoted DNA duplex formation through ordering of single-stranded helices, *Biophys. J.* **105**, 756 (2013).
 - [66] N. Conrad, T. Kennedy, D. K. Fygenson, and O. A. Saleh, Increasing valence pushes DNA nanostar networks to the isostatic point, *Proc. Natl. Acad. Sci. USA* **116**, 7238 (2019).
 - [67] G. Palombo, S. Weir, D. Michieletto, and Y. A. Gutiérrez Fosado, Topological linking determines elasticity in limited valence networks, *Nat. Mater.* **24**, 454 (2025).
 - [68] K. Bhandari, M. A. Cotten, J. Kim, M. K. Rosen, and J. D. Schmit, Structure–function properties in disordered condensates, *J. Phys. Chem. B* **125**, 467 (2021).
 - [69] J. Zenk, C. Tuntivate, and R. Schulman, Kinetics and thermodynamics of Watson–Crick base pairing driven DNA origami dimerization, *J. Am. Chem. Soc.* **138**, 3346 (2016).
 - [70] F. Bomboi, S. Biffi, R. Cerbino, T. Bellini, F. Bordi, and F. Sciortino, Equilibrium gels of trivalent DNA-nanostars: Effect of the ionic strength on the dynamics, *Eur. Phys. J. E* **38**, 64 (2015).
 - [71] K. Skipper and S. F. J. Wickham, DNA nanostars that self-assemble into core-shell condensate microdroplets, *Nanoscale Horiz.* **11**, 1320 (2026).
 - [72] S. Do, C. Lee, T. Lee, D.-N. Kim, and Y. Shin, Engineering DNA-based synthetic condensates with programmable material properties, compositions, and functionalities, *Sci. Adv.* **8**, eabj1771 (2022).
 - [73] D. Y. Zhang and E. Winfree, Control of DNA strand displacement kinetics using toehold exchange, *J. Am. Chem. Soc.* **131**, 17303 (2009).
 - [74] S. Jiang, H. Yan, and Y. Liu, Kinetics of DNA tile dimerization, *ACS Nano* **8**, 5826 (2014).
 - [75] R. Takaki, L. Jawerth, M. Popović, and F. Jülicher, Theory of rheology and aging of protein condensates, *PRX Life* **1**, 013006 (2023).
 - [76] W. M. Jacobs and D. Frenkel, Phase transitions in biological systems with many components, *Biophys. J.* **112**, 683 (2017).
 - [77] D. Zwicker and L. Laan, Evolved interactions stabilize many coexisting phases in multicomponent liquids, *Proc. Natl. Acad. Sci. USA* **119**, e2201250119 (2022).
 - [78] W. M. Jacobs, Theory and simulation of multiphase coexistence in biomolecular mixtures, *J. Chem. Theory Comput.* **19**, 3429 (2023).
 - [79] Y. Qiang, C. Luo, and D. Zwicker, Scaling laws for phase coexistence in multicomponent mixtures, *Phys. Rev. Res.* **7**, 043008 (2025).
 - [80] J. A. Ditlev, L. B. Case, and M. K. Rosen, Who’s in and who’s out—compositional control of biomolecular condensates, *J. Mol. Biol.* **430**, 4666 (2018).
 - [81] J. D. Schmit and T. C. T. Michaels, Physical limits to acceleration of enzymatic reactions inside phase-separated compartments, *Phys. Rev. E* **109**, 064401 (2024).
 - [82] S. Kent, K. Brown, C.-h. Yang, N. Alsaihati, C. Tian, H. Wang, and X. Ren, Phase-separated transcriptional condensates accelerate target-search process revealed by live-cell single-molecule imaging, *Cell Rep.* **33**, 108248 (2020).
 - [83] J. Miné-Hattab, M. Heltberg, M. Villemeur, C. Guedj, T. Mora, A. M. Walczak, M. Dahan, and A. Taddei, Single molecule microscopy reveals key physical features of repair foci in living cells, *eLife* **10**, e60577 (2021).
 - [84] M. L. Heltberg, J. Miné-Hattab, A. Taddei, A. M. Walczak, and T. Mora, Physical observables to determine the nature of membrane-less cellular sub-compartments, *eLife* **10**, e69181 (2021).
 - [85] S. Hao, Y. J. Lee, N. B. Goldfajn, E. Flores, J. Liang, H. Fuehrer, J. Demmerle, J. Lippincott-Schwartz, Z. Liu, S. Sukenik, *et al.*, YAP condensates are highly organized hubs, *iScience* **27**, 109927 (2024).
 - [86] M.-T. Wei, S. Elbaum-Garfinkle, A. S. Holehouse, C. C.-H. Chen, M. Feric, C. B. Arnold, R. D. Priestley, R. V. Pappu, and C. P. Brangwynne, Phase behaviour of disordered proteins underlying low density and high permeability of liquid organelles, *Nat. Chem.* **9**, 1118 (2017).

- [87] N. Galvanetto, M. T. Ivanović, A. Chowdhury, A. Sottini, M. F. Nüesch, D. Nettels, R. B. Best, and B. Schuler, Extreme dynamics in a biomolecular condensate, *Nature (London)* **619**, 876 (2023).
- [88] V. Grigorev, N. S. Wingreen, and Y. Zhang, Conformational entropy of intrinsically disordered proteins bars intruders from biomolecular condensates, *PRX Life* **3**, 013011 (2025).
- [89] B. Wang, L. Zhang, T. Dai, Z. Qin, H. Lu, L. Zhang, and F. Zhou, Liquid–liquid phase separation in human health and diseases, *Signal Transduction Targeted Ther.* **6**, 290 (2021).
- [90] D. M. Mitrea, M. Mittasch, B. F. Gomes, I. A. Klein, and M. A. Murcko, Modulating biomolecular condensates: A novel approach to drug discovery, *Nat. Rev. Drug Discov.* **21**, 841 (2022).
- [91] R. Kliegman, E. Kengmana, R. Schulman, and Y. Zhang, Dissecting rate-limiting processes in biomolecular condensate exchange dynamics, [dataset and code] Johns Hopkins Research Data Repository, 2026, <https://doi.org/10.7281/T1WY5RSP>.
- [92] N. Otsu, A threshold selection method from gray-level histograms, *IEEE Trans. Syst. Man. Cybern.* **9**, 62 (1979).